



wwPDB X-ray Structure Validation Summary Report ⓘ

Apr 10, 2024 – 01:17 pm BST

PDB ID : 8RUD
Title : Crystal structure of Rhizobium etli L-asparaginase ReAV K138A mutant
Authors : Pokrywka, K.; Grzechowiak, M.; Sliwiak, J.; Worsztynowicz, P.; Loch, J.I.;
Ruszkowski, M.; Gilski, M.; Jaskolski, M.
Deposited on : 2024-01-30
Resolution : 2.10 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	373	 84% 9% 6%
1	B	373	 82% 11% 6%
1	C	373	 86% 8% 6%
1	D	373	 81% 11% 6%

2 Entry composition i

There are 6 unique types of molecules in this entry. The entry contains 11349 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called L-asparaginase II protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	350	2650	1639	489	500	22	0	10	0
1	B	349	2607	1609	480	497	21	0	5	0
1	C	349	2635	1628	486	497	24	0	10	0
1	D	349	2629	1626	482	498	23	0	9	0

There are 28 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	GLY	-	expression tag	UNP Q2K0Z2
A	-4	ILE	-	expression tag	UNP Q2K0Z2
A	-3	ASP	-	expression tag	UNP Q2K0Z2
A	-2	PRO	-	expression tag	UNP Q2K0Z2
A	-1	PHE	-	expression tag	UNP Q2K0Z2
A	0	THR	-	expression tag	UNP Q2K0Z2
A	138	ALA	LYS	engineered mutation	UNP Q2K0Z2
B	-5	GLY	-	expression tag	UNP Q2K0Z2
B	-4	ILE	-	expression tag	UNP Q2K0Z2
B	-3	ASP	-	expression tag	UNP Q2K0Z2
B	-2	PRO	-	expression tag	UNP Q2K0Z2
B	-1	PHE	-	expression tag	UNP Q2K0Z2
B	0	THR	-	expression tag	UNP Q2K0Z2
B	138	ALA	LYS	engineered mutation	UNP Q2K0Z2
C	-5	GLY	-	expression tag	UNP Q2K0Z2
C	-4	ILE	-	expression tag	UNP Q2K0Z2
C	-3	ASP	-	expression tag	UNP Q2K0Z2
C	-2	PRO	-	expression tag	UNP Q2K0Z2
C	-1	PHE	-	expression tag	UNP Q2K0Z2
C	0	THR	-	expression tag	UNP Q2K0Z2
C	138	ALA	LYS	engineered mutation	UNP Q2K0Z2

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	-5	GLY	-	expression tag	UNP Q2K0Z2
D	-4	ILE	-	expression tag	UNP Q2K0Z2
D	-3	ASP	-	expression tag	UNP Q2K0Z2
D	-2	PRO	-	expression tag	UNP Q2K0Z2
D	-1	PHE	-	expression tag	UNP Q2K0Z2
D	0	THR	-	expression tag	UNP Q2K0Z2
D	138	ALA	LYS	engineered mutation	UNP Q2K0Z2

- Molecule 2 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total C O 4 2 2	0	0
2	C	1	Total C O 4 2 2	0	0
2	C	1	Total C O 4 2 2	0	0
2	C	1	Total C O 4 2 2	0	0
2	C	1	Total C O 4 2 2	0	0
2	C	1	Total C O 4 2 2	0	0
2	C	1	Total C O 4 2 2	0	0
2	C	1	Total C O 4 2 2	0	0
2	D	1	Total C O 4 2 2	0	0
2	D	1	Total C O 4 2 2	0	0
2	D	1	Total C O 4 2 2	0	0
2	D	1	Total C O 4 2 2	0	0
2	D	1	Total C O 4 2 2	0	0
2	D	1	Total C O 4 2 2	0	0

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Cl 1 1	0	0

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total C O 6 3 3	0	0
4	B	1	Total C O 6 3 3	0	0

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	D	1	Total O S 5 4 1	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	169	Total 169	O 169	0	0
6	B	192	Total 192	O 192	0	0
6	C	176	Total 176	O 176	0	0
6	D	185	Total 185	O 185	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: L-asparaginase II protein




- Molecule 1: L-asparaginase II protein



- Molecule 1: L-asparaginase II protein



- Molecule 1: L-asparaginase II protein





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	78.07Å 91.39Å 114.55Å 90.00° 96.90° 90.00°	Depositor
Resolution (Å)	56.86 – 2.10 56.86 – 2.10	Depositor EDS
% Data completeness (in resolution range)	97.8 (56.86-2.10) 97.8 (56.86-2.10)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.39 (at 2.10Å)	Xtrriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.206 , 0.262 0.204 , 0.260	Depositor DCC
R_{free} test set	999 reflections (1.10%)	wwPDB-VP
Wilson B-factor (Å ²)	23.1	Xtrriage
Anisotropy	0.306	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 43.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	11349	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 36.64 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 4.8430e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, CL, GOL, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.48	0/2725	0.69	0/3681
1	B	0.49	0/2666	0.70	0/3602
1	C	0.51	0/2709	0.70	0/3657
1	D	0.49	0/2700	0.69	0/3646
All	All	0.49	0/10800	0.69	0/14586

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2650	0	2641	20	0
1	B	2607	0	2580	32	0
1	C	2635	0	2625	21	0
1	D	2629	0	2619	31	0
2	A	8	0	12	4	0
2	B	28	0	42	5	0
2	C	28	0	42	6	0
2	D	24	0	36	7	0
3	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	12	0	16	2	0
5	D	5	0	0	0	0
6	A	169	0	0	2	0
6	B	192	0	0	8	0
6	C	176	0	0	4	0
6	D	185	0	0	6	0
All	All	11349	0	10613	106	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 106 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:189[A]:CYS:SG	6:C:644:HOH:O	2.22	0.96
1:B:71:ASP:HB2	2:B:404:EDO:H11	1.65	0.79
1:B:130:ALA:HB2	4:B:403:GOL:H11	1.69	0.75
1:D:224:ARG:HH11	2:D:406:EDO:H21	1.52	0.75
1:C:71:ASP:HB2	2:C:403:EDO:H22	1.69	0.74

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	358/373 (96%)	349 (98%)	9 (2%)	0	100	100
1	B	352/373 (94%)	341 (97%)	11 (3%)	0	100	100
1	C	357/373 (96%)	349 (98%)	8 (2%)	0	100	100
1	D	356/373 (95%)	345 (97%)	11 (3%)	0	100	100
All	All	1423/1492 (95%)	1384 (97%)	39 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	273/282 (97%)	262 (96%)	11 (4%)	31	32
1	B	267/282 (95%)	258 (97%)	9 (3%)	37	39
1	C	272/282 (96%)	267 (98%)	5 (2%)	59	65
1	D	271/282 (96%)	263 (97%)	8 (3%)	41	44
All	All	1083/1128 (96%)	1050 (97%)	33 (3%)	42	44

5 of 33 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	131	VAL
1	D	147	ARG
1	D	320	ASP
1	B	32	ARG
1	B	21	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry

Of 26 ligands modelled in this entry, 1 is monoatomic - leaving 25 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	EDO	A	402	-	3,3,3	0.54	0	2,2,2	0.31	0
2	EDO	C	406	-	3,3,3	0.42	0	2,2,2	0.80	0
2	EDO	A	401	-	3,3,3	0.46	0	2,2,2	0.70	0
2	EDO	D	405	-	3,3,3	0.49	0	2,2,2	0.80	0
4	GOL	B	403	-	5,5,5	1.22	0	5,5,5	0.82	0
2	EDO	C	402	-	3,3,3	0.35	0	2,2,2	0.87	0
2	EDO	D	406	-	3,3,3	0.38	0	2,2,2	0.73	0
5	SO4	D	407	-	4,4,4	0.19	0	6,6,6	0.21	0
2	EDO	D	401	-	3,3,3	0.39	0	2,2,2	0.40	0
2	EDO	D	402	-	3,3,3	0.39	0	2,2,2	0.61	0
4	GOL	B	409	-	5,5,5	1.07	0	5,5,5	0.78	0
2	EDO	B	405	-	3,3,3	0.66	0	2,2,2	0.37	0
2	EDO	B	408	-	3,3,3	0.60	0	2,2,2	0.16	0
2	EDO	C	404	-	3,3,3	0.85	0	2,2,2	0.43	0
2	EDO	B	407	-	3,3,3	0.53	0	2,2,2	0.54	0
2	EDO	D	404	-	3,3,3	0.62	0	2,2,2	0.07	0
2	EDO	B	401	-	3,3,3	0.48	0	2,2,2	0.41	0
2	EDO	B	404	-	3,3,3	0.59	0	2,2,2	0.38	0
2	EDO	C	407	-	3,3,3	0.55	0	2,2,2	0.26	0
2	EDO	D	403	-	3,3,3	0.55	0	2,2,2	0.34	0
2	EDO	B	406	-	3,3,3	0.52	0	2,2,2	0.45	0
2	EDO	C	403	-	3,3,3	0.64	0	2,2,2	0.28	0
2	EDO	C	405	-	3,3,3	0.68	0	2,2,2	0.17	0
2	EDO	C	401	-	3,3,3	0.58	0	2,2,2	0.56	0
2	EDO	B	402	-	3,3,3	0.53	0	2,2,2	0.39	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	A	402	-	-	0/1/1/1	-
2	EDO	C	406	-	-	1/1/1/1	-
2	EDO	A	401	-	-	1/1/1/1	-
2	EDO	D	405	-	-	1/1/1/1	-
4	GOL	B	403	-	-	0/4/4/4	-
2	EDO	C	402	-	-	0/1/1/1	-
2	EDO	D	406	-	-	1/1/1/1	-
2	EDO	D	401	-	-	0/1/1/1	-
2	EDO	D	402	-	-	0/1/1/1	-
4	GOL	B	409	-	-	2/4/4/4	-
2	EDO	B	405	-	-	1/1/1/1	-
2	EDO	B	408	-	-	1/1/1/1	-
2	EDO	C	404	-	-	0/1/1/1	-
2	EDO	B	407	-	-	1/1/1/1	-
2	EDO	D	404	-	-	0/1/1/1	-
2	EDO	B	401	-	-	1/1/1/1	-
2	EDO	B	404	-	-	1/1/1/1	-
2	EDO	C	407	-	-	0/1/1/1	-
2	EDO	D	403	-	-	1/1/1/1	-
2	EDO	B	406	-	-	1/1/1/1	-
2	EDO	C	403	-	-	1/1/1/1	-
2	EDO	C	405	-	-	0/1/1/1	-
2	EDO	C	401	-	-	0/1/1/1	-
2	EDO	B	402	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 15 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	408	EDO	O1-C1-C2-O2
4	B	409	GOL	C1-C2-C3-O3
2	B	405	EDO	O1-C1-C2-O2
2	D	405	EDO	O1-C1-C2-O2
4	B	409	GOL	O2-C2-C3-O3

There are no ring outliers.

17 monomers are involved in 24 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	402	EDO	2	0
2	A	401	EDO	2	0
2	D	405	EDO	1	0
4	B	403	GOL	2	0
2	C	402	EDO	1	0
2	D	406	EDO	2	0
2	D	402	EDO	1	0
2	B	405	EDO	1	0
2	B	408	EDO	1	0
2	C	404	EDO	2	0
2	D	404	EDO	1	0
2	B	404	EDO	2	0
2	D	403	EDO	2	0
2	B	406	EDO	1	0
2	C	403	EDO	1	0
2	C	405	EDO	1	0
2	C	401	EDO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	350/373 (93%)	-0.01	5 (1%) 75 78	16, 23, 42, 69	0
1	B	349/373 (93%)	-0.06	1 (0%) 94 94	16, 21, 36, 53	0
1	C	349/373 (93%)	-0.02	2 (0%) 89 91	19, 25, 41, 60	0
1	D	349/373 (93%)	-0.03	3 (0%) 84 86	18, 23, 37, 63	0
All	All	1397/1492 (93%)	-0.03	11 (0%) 86 88	16, 23, 40, 69	0

The worst 5 of 11 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	218	GLY	3.8
1	A	353	SER	3.3
1	A	220	GLY	3.0
1	A	216	ASP	2.9
1	D	5	GLU	2.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	EDO	C	407	4/4	0.70	0.30	34,42,45,50	0
2	EDO	C	405	4/4	0.74	0.20	31,31,33,35	0
2	EDO	C	404	4/4	0.79	0.25	25,30,32,33	0
2	EDO	B	407	4/4	0.81	0.19	30,36,37,39	0
2	EDO	B	405	4/4	0.81	0.25	25,26,29,33	0
2	EDO	A	402	4/4	0.84	0.21	30,37,37,38	0
2	EDO	A	401	4/4	0.85	0.17	29,30,32,35	0
2	EDO	D	404	4/4	0.85	0.18	27,33,33,33	0
2	EDO	B	404	4/4	0.88	0.14	32,32,32,35	0
2	EDO	C	401	4/4	0.88	0.15	21,32,34,34	0
2	EDO	D	406	4/4	0.88	0.15	28,30,33,38	0
4	GOL	B	403	6/6	0.88	0.37	26,27,30,30	0
2	EDO	D	402	4/4	0.89	0.32	24,30,32,45	0
2	EDO	C	403	4/4	0.89	0.22	32,34,34,34	0
5	SO4	D	407	5/5	0.89	0.18	24,26,34,34	5
2	EDO	D	405	4/4	0.90	0.25	34,38,39,42	0
2	EDO	B	408	4/4	0.90	0.12	33,33,34,36	0
2	EDO	D	403	4/4	0.90	0.30	25,30,33,33	0
2	EDO	C	402	4/4	0.90	0.13	35,38,40,45	0
4	GOL	B	409	6/6	0.91	0.24	22,28,34,42	0
2	EDO	B	406	4/4	0.91	0.32	26,30,32,37	0
2	EDO	C	406	4/4	0.93	0.15	34,41,44,44	0
2	EDO	B	401	4/4	0.93	0.23	30,32,33,39	0
2	EDO	B	402	4/4	0.95	0.09	25,26,28,33	0
3	CL	A	403	1/1	0.96	0.10	36,36,36,36	0
2	EDO	D	401	4/4	0.97	0.11	25,30,30,33	0

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.