



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 30, 2023 – 10:14 AM EST

PDB ID : 4RWD  
Title : XFEL structure of the human delta opioid receptor in complex with a bifunctional peptide  
Authors : Fenalti, G.; Zatsepin, N.A.; Betti, C.; Giguere, P.; Han, G.W.; Ishchenko, A.; Liu, W.; Guillemy, K.; Zhang, H.; James, D.; Wang, D.; Weierstall, U.; Spence, J.C.H.; Boutet, S.; Messerschmidt, M.; Williams, G.J.; Gati, C.; Yefanov, O.M.; White, T.A.; Oberthuer, D.; Metz, M.; Yoon, C.H.; Barty, A.; Chapman, H.N.; Basu, S.; Coe, J.; Conrad, C.E.; Fromme, R.; Fromme, P.; Tourwe, D.; Schiller, P.W.; Roth, B.L.; Ballet, S.; Katritch, V.; Stevens, R.C.; Cherezov, V.; GPCR Network (GPCR)  
Deposited on : 2014-12-02  
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)



The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	411	
1	B	411	

*Continued on next page...*

Refmac : 5.8.0158  
 CCP4 : 7.0.044 (Gargrove)  
 Ideal geometry (proteins) : Engh & Huber (2001)  
 Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
 Validation Pipeline (wwPDB-VP) : 2.36

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	G	5	
2	H	5	

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 6326 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Soluble cytochrome b562,Delta-type opioid receptor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	401	3030	1981	489	537	23	0	0	0
1	B	400	3035	1980	490	542	23	0	0	0


There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	996	GLY	-	expression tag	UNP P0ABE7
A	997	GLY	-	expression tag	UNP P0ABE7
A	998	THR	-	expression tag	UNP P0ABE7
A	999	THR	-	expression tag	UNP P0ABE7
A	1000	MET	-	expression tag	UNP P0ABE7
A	1007	TRP	MET	engineered mutation	UNP P0ABE7
A	1102	ILE	HIS	engineered mutation	UNP P0ABE7
A	1106	LEU	ARG	engineered mutation	UNP P0ABE7
B	996	GLY	-	expression tag	UNP P0ABE7
B	997	GLY	-	expression tag	UNP P0ABE7
B	998	THR	-	expression tag	UNP P0ABE7
B	999	THR	-	expression tag	UNP P0ABE7
B	1000	MET	-	expression tag	UNP P0ABE7
B	1007	TRP	MET	engineered mutation	UNP P0ABE7
B	1102	ILE	HIS	engineered mutation	UNP P0ABE7
B	1106	LEU	ARG	engineered mutation	UNP P0ABE7

- Molecule 2 is a protein called bifunctional peptide.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	G	5	49	39	5	5	0	0	1
2	H	5	49	39	5	5	0	0	1

- Molecule 3 is OLEIC ACID (three-letter code: OLA) (formula:  $C_{18}H_{34}O_2$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 18 16 2	0	0
3	A	1	Total C 13 13	0	0
3	A	1	Total C O 12 10 2	0	0
3	B	1	Total C O 16 14 2	0	0
3	B	1	Total C 11 11	0	0
3	B	1	Total C O 12 10 2	0	0
3	B	1	Total C 13 13	0	0

- Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Na 1 1	0	0
4	B	1	Total Na 1 1	0	0

- Molecule 5 is (2R)-2,3-dihydroxypropyl (9Z)-octadec-9-enoate (three-letter code: OLC) (formula:  $C_{21}H_{40}O_4$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 13 9 4	0	0
5	A	1	Total C O 13 9 4	0	0
5	B	1	Total C O 13 9 4	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	11	Total O 11 11	0	0
6	B	14	Total O 14 14	0	0
6	G	1	Total O 1 1	0	0
6	H	1	Total O 1 1	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Soluble cytochrome b562,Delta-type opioid receptor



- Molecule 1: Soluble cytochrome b562,Delta-type opioid receptor



- Molecule 2: bifunctional peptide



- Molecule 2: bifunctional peptide



## 4 Data and refinement statistics i

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	156.23Å 89.29Å 96.42Å 90.00° 92.30° 90.00°	Depositor
Resolution (Å)	33.45 – 2.70 32.74 – 2.70	Depositor EDS
% Data completeness (in resolution range)	99.9 (33.45-2.70) 89.3 (32.74-2.70)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.36 (at 2.68Å)	Xtriage
Refinement program	REFMAC 5.8.0049	Depositor
R, $R_{free}$	0.208 , 0.238 0.211 , 0.237	Depositor DCC
$R_{free}$ test set	1831 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	63.7	Xtriage
Anisotropy	0.066	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 41.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	0.033 for -1/2*h-3/2*k,-1/2*h+1/2*k,-l 0.033 for -1/2*h+3/2*k,1/2*h+1/2*k,-l 0.035 for 1/2*h-3/2*k,-1/2*h-1/2*k,-l 0.034 for 1/2*h+3/2*k,1/2*h-1/2*k,-l 0.258 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	6326	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	66.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.12% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: OLC, OLA, DI8, DI7, NA, NH2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.32	0/3094	0.48	0/4221
1	B	0.33	0/3099	0.48	0/4226
2	G	1.90	2/23 (8.7%)	1.49	0/29
2	H	1.82	2/23 (8.7%)	1.01	0/29
All	All	0.36	4/6239 (0.1%)	0.49	0/8505

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	G	3	PHE	CB-CG	-5.77	1.41	1.51
2	G	4	PHE	CB-CG	-5.64	1.41	1.51
2	H	4	PHE	CB-CG	-5.60	1.41	1.51
2	H	3	PHE	CB-CG	-5.37	1.42	1.51

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3030	0	3043	9	0
1	B	3035	0	3052	6	0
2	G	49	0	41	1	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	H	49	0	40	2	0
3	A	43	0	61	0	0
3	B	52	0	76	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	26	0	30	0	0
5	B	13	0	15	0	0
6	A	11	0	0	0	0
6	B	14	0	0	0	0
6	G	1	0	0	0	0
6	H	1	0	0	0	0
All	All	6326	0	6358	15	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (15) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1034:ARG:CD	1:B:1031:THR:HG21	2.39	0.53
2:H:1:DI7:HA	2:H:1:DI7:HE4B	1.95	0.48
1:A:1034:ARG:HD2	1:B:1031:THR:HG21	1.98	0.46
2:G:1:DI7:HE4B	2:G:1:DI7:HA	1.99	0.45
1:A:181:VAL:HG22	1:A:184:MET:CE	2.49	0.43
1:B:224:VAL:HB	1:B:225:PRO:HD3	2.02	0.42
1:A:224:VAL:HB	1:A:225:PRO:HD3	2.02	0.42
1:A:1005:ASP:O	1:A:1009:THR:HG23	2.21	0.41
1:A:275:ALA:N	1:A:276:PRO:CD	2.84	0.41
1:B:1005:ASP:O	1:B:1009:THR:HG23	2.21	0.41
1:B:86:ILE:HG22	1:B:169:ASN:ND2	2.36	0.41
1:B:275:ALA:N	1:B:276:PRO:CD	2.84	0.41
1:A:128:ASP:OD2	2:H:1:DI7:N	2.50	0.40
1:A:181:VAL:HA	1:A:184:MET:HE3	2.03	0.40
1:A:1099:ASN:HA	1:A:1103:GLN:HB2	2.04	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	399/411 (97%)	390 (98%)	9 (2%)	0	100	100
1	B	398/411 (97%)	389 (98%)	9 (2%)	0	100	100
2	G	2/5 (40%)	2 (100%)	0	0	100	100
2	H	2/5 (40%)	2 (100%)	0	0	100	100
All	All	801/832 (96%)	783 (98%)	18 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	310/346 (90%)	300 (97%)	10 (3%)	39	68
1	B	314/346 (91%)	304 (97%)	10 (3%)	39	68
2	G	2/2 (100%)	2 (100%)	0	100	100
2	H	2/2 (100%)	2 (100%)	0	100	100
All	All	628/696 (90%)	608 (97%)	20 (3%)	39	68

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1038	LEU
1	A	1039	ASP

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	41	ARG
1	A	69	LEU
1	A	108	LYS
1	A	128	ASP
1	A	146	ARG
1	A	249	SER
1	A	261	ARG
1	A	280	PHE
1	B	1039	ASP
1	B	41	ARG
1	B	69	LEU
1	B	108	LYS
1	B	128	ASP
1	B	146	ARG
1	B	155	LYS
1	B	249	SER
1	B	261	ARG
1	B	280	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	152	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	DI8	G	2	2	11,13,14	2.62	3 (27%)	13,17,19	1.32	2 (15%)
2	DI7	G	1	2	13,14,15	3.40	3 (23%)	15,19,21	1.03	2 (13%)
2	DI8	H	2	2	11,13,14	2.44	2 (18%)	13,17,19	1.29	2 (15%)
2	DI7	H	1	2	13,14,15	3.47	3 (23%)	15,19,21	1.22	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	DI8	G	2	2	-	0/1/11/13	0/2/2/2
2	DI7	G	1	2	-	0/5/6/8	0/1/1/1
2	DI8	H	2	2	-	0/1/11/13	0/2/2/2
2	DI7	H	1	2	-	0/5/6/8	0/1/1/1

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	H	1	DI7	CB-CD	-10.06	1.39	1.51
2	G	1	DI7	CB-CD	-9.86	1.39	1.51
2	G	2	DI8	C1-C2	-7.67	1.37	1.51
2	H	2	DI8	C1-C2	-6.97	1.38	1.51
2	G	1	DI7	CE4-CD2	-5.32	1.40	1.51
2	H	1	DI7	CE3-CD1	-5.16	1.40	1.51
2	H	1	DI7	CE4-CD2	-4.96	1.41	1.51
2	G	1	DI7	CE3-CD1	-4.61	1.41	1.51
2	H	2	DI8	C8-C7	-3.40	1.39	1.50
2	G	2	DI8	C8-C7	-3.34	1.40	1.50
2	G	2	DI8	C8-N	2.10	1.49	1.46

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	G	2	DI8	C1-CA-N	-3.31	105.66	112.50
2	H	2	DI8	C1-CA-N	-3.15	106.00	112.50
2	H	1	DI7	CE4-CD2-CD	2.98	125.67	122.01
2	H	1	DI7	CE4-CD2-CE2	-2.71	114.47	119.49
2	G	1	DI7	CE4-CD2-CD	2.39	124.94	122.01
2	G	2	DI8	C2-C1-CA	-2.34	108.40	111.29
2	G	1	DI7	CE4-CD2-CE2	-2.19	115.44	119.49

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	2	DI8	C2-C1-CA	-2.10	108.70	111.29

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	G	1	DI7	1	0
2	H	1	DI7	2	0

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 2 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	OLA	B	1201	-	15,15,19	0.57	0	15,15,19	0.69	0
3	OLA	B	1204	-	11,11,19	0.70	0	11,11,19	0.79	0
3	OLA	A	1206	-	11,11,19	0.67	0	11,11,19	0.74	0
3	OLA	B	1206	-	12,12,19	0.34	0	11,11,19	0.35	0
5	OLC	A	1205	-	12,12,24	1.36	1 (8%)	13,13,25	1.06	2 (15%)
5	OLC	B	1205	-	12,12,24	1.27	1 (8%)	13,13,25	1.05	2 (15%)
5	OLC	A	1204	-	12,12,24	1.26	1 (8%)	13,13,25	1.04	1 (7%)
3	OLA	A	1203	-	12,12,19	0.33	0	11,11,19	0.36	0
3	OLA	B	1203	-	10,10,19	0.35	0	9,9,19	0.57	0
3	OLA	A	1201	-	17,17,19	0.58	0	17,17,19	0.58	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	OLA	B	1201	-	-	5/13/13/17	-
3	OLA	B	1204	-	-	3/9/9/17	-
3	OLA	A	1206	-	-	6/9/9/17	-
3	OLA	B	1206	-	-	5/10/10/17	-
5	OLC	A	1205	-	-	5/12/12/24	-
5	OLC	B	1205	-	-	8/12/12/24	-
5	OLC	A	1204	-	-	5/12/12/24	-
3	OLA	A	1203	-	-	6/10/10/17	-
3	OLA	B	1203	-	-	3/8/8/17	-
3	OLA	A	1201	-	-	8/15/15/17	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	1205	OLC	O20-C1	4.53	1.46	1.33
5	B	1205	OLC	O20-C1	4.25	1.45	1.33
5	A	1204	OLC	O20-C1	4.17	1.45	1.33

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	1205	OLC	O20-C1-C2	2.78	120.62	111.91
5	B	1205	OLC	O20-C1-C2	2.75	120.53	111.91
5	A	1204	OLC	O20-C1-C2	2.64	120.21	111.91
5	A	1205	OLC	O20-C1-O19	-2.08	118.34	123.59
5	B	1205	OLC	O20-C1-O19	-2.05	118.41	123.59

There are no chirality outliers.

All (54) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	1204	OLC	C21-C22-C24-O25
5	B	1205	OLC	C21-C22-C24-O25
5	B	1205	OLC	O20-C21-C22-C24
5	B	1205	OLC	O20-C21-C22-O23

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
5	A	1205	OLC	C2-C1-O20-C21
5	B	1205	OLC	C2-C1-O20-C21
5	A	1205	OLC	O19-C1-O20-C21
5	B	1205	OLC	O19-C1-O20-C21
3	A	1206	OLA	C5-C6-C7-C8
3	A	1206	OLA	C2-C3-C4-C5
3	B	1203	OLA	C11-C12-C13-C14
3	A	1201	OLA	C2-C3-C4-C5
3	A	1201	OLA	C5-C6-C7-C8
3	A	1201	OLA	C11-C12-C13-C14
3	B	1206	OLA	C11-C12-C13-C14
5	B	1205	OLC	C2-C3-C4-C5
3	A	1203	OLA	C11-C12-C13-C14
5	A	1204	OLC	O23-C22-C24-O25
5	B	1205	OLC	O23-C22-C24-O25
3	A	1206	OLA	C4-C5-C6-C7
3	B	1201	OLA	C3-C4-C5-C6
3	B	1204	OLA	C5-C6-C7-C8
3	B	1204	OLA	C1-C2-C3-C4
3	A	1203	OLA	C6-C7-C8-C9
3	B	1206	OLA	C5-C6-C7-C8
3	A	1206	OLA	C6-C7-C8-C9
3	B	1201	OLA	C2-C3-C4-C5
3	A	1201	OLA	C6-C7-C8-C9
5	B	1205	OLC	C1-C2-C3-C4
5	A	1205	OLC	C1-C2-C3-C4
5	A	1205	OLC	C2-C3-C4-C5
3	B	1206	OLA	C10-C11-C12-C13
3	A	1203	OLA	C4-C5-C6-C7
5	A	1204	OLC	C2-C3-C4-C5
3	B	1204	OLA	C2-C3-C4-C5
3	B	1203	OLA	C15-C16-C17-C18
3	B	1201	OLA	C11-C12-C13-C14
3	A	1206	OLA	C1-C2-C3-C4
5	A	1205	OLC	C3-C4-C5-C6
3	A	1201	OLA	O2-C1-C2-C3
3	A	1206	OLA	C7-C8-C9-C10
3	A	1201	OLA	O1-C1-C2-C3
3	A	1203	OLA	C7-C8-C9-C10
3	A	1201	OLA	C10-C11-C12-C13
3	A	1203	OLA	C10-C11-C12-C13
3	B	1201	OLA	C9-C10-C11-C12

*Continued on next page...*




Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	A	1201	OLA	C9-C10-C11-C12
3	A	1203	OLA	C9-C10-C11-C12
3	B	1206	OLA	C9-C10-C11-C12
3	B	1206	OLA	C3-C4-C5-C6
3	B	1201	OLA	C1-C2-C3-C4
5	A	1204	OLC	O20-C1-C2-C3
3	B	1203	OLA	C9-C10-C11-C12
5	A	1204	OLC	O19-C1-C2-C3




There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	401/411 (97%)	-0.33	4 (0%) 82 83	44, 63, 101, 138	0
1	B	400/411 (97%)	-0.28	5 (1%) 77 78	41, 62, 106, 153	0
2	G	2/5 (40%)	-0.61	0 100 100	56, 56, 56, 77	0
2	H	2/5 (40%)	0.63	0 100 100	60, 60, 60, 88	0
All	All	805/832 (96%)	-0.30	9 (1%) 80 82	41, 62, 101, 153	0

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	242	SER	8.9
1	B	238	LEU	4.8
1	A	1001	ALA	3.2
1	A	242	SER	3.2
1	A	241	ARG	2.2
1	B	241	ARG	2.2
1	B	329	PHE	2.1
1	A	244	ARG	2.1
1	B	244	ARG	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	DI7	H	1	14/15	0.97	0.12	43,45,53,55	0
2	DI7	G	1	14/15	0.98	0.14	39,42,49,52	0
2	DI8	G	2	12/13	0.98	0.17	49,52,54,56	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	DI8	H	2	12/13	0.98	0.17	55,56,59,60	0

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	OLA	A	1206	12/20	0.78	0.29	85,90,93,94	0
3	OLA	B	1201	16/20	0.78	0.27	76,89,94,97	0
3	OLA	B	1204	12/20	0.85	0.26	81,85,92,94	0
3	OLA	B	1203	11/20	0.89	0.19	80,84,93,94	0
3	OLA	A	1203	13/20	0.89	0.27	78,87,98,99	0
3	OLA	A	1201	18/20	0.90	0.23	72,83,91,91	0
5	OLC	A	1205	13/25	0.91	0.20	65,76,85,86	0
5	OLC	B	1205	13/25	0.91	0.24	68,74,78,86	0
4	NA	B	1202	1/1	0.92	0.08	62,62,62,62	0
3	OLA	B	1206	13/20	0.93	0.27	69,72,81,81	0
4	NA	A	1202	1/1	0.94	0.10	66,66,66,66	0
5	OLC	A	1204	13/25	0.94	0.25	66,70,80,82	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around OLA A 1206:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around OLA B 1201:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)













**Electron density around OLA A 1203:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around OLA A 1201:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around OLC A 1205:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around OLC B 1205:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around OLA B 1206:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around OLC A 1204:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.