

# Full wwPDB X-ray Structure Validation Report (i)

#### Sep 14, 2023 – 07:11 PM EDT

PDB ID	:	4S1S
Title	:	Crystal structure of a VRC01-lineage antibody, 45-VRC01.H5.F-185917, in
		complex with clade A/E HIV-1 gp120 core
Authors	:	Kwon, Y.D.; Yang, Y.; Zhang, B.; Kwong, P.D.
Deposited on	:	2015-01-14
Resolution	:	3.39  Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.35.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35.1

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $X\text{-}RAY \, DIFFRACTION$ 

The reported resolution of this entry is 3.39 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
	$(\# { m Entries})$	$(\# { m Entries},  { m resolution}  { m range}({ m \AA}))$
$R_{free}$	130704	1026 (3.48-3.32)
Clashscore	141614	1055 (3.48-3.32)
Ramachandran outliers	138981	1038 (3.48-3.32)
Sidechain outliers	138945	1038 (3.48-3.32)
RSRZ outliers	127900	2173 (3.50-3.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	G	353	7%	18%	·
2	Н	228	75%	23%	•
3	L	210	80%	19%	·



# 2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 6187 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called clade A/E 93TH057 HIV-1 gp120 core.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	G	342	Total 2677	C 1680	N 466	O 509	S 22	0	0	0

• Molecule 2 is a protein called Fab of VRC01-lineage antibody,45-VRC01.H5.F-185917 heavy chain.

Mol	Chain	Residues		Ate	oms			ZeroOcc	AltConf	Trace
2	Н	228	Total 1746	C 1108	N 303	O 327	S 8	0	0	0

• Molecule 3 is a protein called Fab of VRC01 light chain.

Mol	Chain	Residues		Ate	oms			ZeroOcc	AltConf	Trace
3	L	208	Total 1609	C 1008	N 273	0 323	${ m S}{ m 5}$	0	0	0

• Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	G	1	Total         C         N         O           14         8         1         5	0	0
4	G	1	Total         C         N         O           14         8         1         5	0	0
4	G	1	Total         C         N         O           14         8         1         5	0	0
4	G	1	Total         C         N         O           14         8         1         5	0	0
4	G	1	Total         C         N         O           14         8         1         5	0	0
4	G	1	Total         C         N         O           14         8         1         5	0	0
4	G	1	Total         C         N         O           14         8         1         5	0	0
4	G	1	Total         C         N         O           14         8         1         5	0	0
4	G	1	Total         C         N         O           14         8         1         5	0	0
4	G	1	Total         C         N         O           14         8         1         5	0	0

• Molecule 5 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (three-letter code: EPE) (formula:  $C_8H_{18}N_2O_4S$ ).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf		
5	G	1	Total 15	C 8	N 2	0 4	S 1	0	0



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.





• Molecule 2: Fab of VRC01-lineage antibody,45-VRC01.H5.F-185917 heavy chain



• Molecule 3: Fab of VRC01 light chain





## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	55.39Å $67.66$ Å $266.75$ Å	Deperitor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.00^{\circ}$ $90.00^{\circ}$	Depositor
$\mathbf{P}_{\mathrm{oscolution}}(\mathbf{\hat{A}})$	47.50 - 3.39	Depositor
Resolution (A)	47.50 - 3.39	EDS
% Data completeness	96.8 (47.50-3.39)	Depositor
(in resolution range)	$92.2 \ (47.50-3.39)$	EDS
R <sub>merge</sub>	0.07	Depositor
R <sub>sym</sub>	0.10	Depositor
$< I/\sigma(I) > 1$	$1.95 (at 3.40 \text{\AA})$	Xtriage
Refinement program	PHENIX (phenix.refine: dev_1839)	Depositor
D D.	0.262 , $0.300$	Depositor
$\Pi, \Pi_{free}$	0.265 , $0.305$	DCC
$R_{free}$ test set	698 reflections $(4.92%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	83.8	Xtriage
Anisotropy	0.236	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.27 , $47.2$	EDS
L-test for twinning <sup>2</sup>	$< L >=0.46, < L^2>=0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.88	EDS
Total number of atoms	6187	wwPDB-VP
Average B, all atoms $(Å^2)$	110.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 4.01% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

### 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, EPE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond	lengths	Bond angles		
	Unam	RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	G	0.25	0/2733	0.54	0/3709	
2	Н	0.23	0/1797	0.46	0/2449	
3	L	0.20	0/1646	0.37	0/2235	
All	All	0.23	0/6176	0.48	0/8393	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

#### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	G	2677	0	2611	37	0
2	Н	1746	0	1709	50	0
3	L	1609	0	1547	22	0
4	G	140	0	130	3	0
5	G	15	0	17	0	0
All	All	6187	0	6014	106	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (106) close contacts within the same asymmetric unit are listed below, sorted by their clash



magnitude.

Atom 1	Atom 2	Interatomic	Clash	
Atom-1	Atom-2	distance $(\text{\AA})$	overlap (Å)	
2:H:138:LEU:HD11	2:H:211:VAL:CG1	1.13	1.54	
2:H:138:LEU:CD1	2:H:211:VAL:CG1	2.06	1.32	
2:H:138:LEU:CD1	2:H:211:VAL:HG11	1.63	1.22	
2:H:138:LEU:HD11	2:H:211:VAL:HG12	1.11	1.06	
2:H:138:LEU:HD11	2:H:211:VAL:HG11	1.05	1.05	
2:H:97:ASP:HB3	2:H:98:CYS:HA	1.39	1.02	
2:H:138:LEU:HD11	2:H:211:VAL:HG13	1.50	0.88	
2:H:138:LEU:CD1	2:H:211:VAL:HG12	1.89	0.79	
2:H:126:PRO:HD3	2:H:138:LEU:HD13	1.67	0.76	
2:H:138:LEU:CG	2:H:211:VAL:HG11	2.17	0.74	
1:G:298:ARG:NH2	1:G:439:ILE:O	2.25	0.69	
2:H:82:VAL:HG12	2:H:82(C):LEU:CD2	2.24	0.68	
1:G:101:VAL:HG21	1:G:480:ARG:HG2	1.78	0.65	
1:G:277:LEU:O	1:G:456:ARG:NH2	2.29	0.65	
2:H:97:ASP:HB3	2:H:98:CYS:CA	2.24	0.65	
2:H:138:LEU:CG	2:H:211:VAL:CG1	2.74	0.65	
2:H:197:ASN:ND2	2:H:208:ASP:OD1	2.30	0.65	
2:H:67:PHE:CE1	2:H:82:VAL:HG22	2.34	0.63	
3:L:79:GLU:O	3:L:81:GLY:N	2.34	0.61	
1:G:280:ASN:ND2	3:L:96:GLU:OE1	2.34	0.61	
2:H:126:PRO:HB2	2:H:127:SER:HA	1.82	0.60	
1:G:233:PHE:O	1:G:273:ARG:NH2	2.30	0.59	
2:H:12:LYS:HE3	2:H:18:VAL:HG22	1.84	0.59	
2:H:115:SER:HB3	2:H:116:THR:HA	1.84	0.59	
3:L:63:SER:OG	3:L:74:THR:OG1	2.20	0.58	
2:H:82:VAL:HG12	2:H:82(C):LEU:HD21	1.85	0.57	
2:H:82:VAL:CG1	2:H:82(C):LEU:HD21	2.34	0.57	
2:H:138:LEU:HD21	2:H:211:VAL:HG12	1.87	0.56	
1:G:343:LYS:O	1:G:346:THR:OG1	2.24	0.56	
1:G:298:ARG:NH1	1:G:326:ILE:O	2.35	0.56	
1:G:461:ASN:OD1	1:G:462:ASN:N	2.37	0.56	
2:H:159:LEU:HD21	2:H:182:VAL:HG11	1.88	0.56	
1:G:361:PHE:HB3	1:G:391:PHE:HB3	1.88	0.55	
2:H:82:VAL:HG12	2:H:82(C):LEU:HD23	1.87	0.55	
2:H:67:PHE:CD1	2:H:80:LEU:HD11	2.42	0.55	
3:L:191:HIS:O	3:L:213:ARG:NH1	2.39	0.55	
1:G:260:LEU:HD12	1:G:451:GLY:HA3	1.90	0.54	
2:H:82:VAL:CG1	2:H:82(C):LEU:CD2	2.85	0.54	
1:G:121:LYS:N	1:G:201:ILE:O	2.38	0.53	
2:H:67:PHE:HD1	2:H:80:LEU:HD11	1.73	0.53	
2:H:100(A):ASP:HB3	2:H:100(B):PRO:HD3	1.89	0.53	



Atom 1	Atom 2	Interatomic	Clash	
Atom-1	Atom-2	distance $(\text{\AA})$	overlap (Å)	
1:G:291:SER:OG	4:G:510:NAG:O6	2.28	0.52	
2:H:47:TRP:CZ2	2:H:49:GLY:HA2	2.46	0.51	
3:L:47:VAL:HG12	3:L:48:ILE:HG12	1.93	0.50	
1:G:120:VAL:HB	1:G:434:MET:HE3	1.94	0.50	
1:G:454:LEU:HD23	1:G:470:PRO:HA	1.93	0.50	
2:H:138:LEU:CD2	2:H:211:VAL:HG12	2.42	0.49	
2:H:68:THR:OG1	2:H:81:ASP:HB2	2.13	0.49	
2:H:200:HIS:CD2	2:H:202:PRO:HD2	2.47	0.49	
1:G:119:CYS:N	1:G:205:CYS:HB3	2.27	0.49	
1:G:455:THR:HG23	1:G:471:GLY:HA3	1.93	0.49	
1:G:241:ASN:HB3	4:G:502:NAG:O5	2.12	0.49	
3:L:120:PHE:HB2	3:L:135:VAL:HB	1.95	0.49	
3:L:47:VAL:HA	3:L:58:ILE:HG13	1.96	0.48	
3:L:130:GLY:HA2	3:L:185:LYS:HB2	1.94	0.48	
1:G:52:LEU:HD11	1:G:100:MET:HG2	1.95	0.47	
2:H:18:VAL:HG23	2:H:82(C):LEU:HD11	1.96	0.47	
3:L:24:ARG:HG2	3:L:70:ASP:OD2	2.15	0.47	
1:G:64:GLU:HA	1:G:209:SER:HB3	1.96	0.47	
1:G:266:ALA:HB3	1:G:289:ASN:HB3	1.96	0.47	
3:L:115:PRO:HD3	3:L:200:HIS:HD2	1.79	0.46	
1:G:476:LYS:HA	1:G:479:TRP:CD1	2.50	0.46	
2:H:136:ALA:N	2:H:184:VAL:O	2.49	0.46	
1:G:104:MET:HE3	1:G:104:MET:HB2	1.87	0.45	
3:L:78:LEU:HD23	3:L:83:PHE:CE1	2.52	0.45	
2:H:193:THR:HG23	2:H:210:LYS:HE3	1.99	0.45	
3:L:157:GLN:HE21	3:L:160:ASN:HD21	1.64	0.45	
1:G:119:CYS:N	1:G:434:MET:O	2.47	0.44	
2:H:96:ALA:HB2	2:H:101:PRO:HG3	2.00	0.44	
1:G:277:LEU:HD13	1:G:352:HIS:HB3	1.99	0.44	
2:H:18:VAL:O	2:H:81:ASP:HA	2.18	0.44	
2:H:89:VAL:HG12	2:H:108:LEU:HD23	1.99	0.44	
1:G:54:CYS:O	1:G:75:VAL:HG22	2.18	0.43	
1:G:396:ILE:O	1:G:396:ILE:HG22	2.18	0.43	
1:G:474:ASN:OD1	2:H:100(B):PRO:HG3	2.17	0.43	
2:H:68:THR:O	2:H:80:LEU:HD12	2.18	0.43	
3:L:142:TYR:CG	3:L:143:PRO:HA	2.54	0.43	
3:L:110:ARG:HD2	3:L:173:SER:HB2	2.00	0.43	
1:G:348:LYS:HD3	1:G:348:LYS:HA	1.90	0.43	
2:H:123:PRO:HD3	2:H:209:LYS:HE2	2.00	0.43	
3:L:30:SER:HB2	3:L:50:SER:HA	2.00	0.43	
3:L:192:LYS:HD3	3:L:213:ARG:HH12	1.83	0.43	

Continued from previous page...



Atom 1	Atom 2	Interatomic	Clash	
Atom-1	Atom-2	distance (Å)	overlap (Å)	
3:L:60:ASP:OD1	3:L:60:ASP:N	2.52	0.43	
2:H:27:TYR:CZ	2:H:94:ARG:HD2	2.54	0.42	
2:H:38:ARG:HB3	2:H:48:ILE:HD11	2.00	0.42	
1:G:49:ASP:OD2	1:G:99:ASN:ND2	2.52	0.42	
1:G:429:GLY:HA3	2:H:53:ARG:HH11	1.84	0.42	
1:G:475:ILE:O	1:G:478:ASN:HB2	2.19	0.42	
2:H:148:GLU:HA	2:H:149:PRO:HA	1.83	0.42	
1:G:390:LEU:HD23	1:G:390:LEU:HA	1.89	0.42	
3:L:5:THR:HB	3:L:24:ARG:HB2	2.01	0.41	
2:H:2:VAL:HB	2:H:102:HIS:CD2	2.55	0.41	
2:H:138:LEU:HD21	2:H:211:VAL:CG1	2.49	0.41	
3:L:121:PRO:HA	3:L:122:PRO:HD3	1.97	0.41	
2:H:152:VAL:HG22	2:H:198:VAL:HG22	2.03	0.41	
3:L:115:PRO:HD3	3:L:200:HIS:CD2	2.56	0.41	
1:G:346:THR:O	1:G:350:LYS:HG3	2.20	0.41	
2:H:97:ASP:CB	2:H:98:CYS:HA	2.23	0.41	
1:G:78:ASP:HA	1:G:79:PRO:HD3	1.86	0.41	
3:L:34:ALA:HA	3:L:49:TYR:HA	2.03	0.41	
1:G:420:ILE:HG21	1:G:438:PRO:HG3	2.03	0.41	
2:H:146:PHE:HA	2:H:147:PRO:HA	1.87	0.41	
2:H:67:PHE:HE1	2:H:82:VAL:HG22	1.84	0.41	
1:G:207:LYS:HE2	1:G:436:ALA:HB3	2.04	0.40	
1:G:389:GLN:HG2	4:G:509:NAG:C8	2.52	0.40	
3:L:203:LEU:HD13	3:L:207:VAL:HG23	2.04	0.40	

Continued from previous page...

There are no symmetry-related clashes.

#### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Analysed Favoured Allowed		Outliers	Percentiles	
1	G	336/353~(95%)	315 (94%)	18 (5%)	3 (1%)	17	49



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	Н	226/228~(99%)	194 (86%)	24 (11%)	8 (4%)	3	21
3	L	206/210~(98%)	189 (92%)	15 (7%)	2(1%)	15	46
All	All	768/791~(97%)	698 (91%)	57 (7%)	13 (2%)	9	34

Continued from previous page...

All (13) Ramachandran outliers are listed below:

Mol	Chain	$\operatorname{Res}$	Type	
3	L	80	SER	
1	G	232	ASN	
2	Н	96	ALA	
2	Н	100(E)	GLY	
2	Н	135	THR	
1	G	204	ALA	
2	Н	53	ARG	
2	Н	129	LYS	
2	Н	149	PRO	
1	G	411	ASN	
2	Н	14	PRO	
2	Н	100(B)	PRO	
3	Ĺ	206	PRO	

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the side chain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric Outliers		Percentiles		
1	G	305/311~(98%)	301~(99%)	4 (1%)	69	84	
2	Н	194/194~(100%)	189~(97%)	5(3%)	46	72	
3	L	180/182~(99%)	179~(99%)	1 (1%)	86	94	
All	All	679/687~(99%)	669~(98%)	10 (2%)	65	82	

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	65	VAL
· · · · · · · · · · · · · · · · · · ·	ã	-	



Continuaca from precioas page								
Mol	l Chain Res		Type					
1	G	231	LYS					
1	G	410	CYS					
1	G	434	MET					
2	Н	71	ARG					
2	Н	93	VAL					
2	Н	98	CYS					
2	Н	100(C)	CYS					
2	Н	216	CYS					
3	L	216	CYS					

Continued from previous page..

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
3	L	157	GLN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

#### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

#### 5.6 Ligand geometry (i)

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



4S1S
------

Mal	Tuno	Chain	Dog	Link	Bo	ond leng	$_{\rm ths}$	Bond angles		
WIOI	туре	Ullalli	nes		Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2
4	NAG	G	507	1	14,14,15	0.24	0	17,19,21	0.35	0
4	NAG	G	508	1	14,14,15	0.28	0	17,19,21	0.52	0
4	NAG	G	502	1	$14,\!14,\!15$	0.21	0	17,19,21	0.47	0
4	NAG	G	506	1	$14,\!14,\!15$	0.18	0	17,19,21	0.38	0
5	EPE	G	511	-	$15,\!15,\!15$	0.80	1 (6%)	18,20,20	1.84	5 (27%)
4	NAG	G	505	1	14,14,15	0.16	0	17,19,21	0.70	1 (5%)
4	NAG	G	504	1	14,14,15	0.30	0	17,19,21	0.38	0
4	NAG	G	509	1	14,14,15	0.21	0	17,19,21	0.44	0
4	NAG	G	510	1	14,14,15	0.16	0	17,19,21	0.57	0
4	NAG	G	501	1	14,14,15	0.30	0	17,19,21	0.35	0
4	NAG	G	503	1	14,14,15	0.24	0	17,19,21	0.50	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	G	507	1	-	2/6/23/26	0/1/1/1
4	NAG	G	508	1	-	2/6/23/26	0/1/1/1
4	NAG	G	502	1	-	2/6/23/26	0/1/1/1
4	NAG	G	506	1	-	2/6/23/26	0/1/1/1
5	EPE	G	511	-	-	5/9/19/19	0/1/1/1
4	NAG	G	505	1	-	4/6/23/26	0/1/1/1
4	NAG	G	504	1	-	2/6/23/26	0/1/1/1
4	NAG	G	509	1	-	2/6/23/26	0/1/1/1
4	NAG	G	510	1	-	4/6/23/26	0/1/1/1
4	NAG	G	501	1	-	0/6/23/26	0/1/1/1
4	NAG	G	503	1	_	2/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	G	511	EPE	C10-S	2.75	1.81	1.77

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
5	G	511	EPE	C5-N4-C3	4.83	119.69	108.83



4S15	5
------	---

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
5	G	511	EPE	O1S-S-C10	2.59	110.04	106.92
4	G	505	NAG	C1-O5-C5	2.49	115.57	112.19
5	G	511	EPE	C7-N4-C5	2.36	117.28	111.23
5	G	511	EPE	C2-C3-N4	2.13	115.01	110.64
5	G	511	EPE	O2S-S-C10	2.06	109.39	106.92

Continued from previous page...

There are no chirality outliers.

Mol	Chain	Res	Type	Atoms
5	G	511	EPE	C10-C9-N1-C2
4	G	502	NAG	C4-C5-C6-O6
4	G	506	NAG	O5-C5-C6-O6
4	G	502	NAG	O5-C5-C6-O6
4	G	505	NAG	O5-C5-C6-O6
4	G	510	NAG	O5-C5-C6-O6
4	G	506	NAG	C4-C5-C6-O6
4	G	507	NAG	C4-C5-C6-O6
4	G	507	NAG	O5-C5-C6-O6
4	G	508	NAG	C4-C5-C6-O6
4	G	510	NAG	C4-C5-C6-O6
4	G	505	NAG	C4-C5-C6-O6
4	G	503	NAG	O5-C5-C6-O6
4	G	505	NAG	C8-C7-N2-C2
4	G	505	NAG	O7-C7-N2-C2
4	G	509	NAG	C8-C7-N2-C2
4	G	509	NAG	O7-C7-N2-C2
4	G	510	NAG	C8-C7-N2-C2
4	G	510	NAG	O7-C7-N2-C2
4	G	504	NAG	O5-C5-C6-O6
4	G	508	NAG	O5-C5-C6-O6
4	G	503	NAG	C4-C5-C6-O6
5	G	511	EPE	C9-C10-S-O3S
4	G	504	NAG	C4-C5-C6-O6
5	G	511	EPE	N4-C7-C8-O8
5	G	511	EPE	C10-C9-N1-C6
5	G	511	EPE	C9-C10-S-O1S

All (27) torsion outliers are listed below:

There are no ring outliers.

3 monomers are involved in 3 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	G	502	NAG	1	0
4	G	509	NAG	1	0
4	G	510	NAG	1	0

## 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 6 Fit of model and data (i)

## 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle RSRZ \rangle$	#RSRZ>2	$OWAB(Å^2)$	Q < 0.9
1	G	342/353~(96%)	0.61	23 (6%) 17 19	48, 109, 156, 179	0
2	Н	228/228~(100%)	0.65	26 (11%) 5 6	42, 96, 160, 204	0
3	L	208/210~(99%)	0.55	21 (10%) 7 8	69, 126, 181, 198	0
All	All	778/791~(98%)	0.61	70 (8%) 9 11	42, 112, 169, 204	0

All (70) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	123	THR	10.0
1	G	124	GLY	9.5
3	L	132	ALA	8.3
1	G	199	SER	8.2
1	G	198	GLY	7.2
1	G	460	ALA	7.2
2	Н	132	SER	7.2
1	G	201	ILE	6.9
1	G	202	LYS	5.9
3	L	134	VAL	5.0
1	G	122	LEU	4.5
1	G	200	VAL	4.3
1	G	338	TRP	4.3
2	Н	133	GLY	4.2
1	G	204	ALA	4.0
2	Н	125	ALA	4.0
3	L	133	SER	4.0
2	Н	126	PRO	4.0
2	Н	211	VAL	3.9
3	L	131	THR	3.8
2	Н	138	LEU	3.6
2	Н	139	GLY	3.4
3	L	195	ALA	3.4



4S1	S
-----	---

Mol	Chain	Res	Type	RSRZ
2	Н	141	LEU	3.3
3	L	151	LYS	3.3
3	L	196	CYS	3.3
1	G	203	GLN	3.3
1	G	271	ILE	3.3
2	Н	122	PHE	3.2
3	L	183	LEU	3.2
1	G	442	LYS	3.2
3	L	137	LEU	3.2
1	G	121	LYS	3.2
3	L	198	VAL	3.2
3	L	135	VAL	3.1
2	Н	150	VAL	3.1
2	Н	140	CYS	3.0
2	Н	115	SER	3.0
2	Н	183	THR	2.9
2	Н	124	LEU	2.9
2	Н	178	LEU	2.9
3	L	156	LEU	2.7
1	G	347	GLU	2.7
1	G	353	PHE	2.6
3	L	181	LEU	2.6
3	L	149	GLN	2.6
2	Н	198	VAL	2.6
2	Н	131	THR	2.5
3	L	197	GLU	2.5
3	L	150	TRP	2.4
2	Н	123	PRO	2.4
1	G	333	ILE	2.4
2	Н	194	TYR	2.4
3	L	136	CYS	2.3
3	L	188	TYR	2.3
2	Н	120	SER	2.3
2	Н	97	ASP	2.2
2	Н	142	VAL	2.2
3	L	194	TYR	2.2
1	G	349	LEU	2.2
1	G	288	LEU	2.2
3	L	126	GLN	2.2
3	L	124	ASP	2.2
1	G	462	ASN	2.2
2	Н	143	LYS	2.2

Continued from previous page...



Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	G	461	ASN	2.1
2	Н	130	SER	2.0
2	Н	169	VAL	2.0
2	Н	181	VAL	2.0
1	G	61	HIS	2.0

#### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

#### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(Å^2)$	Q<0.9
4	NAG	G	510	14/15	0.69	0.39	119,132,149,152	0
4	NAG	G	509	14/15	0.71	0.31	164,167,170,170	0
4	NAG	G	504	14/15	0.71	0.28	115,122,130,130	0
4	NAG	G	502	14/15	0.76	0.39	138,149,176,180	0
4	NAG	G	506	14/15	0.78	0.24	125,149,158,159	0
4	NAG	G	507	14/15	0.80	0.32	143,154,161,163	0
4	NAG	G	505	14/15	0.84	0.34	114,127,138,145	0
4	NAG	G	501	14/15	0.85	0.21	91,101,115,118	0
5	EPE	G	511	15/15	0.89	0.31	67,79,88,94	0
4	NAG	G	503	14/15	0.90	0.24	98,117,125,127	0
4	NAG	G	508	14/15	0.92	0.25	80,94,105,106	0

#### 6.5 Other polymers (i)

There are no such residues in this entry.

