



## wwPDB EM Validation Summary Report ⓘ

Apr 18, 2024 – 02:53 pm BST

PDB ID : 8S51  
EMDB ID : EMD-19718  
Title : RNA polymerase II core initially transcribing complex with an ordered RNA of 8 nt  
Authors : Zhan, Y.; Grabbe, F.; Oberbeckmann, E.; Dienemann, C.; Cramer, P.  
Deposited on : 2024-02-22  
Resolution : 3.10 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev92  
MolProbity : 4.02b-467  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
MapQ : 1.9.13  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36

# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:  
*ELECTRON MICROSCOPY*

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826
RNA backbone	4643	859

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion  $< 40\%$ ). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1970	
2	B	1300	
3	C	275	
4	D	184	
5	E	210	
6	F	127	
7	G	172	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
8	H	150	 89% 10%
9	I	125	 81% 10% 9%
10	J	67	 85% 10%
11	K	117	 83% 15%
12	L	58	 48% 28% 24%
13	M	316	 84% 9% 8%
14	N	139	 29% 9% 62%
15	O	339	 46% 7% 47%
16	P	8	 62% 38%
17	Q	517	 8% 22% 73%
18	R	249	 35% 83% 6% 11%
19	T	139	 36% 9% 55%
20	U	376	 27% 27% 70%
21	V	109	 77% 78% 13% 9%
22	W	439	 35% 38% 57%
23	X	291	 53% 57% 41%

## 2 Entry composition

There are 25 unique types of molecules in this entry. The entry contains 44796 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA-directed RNA polymerase subunit.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	1421	11257	7083	2013	2089	72	0	0

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	?	-	TYR	deletion	UNP A0A7M4DUC2
A	?	-	SER	deletion	UNP A0A7M4DUC2
A	?	-	PRO	deletion	UNP A0A7M4DUC2
A	?	-	THR	deletion	UNP A0A7M4DUC2
A	?	-	SER	deletion	UNP A0A7M4DUC2
A	?	-	PRO	deletion	UNP A0A7M4DUC2
A	?	-	SER	deletion	UNP A0A7M4DUC2
A	?	-	TYR	deletion	UNP A0A7M4DUC2
A	?	-	SER	deletion	UNP A0A7M4DUC2
A	?	-	PRO	deletion	UNP A0A7M4DUC2
A	?	-	THR	deletion	UNP A0A7M4DUC2
A	?	-	SER	deletion	UNP A0A7M4DUC2
A	?	-	PRO	deletion	UNP A0A7M4DUC2
A	?	-	SER	deletion	UNP A0A7M4DUC2

- Molecule 2 is a protein called DNA-directed RNA polymerase subunit beta.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	B	1121	8971	5675	1574	1658	64	0	0

- Molecule 3 is a protein called DNA-directed RNA polymerase II subunit RPB3.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
3	C	257	2059	1294	351	408	6	0	0

- Molecule 4 is a protein called RNA polymerase II subunit D.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
4	D	128	Total	C	N	O	S	0	0
			1050	656	178	212	4		

- Molecule 5 is a protein called DNA-directed RNA polymerase II subunit E.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
5	E	209	Total	C	N	O	S	0	0
			1721	1089	300	324	8		

- Molecule 6 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC2.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
6	F	79	Total	C	N	O	S	0	0
			636	406	108	117	5		

- Molecule 7 is a protein called DNA-directed RNA polymerase II subunit RPB7.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
7	G	171	Total	C	N	O	S	0	0
			1351	875	219	249	8		

- Molecule 8 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC3.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
8	H	148	Total	C	N	O	S	0	0
			1186	750	194	237	5		

- Molecule 9 is a protein called DNA-directed RNA polymerase II subunit RPB9.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
9	I	114	Total	C	N	O	S	0	0
			928	571	166	180	11		

- Molecule 10 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC5.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
10	J	64	Total	C	N	O	S	0	0
			507	328	86	87	6		

- Molecule 11 is a protein called DNA-directed RNA polymerase II subunit RPB11-a.

Mol	Chain	Residues	Atoms					AltConf	Trace
11	K	115	Total	C	N	O	S	0	0
			920	593	152	173	2		

- Molecule 12 is a protein called RNA polymerase II subunit K.

Mol	Chain	Residues	Atoms					AltConf	Trace
12	L	44	Total	C	N	O	S	0	0
			373	231	72	64	6		

- Molecule 13 is a protein called Transcription initiation factor IIB.

Mol	Chain	Residues	Atoms					AltConf	Trace
13	M	292	Total	C	N	O	S	0	0
			2252	1403	402	429	18		

- Molecule 14 is a DNA chain called Non-template DNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
14	N	53	Total	C	N	O	P	0	0
			1113	521	226	313	53		

- Molecule 15 is a protein called TATA-box-binding protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
15	O	179	Total	C	N	O	S	0	0
			1422	923	251	241	7		

- Molecule 16 is a RNA chain called RNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
16	P	8	Total	C	N	O	P	0	0
			169	76	32	53	8		

- Molecule 17 is a protein called General transcription factor IIF subunit 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
17	Q	138	Total	C	N	O	S	0	0
			1138	719	208	208	3		

- Molecule 18 is a protein called General transcription factor IIF subunit 2.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
18	R	222	1788	1127	320	338	3	0	0

- Molecule 19 is a DNA chain called Template DNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	P		
19	T	63	1270	604	218	385	63	0	0

- Molecule 20 is a protein called Transcription initiation factor IIA subunit 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
20	U	113	930	585	152	189	4	0	0

- Molecule 21 is a protein called Transcription initiation factor IIA subunit 2.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
21	V	99	806	510	142	151	3	0	0

- Molecule 22 is a protein called General transcription factor IIE subunit 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
22	W	187	1535	964	275	285	11	0	0

- Molecule 23 is a protein called Transcription initiation factor IIE subunit beta.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
23	X	171	1403	895	243	261	4	0	0

- Molecule 24 is ZINC ION (three-letter code: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		AltConf
24	A	2	Total	Zn	0
			2	2	
24	B	1	Total	Zn	0
			1	1	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms		AltConf
24	C	1	Total 1	Zn 1	0
24	I	2	Total 2	Zn 2	0
24	J	1	Total 1	Zn 1	0
24	L	1	Total 1	Zn 1	0
24	M	1	Total 1	Zn 1	0
24	W	1	Total 1	Zn 1	0


- Molecule 25 is MAGNESIUM ION (three-letter code: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		AltConf
25	A	1	Total 1	Mg 1	0












• Molecule 5: DNA-directed RNA polymerase II subunit E




• Molecule 6: DNA-directed RNA polymerases I, II, and III subunit RPABC2



• Molecule 7: DNA-directed RNA polymerase II subunit RPB7




• Molecule 8: DNA-directed RNA polymerases I, II, and III subunit RPABC3













• Molecule 22: General transcription factor IIE subunit 1



• Molecule 23: Transcription initiation factor IIE subunit beta



K183	L184	L185	G186	D187	Q188	I189	L190	F191	V192	N193	R194	P195	D196	K197	K198	K199	I200	L201	F202	F203	N204	D205	K206	S207	C208	Q209	F210	S211	V212	D213	E214	E215	F216	Q217	K218	L219	W220	R221	S222	V223	T224	V225	D226	S227	M228	D229	E230	E231	K232	I233	E234	E235	Y236	L237	K238	R239	Q240	G241	I242
SER	SER	MET	GLN	GLU	SER	GLY	PRO	LYS	LYS	VAL	ALA	PRO	ILE	GLN	ARG	ARG	LYS	LYS	PRO	ALA	SER	GLN	LYS	ARG	ARG	PHE	LYS	THR	HIS	ASN	GLU	HIS	LEU	ALA	GLY	VAL	LEU	LYS	ASP	TYR	SER	ASP	ILE	THR	SER	SER	LYS	LYS											



## 4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	87097	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	40	Depositor
Minimum defocus (nm)	700	Depositor
Maximum defocus (nm)	1700	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	54.772	Depositor
Minimum map value	-35.910	Depositor
Average map value	-0.000	Depositor
Map value standard deviation	1.000	Depositor
Recommended contour level	3.29	Depositor
Map size (Å)	419.99997, 419.99997, 419.99997	wwPDB
Map dimensions	400, 400, 400	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.05, 1.05, 1.05	Depositor

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.41	0/11462	0.58	0/15473
2	B	0.46	0/9149	0.58	0/12349
3	C	0.47	0/2102	0.54	0/2857
4	D	0.27	0/1064	0.51	0/1428
5	E	0.38	0/1752	0.59	0/2366
6	F	0.42	0/646	0.57	0/871
7	G	0.30	0/1382	0.53	0/1874
8	H	0.43	0/1207	0.58	0/1628
9	I	0.34	0/949	0.58	0/1284
10	J	0.54	0/516	0.55	0/696
11	K	0.41	0/939	0.50	0/1271
12	L	0.43	0/378	0.65	0/500
13	M	0.38	0/2286	0.65	0/3084
14	N	0.95	0/1254	1.19	0/1937
15	O	0.34	0/1448	0.66	0/1948
16	P	0.75	0/188	1.07	0/290
17	Q	0.26	0/1167	0.53	0/1576
18	R	0.30	0/1817	0.57	0/2445
19	T	1.07	0/1416	1.24	0/2176
20	U	0.25	0/945	0.51	0/1274
21	V	0.25	0/816	0.53	0/1105
22	W	0.25	0/1560	0.56	0/2097
23	X	0.24	0/1427	0.48	0/1916
All	All	0.45	0/45870	0.64	0/62445

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	11257	0	11394	110	0
2	B	8971	0	8999	98	0
3	C	2059	0	2007	24	0
4	D	1050	0	1033	3	0
5	E	1721	0	1737	6	0
6	F	636	0	665	6	0
7	G	1351	0	1358	10	0
8	H	1186	0	1147	11	0
9	I	928	0	859	10	0
10	J	507	0	523	4	0
11	K	920	0	942	12	0
12	L	373	0	378	13	0
13	M	2252	0	2269	18	0
14	N	1113	0	592	12	0
15	O	1422	0	1514	13	0
16	P	169	0	89	3	0
17	Q	1138	0	1103	17	0
18	R	1788	0	1819	10	0
19	T	1270	0	710	16	0
20	U	930	0	888	8	0
21	V	806	0	818	8	0
22	W	1535	0	1540	11	0
23	X	1403	0	1428	4	0
24	A	2	0	0	0	0
24	B	1	0	0	0	0
24	C	1	0	0	0	0
24	I	2	0	0	0	0
24	J	1	0	0	0	0
24	L	1	0	0	0	0
24	M	1	0	0	0	0
24	W	1	0	0	0	0
25	A	1	0	0	0	0
All	All	44796	0	43812	357	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 357 close contacts within the same asymmetric unit are listed below, sorted by

their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:74:CYS:SG	1:A:84:HIS:CE1	2.73	0.82
14:N:24:DG:H2'	14:N:25:DG:C8	2.16	0.81
1:A:74:CYS:SG	1:A:84:HIS:HE1	2.03	0.78
1:A:11:SER:N	2:B:1135:TYR:HH	1.83	0.77
1:A:801:GLY:H	2:B:503:ASN:HD22	1.36	0.72

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1411/1970 (72%)	1355 (96%)	56 (4%)	0	100	100
2	B	1111/1300 (86%)	1062 (96%)	49 (4%)	0	100	100
3	C	253/275 (92%)	241 (95%)	12 (5%)	0	100	100
4	D	126/184 (68%)	124 (98%)	2 (2%)	0	100	100
5	E	207/210 (99%)	202 (98%)	5 (2%)	0	100	100
6	F	77/127 (61%)	76 (99%)	1 (1%)	0	100	100
7	G	169/172 (98%)	167 (99%)	2 (1%)	0	100	100
8	H	146/150 (97%)	141 (97%)	5 (3%)	0	100	100
9	I	112/125 (90%)	106 (95%)	6 (5%)	0	100	100
10	J	62/67 (92%)	62 (100%)	0	0	100	100
11	K	113/117 (97%)	110 (97%)	3 (3%)	0	100	100
12	L	42/58 (72%)	38 (90%)	4 (10%)	0	100	100
13	M	288/316 (91%)	280 (97%)	8 (3%)	0	100	100
15	O	177/339 (52%)	171 (97%)	6 (3%)	0	100	100
17	Q	134/517 (26%)	128 (96%)	6 (4%)	0	100	100

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
18	R	218/249 (88%)	212 (97%)	5 (2%)	1 (0%)	29	64
20	U	109/376 (29%)	106 (97%)	2 (2%)	1 (1%)	17	52
21	V	97/109 (89%)	96 (99%)	1 (1%)	0	100	100
22	W	185/439 (42%)	182 (98%)	3 (2%)	0	100	100
23	X	169/291 (58%)	162 (96%)	7 (4%)	0	100	100
All	All	5206/7391 (70%)	5021 (96%)	183 (4%)	2 (0%)	100	100

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
20	U	346	LYS
18	R	231	ASN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	1252/1749 (72%)	1246 (100%)	6 (0%)	88	94
2	B	985/1127 (87%)	983 (100%)	2 (0%)	93	97
3	C	234/252 (93%)	234 (100%)	0	100	100
4	D	118/160 (74%)	117 (99%)	1 (1%)	81	92
5	E	191/192 (100%)	191 (100%)	0	100	100
6	F	69/111 (62%)	69 (100%)	0	100	100
7	G	152/153 (99%)	152 (100%)	0	100	100
8	H	129/131 (98%)	129 (100%)	0	100	100
9	I	103/112 (92%)	103 (100%)	0	100	100
10	J	53/56 (95%)	53 (100%)	0	100	100
11	K	104/106 (98%)	104 (100%)	0	100	100
12	L	41/55 (74%)	41 (100%)	0	100	100
13	M	247/268 (92%)	246 (100%)	1 (0%)	91	96

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
15	O	154/293 (53%)	151 (98%)	3 (2%)	57	81
17	Q	121/448 (27%)	120 (99%)	1 (1%)	81	92
18	R	196/218 (90%)	194 (99%)	2 (1%)	76	90
20	U	105/324 (32%)	105 (100%)	0	100	100
21	V	90/98 (92%)	89 (99%)	1 (1%)	73	89
22	W	169/373 (45%)	168 (99%)	1 (1%)	86	94
23	X	154/261 (59%)	154 (100%)	0	100	100
All	All	4667/6487 (72%)	4649 (100%)	18 (0%)	91	96

5 of 18 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
18	R	175	ARG
22	W	153	ARG
21	V	82	ARG
4	D	24	LYS
17	Q	105	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 43 such sidechains are listed below:

Mol	Chain	Res	Type
7	G	4	HIS
11	K	89	ASN
7	G	122	ASN
9	I	22	ASN
15	O	242	GLN

### 5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
16	P	7/8 (87%)	0	0

There are no RNA backbone outliers to report.

There are no RNA pucker outliers to report.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 11 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


## 6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-19718. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

### 6.1 Orthogonal projections [i](#)

#### 6.1.1 Primary map



#### 6.1.2 Raw map




The images above show the map projected in three orthogonal directions.




## 6.2 Central slices [i](#)


### 6.2.1 Primary map



X Index: 200




Y Index: 200




Z Index: 200


### 6.2.2 Raw map



X Index: 200



Y Index: 200




Z Index: 200


The images above show central slices of the map in three orthogonal directions.

## 6.3 Largest variance slices [i](#)


### 6.3.1 Primary map



X Index: 189




Y Index: 202




Z Index: 262


### 6.3.2 Raw map



X Index: 189



Y Index: 190




Z Index: 250


The images above show the largest variance slices of the map in three orthogonal directions.

## 6.4 Orthogonal standard-deviation projections (False-color) [i](#)


### 6.4.1 Primary map



X




Y




Z


### 6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


## 6.5 Orthogonal surface views [i](#)

### 6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 3.29. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

### 6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


## 6.6 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


### 6.6.1 emd\_19718\_msk\_1.map [i](#)



X



Y




Z

## 7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

### 7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


## 7.2 Volume estimate [i](#)



The volume at the recommended contour level is 363 nm<sup>3</sup>; this corresponds to an approximate mass of 328 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

### 7.3 Rotationally averaged power spectrum [i](#)




\*Reported resolution corresponds to spatial frequency of 0.323 Å<sup>-1</sup>



## 8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

### 8.1 FSC [i](#)



\*Reported resolution corresponds to spatial frequency of 0.323 Å<sup>-1</sup>

## 8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.10	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	3.68	6.61	3.78

\*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.68 differs from the reported value 3.1 by more than 10 %

## 9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-19718 and PDB model 8S51. Per-residue inclusion information can be found in section 3 on page 9.

### 9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 3.29 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

## 9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

## 9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (3.29).

















































## 9.4 Atom inclusion [i](#)



At the recommended contour level, 81% of all backbone atoms, 76% of all non-hydrogen atoms, are inside the map.

## 9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (3.29) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.7560	 0.4150
A	 0.8850	 0.4940
B	 0.9020	 0.5130
C	 0.9180	 0.5270
D	 0.3260	 0.1510
E	 0.8730	 0.4750
F	 0.9090	 0.5200
G	 0.5410	 0.2610
H	 0.8960	 0.5010
I	 0.8380	 0.4420
J	 0.9390	 0.5390
K	 0.9060	 0.5190
L	 0.8910	 0.4960
M	 0.8350	 0.4530
N	 0.7920	 0.2980
O	 0.7770	 0.3410
P	 0.8520	 0.4580
Q	 0.5870	 0.2850
R	 0.4960	 0.2520
T	 0.8090	 0.3550
U	 0.1260	 0.1130
V	 0.1370	 0.1430
W	 0.2030	 0.1280
X	 0.1250	 0.1230

