



# wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 10, 2020 – 07:53 AM BST

PDB ID : 3S9L  
Title : Complex between transferrin receptor 1 and transferrin with iron in the N-Lobe, cryocooled 2  
Authors : Eckenroth, B.E.; Steere, A.N.; Mason, A.B.; Everse, S.J.  
Deposited on : 2011-06-01  
Resolution : 3.22 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.13.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.22 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1335 (3.24-3.20)
Clashscore	141614	1460 (3.24-3.20)
Ramachandran outliers	138981	1437 (3.24-3.20)
Sidechain outliers	138945	1436 (3.24-3.20)
RSRZ outliers	127900	1291 (3.24-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	654	
1	B	654	
2	C	693	
2	D	693	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	CO3	D	905	-	-	X	-

## 2 Entry composition i

There are 6 unique types of molecules in this entry. The entry contains 16713 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transferrin receptor protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	638	4949	3169	828	938	14	0	0	0
1	B	639	4800	3064	799	923	14	0	0	0

There are 28 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	107	VAL	-	expression tag	UNP P02786
A	108	PRO	-	expression tag	UNP P02786
A	109	ASP	-	expression tag	UNP P02786
A	110	LYS	-	expression tag	UNP P02786
A	111	HIS	-	expression tag	UNP P02786
A	112	HIS	-	expression tag	UNP P02786
A	113	HIS	-	expression tag	UNP P02786
A	114	HIS	-	expression tag	UNP P02786
A	115	HIS	-	expression tag	UNP P02786
A	116	HIS	-	expression tag	UNP P02786
A	117	ILE	-	expression tag	UNP P02786
A	118	GLU	-	expression tag	UNP P02786
A	119	GLY	-	expression tag	UNP P02786
A	142	SER	GLY	SEE REMARK 999	UNP P02786
B	107	VAL	-	expression tag	UNP P02786
B	108	PRO	-	expression tag	UNP P02786
B	109	ASP	-	expression tag	UNP P02786
B	110	LYS	-	expression tag	UNP P02786
B	111	HIS	-	expression tag	UNP P02786
B	112	HIS	-	expression tag	UNP P02786
B	113	HIS	-	expression tag	UNP P02786
B	114	HIS	-	expression tag	UNP P02786
B	115	HIS	-	expression tag	UNP P02786
B	116	HIS	-	expression tag	UNP P02786
B	117	ILE	-	expression tag	UNP P02786

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	118	GLU	-	expression tag	UNP P02786
B	119	GLY	-	expression tag	UNP P02786
B	142	SER	GLY	SEE REMARK 999	UNP P02786

- Molecule 2 is a protein called Serotransferrin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	515	Total	C	N	O	S	0	0	0
			3650	2275	615	725	35			
2	D	487	Total	C	N	O	S	0	0	0
			3274	1997	569	675	33			

There are 38 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
C	-13	VAL	-	expression tag	UNP P02787
C	-12	PRO	-	expression tag	UNP P02787
C	-11	ASP	-	expression tag	UNP P02787
C	-10	LYS	-	expression tag	UNP P02787
C	-9	HIS	-	expression tag	UNP P02787
C	-8	HIS	-	expression tag	UNP P02787
C	-7	HIS	-	expression tag	UNP P02787
C	-6	HIS	-	expression tag	UNP P02787
C	-5	HIS	-	expression tag	UNP P02787
C	-4	HIS	-	expression tag	UNP P02787
C	-3	ILE	-	expression tag	UNP P02787
C	-2	GLU	-	expression tag	UNP P02787
C	-1	GLY	-	expression tag	UNP P02787
C	0	ARG	-	expression tag	UNP P02787
C	413	ASP	ASN	engineered mutation	UNP P02787
C	426	PHE	TYR	engineered mutation	UNP P02787
C	429	VAL	ILE	SEE REMARK 999	UNP P02787
C	517	PHE	TYR	engineered mutation	UNP P02787
C	611	ASP	ASN	engineered mutation	UNP P02787
D	-13	VAL	-	expression tag	UNP P02787
D	-12	PRO	-	expression tag	UNP P02787
D	-11	ASP	-	expression tag	UNP P02787
D	-10	LYS	-	expression tag	UNP P02787
D	-9	HIS	-	expression tag	UNP P02787
D	-8	HIS	-	expression tag	UNP P02787
D	-7	HIS	-	expression tag	UNP P02787
D	-6	HIS	-	expression tag	UNP P02787

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	-5	HIS	-	expression tag	UNP P02787
D	-4	HIS	-	expression tag	UNP P02787
D	-3	ILE	-	expression tag	UNP P02787
D	-2	GLU	-	expression tag	UNP P02787
D	-1	GLY	-	expression tag	UNP P02787
D	0	ARG	-	expression tag	UNP P02787
D	413	ASP	ASN	engineered mutation	UNP P02787
D	426	PHE	TYR	engineered mutation	UNP P02787
D	429	VAL	ILE	SEE REMARK 999	UNP P02787
D	517	PHE	TYR	engineered mutation	UNP P02787
D	611	ASP	ASN	engineered mutation	UNP P02787

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf			
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	
4	B	1	Total	Ca	0	0
			1	1		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Ca 1 1	0	0

- Molecule 5 is FE (III) ION (three-letter code: FE) (formula: Fe).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	D	1	Total Fe 1 1	0	0
5	C	1	Total Fe 1 1	0	0

- Molecule 6 is CARBONATE ION (three-letter code: CO3) (formula: CO<sub>3</sub>).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	C	1	Total C O 4 1 3	0	0
6	D	1	Total C O 4 1 3	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Transferrin receptor protein 1



#### • Molecule 1: Transferrin receptor protein 1









## 4 Data and refinement statistics

Property	Value	Source
Space group	P 43 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	231.68Å 231.68Å 168.96Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 3.22 29.94 – 3.22	Depositor EDS
% Data completeness (in resolution range)	99.1 (30.00-3.22) 99.5 (29.94-3.22)	Depositor EDS
$R_{merge}$	0.14	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.38 (at 3.24Å)	Xtrriage
Refinement program	CNS	Depositor
R, $R_{free}$	0.271 , 0.314 0.273 , 0.316	Depositor DCC
$R_{free}$ test set	7464 reflections (10.11%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	89.1	Xtrriage
Anisotropy	0.519	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 85.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	16713	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	114.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.28% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CO3, FE, CA, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.42	0/5068	0.60	0/6897
1	B	0.42	0/4916	0.57	1/6718 (0.0%)
2	C	0.34	0/3725	0.52	0/5095
2	D	0.29	0/3327	0.49	0/4546
All	All	0.38	0/17036	0.55	1/23256 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	579	ARG	NE-CZ-NH2	-5.21	117.70	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4949	0	4754	242	0
1	B	4800	0	4434	184	0
2	C	3650	0	3184	121	0
2	D	3274	0	2660	85	0
3	A	14	0	13	0	0
3	B	14	0	13	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	C	1	0	0	0	0
5	D	1	0	0	0	0
6	C	4	0	0	0	0
6	D	4	0	0	3	0
All	All	16713	0	15058	625	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 20.

The worst 5 of 625 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:489:LEU:HD12	1:B:558:CYS:SG	1.56	1.43
1:B:412:TRP:CZ3	1:B:569:THR:HG23	1.52	1.42
1:B:412:TRP:CZ3	1:B:569:THR:CG2	2.21	1.22
2:C:58:ASP:CG	2:C:259:LYS:HZ1	1.49	1.14
1:A:677:ARG:NH1	1:A:751:GLY:HA2	1.63	1.12

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	636/654 (97%)	588 (92%)	43 (7%)	5 (1%)	19 57
1	B	637/654 (97%)	603 (95%)	32 (5%)	2 (0%)	41 74
2	C	509/693 (73%)	464 (91%)	39 (8%)	6 (1%)	13 48
2	D	467/693 (67%)	434 (93%)	29 (6%)	4 (1%)	17 55
All	All	2249/2694 (84%)	2089 (93%)	143 (6%)	17 (1%)	19 57

5 of 17 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	210	VAL
1	A	681	VAL
2	C	125	SER
2	D	613	THR
2	D	629	LEU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	521/562 (93%)	515 (99%)	6 (1%)	71 87
1	B	483/562 (86%)	477 (99%)	6 (1%)	71 87
2	C	353/585 (60%)	349 (99%)	4 (1%)	73 88
2	D	291/585 (50%)	289 (99%)	2 (1%)	84 93
All	All	1648/2294 (72%)	1630 (99%)	18 (1%)	73 88

5 of 18 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	251	ASN
1	B	320	GLN
2	C	241	CYS
1	B	197	GLN
1	B	242	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 38 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	335	GLN
1	B	596	GLN
2	D	585	HIS
1	B	512	ASN
1	B	715	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
6	CO3	C	905	5	0,3,3	0.00	-	0,3,3	0.00	-
3	NAG	B	903	1	14,14,15	0.42	0	17,19,21	1.17	2 (11%)
3	NAG	A	903	1	14,14,15	0.42	0	17,19,21	1.16	2 (11%)
6	CO3	D	905	5	0,3,3	0.00	-	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	B	903	1	-	0/6/23/26	0/1/1/1
3	NAG	A	903	1	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
3	B	903	NAG	C8-C7-N2	2.27	119.94	116.10
3	A	903	NAG	C8-C7-N2	2.27	119.94	116.10
3	B	903	NAG	C2-N2-C7	-2.06	119.97	122.90
3	A	903	NAG	C2-N2-C7	-2.05	119.98	122.90

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	D	905	CO3	3	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	638/654 (97%)	-0.19	9 (1%) 75 64	45, 81, 127, 200	0
1	B	639/654 (97%)	0.13	27 (4%) 36 24	39, 106, 184, 200	1 (0%)
2	C	515/693 (74%)	0.20	23 (4%) 33 21	71, 127, 182, 200	2 (0%)
2	D	487/693 (70%)	0.44	51 (10%) 6 4	68, 150, 199, 200	1 (0%)
All	All	2279/2694 (84%)	0.12	110 (4%) 30 19	39, 111, 185, 200	4 (0%)

The worst 5 of 110 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	195	SER	5.2
2	D	288	PRO	5.2
2	D	12	SER	5.2
2	D	613	THR	5.0
2	C	665	CYS	4.9

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	NAG	A	903	14/15	0.72	0.34	134,134,134,134	0
3	NAG	B	903	14/15	0.80	0.34	154,154,154,154	0
5	FE	D	901	1/1	0.86	0.14	128,128,128,128	0
6	CO3	D	905	4/4	0.93	0.26	163,163,163,163	0
5	FE	C	901	1/1	0.97	0.18	98,98,98,98	0
6	CO3	C	905	4/4	0.98	0.37	78,78,78,78	0
4	CA	B	900	1/1	0.98	0.13	79,79,79,79	0
4	CA	A	900	1/1	0.98	0.23	59,59,59,59	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.