

Full wwPDB X-ray Structure Validation Report (i)

Jun 26, 2024 – 06:39 AM EDT

PDB ID	:	6SCF
Title	:	A viral anti-CRISPR subverts type III CRISPR immunity by rapid degradation
		of cyclic oligoadenylate
Authors	:	McMahon, S.A.; Athukoralage, J.S.; Graham, S.; White, M.F.; Gloster, T.M.
Deposited on	:	2019-07-24
Resolution	:	1.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 1.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\#Entries)$	Similar resolution $(\#\text{Entries}, \text{resolution}, \text{range}(\text{\AA}))$		
	(#Entries)	(#Entries, resolution range(A))		
R _{free}	130704	$1483 \ (1.56-1.56)$		
Clashscore	141614	1529 (1.56-1.56)		
Ramachandran outliers	138981	1498 (1.56-1.56)		
Sidechain outliers	138945	1495 (1.56-1.56)		
RSRZ outliers	127900	1465 (1.56-1.56)		
RNA backbone	3102	1015 (2.36-0.86)		

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	А	138	75% 7%	18%
1	В	138	% 7 6% 6%	6 18%
1	С	138	78%	19%
1	D	138	.% 8 0%	• 19%



Mol	Chain	Length		Quality	of chain		
1	Е	138	3%	76%		••	19%
1	F	138		74%		9%	17%
1	G	138	2%	76%		5%	19%
1	Н	138	3%	75%		•	21%
2	Ι	4	50%		25%		25%
2	K	4	25%	50	9%		25%
2	L	4	50%		25%	_	25%
2	М	4	50%			50%	



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 8312 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace				
1	Δ	119	Total	С	Ν	0	S	0	0	0				
	115	916	587	146	180	3	0	0	0					
1	В	112	Total	С	Ν	0	S	0	4	0				
1	D	110	950	610	151	186	3	0	4	0				
1	C	119	Total	С	Ν	0	S	0	2	0				
		112	928	598	146	181	3	0	0	0				
1	Л	Л	п	Л	Л	119	Total	С	Ν	0	S	S 0	4	0
1	D	112	920	599	145	173	3	0	4	0				
1	F	119	Total	С	Ν	0	S	0	1	0				
1	Ľ	112	902	581	142	176	3		T	0				
1	Б	114	Total	С	Ν	0	S	0	4	0				
	Г	114	950	609	150	187	4	0	4	0				
1	C	119	Total	С	Ν	0	S	0	2	0				
	G	112	917	591	147	176	3	0	2	0				
1	ц	100	Total	С	Ν	0	S	0	0	0				
	п	109	882	570	141	168	3		0	U				

• Molecule 1 is a protein called Uncharacterized protein.

There are 200 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	-23	MET	-	initiating methionine	UNP Q8QL27
А	-22	SER	-	expression tag	UNP Q8QL27
А	-21	TYR	-	expression tag	UNP Q8QL27
А	-20	TYR	-	expression tag	UNP Q8QL27
А	-19	HIS	-	expression tag	UNP Q8QL27
А	-18	HIS	-	expression tag	UNP Q8QL27
А	-17	HIS	-	expression tag	UNP Q8QL27
А	-16	HIS	-	expression tag	UNP Q8QL27
А	-15	HIS	-	expression tag	UNP Q8QL27
А	-14	HIS	-	expression tag	UNP Q8QL27
А	-13	ASP	-	expression tag	UNP Q8QL27
А	-12	TYR	-	expression tag	UNP Q8QL27
A	-11	ASP	-	expression tag	UNP Q8QL27



Chain	Residue	Modelled	Actual	Comment	Reference
А	-10	ILE	-	expression tag	UNP Q8QL27
А	-9	PRO	-	expression tag	UNP Q8QL27
А	-8	THR	-	expression tag	UNP Q8QL27
А	-7	THR	-	expression tag	UNP Q8QL27
А	-6	GLU	-	expression tag	UNP Q8QL27
А	-5	ASN	-	expression tag	UNP Q8QL27
А	-4	LEU	-	expression tag	UNP Q8QL27
А	-3	TYR	-	expression tag	UNP Q8QL27
А	-2	PHE	-	expression tag	UNP Q8QL27
А	-1	GLN	-	expression tag	UNP Q8QL27
А	0	GLY	-	expression tag	UNP Q8QL27
А	47	ALA	HIS	conflict	UNP Q8QL27
В	-23	MET	-	initiating methionine	UNP Q8QL27
В	-22	SER	-	expression tag	UNP Q8QL27
В	-21	TYR	-	expression tag	UNP Q8QL27
В	-20	TYR	_	expression tag	UNP Q8QL27
В	-19	HIS	-	expression tag	UNP Q8QL27
В	-18	HIS	-	expression tag	UNP Q8QL27
В	-17	HIS	_	expression tag	UNP Q8QL27
В	-16	HIS	-	expression tag	UNP Q8QL27
В	-15	HIS	_	expression tag	UNP Q8QL27
В	-14	HIS	-	expression tag	UNP Q8QL27
В	-13	ASP	-	expression tag	UNP Q8QL27
В	-12	TYR	_	expression tag	UNP Q8QL27
В	-11	ASP	-	expression tag	UNP Q8QL27
В	-10	ILE	_	expression tag	UNP Q8QL27
В	-9	PRO	_	expression tag	UNP Q8QL27
В	-8	THR	-	expression tag	UNP Q8QL27
В	-7	THR	-	expression tag	UNP Q8QL27
В	-6	GLU	-	expression tag	UNP Q8QL27
В	-5	ASN	-	expression tag	UNP Q8QL27
В	-4	LEU	-	expression tag	UNP Q8QL27
В	-3	TYR	_	expression tag	UNP Q8QL27
В	-2	PHE	_	expression tag	UNP Q8QL27
В	-1	GLN	-	expression tag	UNP Q8QL27
В	0	GLY	_	expression tag	UNP Q8QL27
В	47	ALA	HIS	conflict	UNP Q8QL27
С	-23	MET	-	initiating methionine	UNP Q8QL27
С	-22	SER	-	expression tag	UNP Q8QL27
С	-21	TYR	-	expression tag	UNP Q8QL27
С	-20	TYR	-	expression tag	UNP Q8QL27
С	-19	HIS	-	expression tag	UNP Q8QL27



С

Actual

-	expression tag	UNP Q8QL27
-	expression tag	UNP Q8QL27
-	expression tag	UNP Q8QL27
-	expression tag	UNP Q8QL27
-	expression tag	UNP Q8QL27
-	expression tag	UNP Q8QL27
-	expression tag	UNP Q8QL27
-	expression tag	UNP Q8QL27
-	expression tag	UNP Q8QL27
-	expression tag	UNP Q8QL27
-	expression tag	UNP Q8QL27
-	expression tag	UNP Q8QL27
-	expression tag	UNP Q8QL27
-	expression tag	UNP Q8QL27
-	expression tag	UNP Q8QL27
-	expression tag	UNP Q8QL27
-	expression tag	UNP Q8QL27
-	expression tag	UNP Q8QL27
-	expression tag	UNP Q8QL27
HIS	conflict	UNP $080L27$

Comment

Continued from previous page... Chain Residue Modelled

HIS

-18

	1				
С	-17	HIS	-	expression tag	UNP Q8QL27
С	-16	HIS	-	expression tag	UNP Q8QL27
С	-15	HIS	-	expression tag	UNP Q8QL27
С	-14	HIS	-	expression tag	UNP Q8QL27
С	-13	ASP	-	expression tag	UNP Q8QL27
С	-12	TYR	-	expression tag	UNP Q8QL27
С	-11	ASP	-	expression tag	UNP Q8QL27
С	-10	ILE	-	expression tag	UNP Q8QL27
С	-9	PRO	-	expression tag	UNP Q8QL27
С	-8	THR	-	expression tag	UNP Q8QL27
С	-7	THR	-	expression tag	UNP Q8QL27
С	-6	GLU	-	expression tag	UNP Q8QL27
С	-5	ASN	-	expression tag	UNP Q8QL27
С	-4	LEU	-	expression tag	UNP Q8QL27
С	-3	TYR	-	expression tag	UNP Q8QL27
С	-2	PHE	-	expression tag	UNP Q8QL27
С	-1	GLN	-	expression tag	UNP Q8QL27
С	0	GLY	-	expression tag	UNP Q8QL27
С	47	ALA	HIS	conflict	UNP Q8QL27
D	-23	MET	-	initiating methionine	UNP Q8QL27
D	-22	SER	-	expression tag	UNP Q8QL27
D	-21	TYR	-	expression tag	UNP Q8QL27
D	-20	TYR	-	expression tag	UNP Q8QL27
D	-19	HIS	-	expression tag	UNP Q8QL27
D	-18	HIS	-	expression tag	UNP Q8QL27
D	-17	HIS	-	expression tag	UNP Q8QL27
D	-16	HIS	-	expression tag	UNP Q8QL27
D	-15	HIS	-	expression tag	UNP Q8QL27
D	-14	HIS	-	expression tag	UNP Q8QL27
D	-13	ASP	-	expression tag	UNP Q8QL27
D	-12	TYR	-	expression tag	UNP Q8QL27
D	-11	ASP	-	expression tag	UNP Q8QL27
D	-10	ILE	-	expression tag	UNP Q8QL27
D	-9	PRO	-	expression tag	UNP Q8QL27
D	-8	THR	-	expression tag	UNP Q8QL27
D	-7	THR	-	expression tag	UNP Q8QL27
D	-6	GLU	-	expression tag	UNP Q8QL27
D	-5	ASN	-	expression tag	UNP Q8QL27
D	-4	LEU	-	expression tag	UNP Q8QL27
D	-3	TYR	-	expression tag	UNP Q8QL27
D	-2	PHE	-	expression tag	UNP Q8QL27

Continued on next page...

Reference



6SCF

Chain	Residue	Modelled	Actual	Comment	Reference
D	-1	GLN	-	expression tag	UNP Q8QL27
D	0	GLY	-	expression tag	UNP Q8QL27
D	47	ALA	HIS	conflict	UNP Q8QL27
Е	-23	MET	-	initiating methionine	UNP Q8QL27
Е	-22	SER	-	expression tag	UNP Q8QL27
Е	-21	TYR	-	expression tag	UNP Q8QL27
Е	-20	TYR	-	expression tag	UNP Q8QL27
Е	-19	HIS	-	expression tag	UNP Q8QL27
Е	-18	HIS	-	expression tag	UNP Q8QL27
Е	-17	HIS	-	expression tag	UNP Q8QL27
Е	-16	HIS	-	expression tag	UNP Q8QL27
Е	-15	HIS	-	expression tag	UNP Q8QL27
Е	-14	HIS	-	expression tag	UNP Q8QL27
Е	-13	ASP	-	expression tag	UNP Q8QL27
Е	-12	TYR	-	expression tag	UNP Q8QL27
Е	-11	ASP	-	expression tag	UNP Q8QL27
Е	-10	ILE	-	expression tag	UNP Q8QL27
Е	-9	PRO	-	expression tag	UNP Q8QL27
Е	-8	THR	-	expression tag	UNP Q8QL27
Е	-7	THR	-	expression tag	UNP Q8QL27
Е	-6	GLU	-	expression tag	UNP Q8QL27
Е	-5	ASN	-	expression tag	UNP Q8QL27
Е	-4	LEU	-	expression tag	UNP Q8QL27
Е	-3	TYR	-	expression tag	UNP Q8QL27
Е	-2	PHE	-	expression tag	UNP Q8QL27
Е	-1	GLN	-	expression tag	UNP Q8QL27
Е	0	GLY	-	expression tag	UNP Q8QL27
Е	47	ALA	HIS	conflict	UNP Q8QL27
F	-23	MET	-	initiating methionine	UNP Q8QL27
F	-22	SER	-	expression tag	UNP Q8QL27
F	-21	TYR	-	expression tag	UNP Q8QL27
F	-20	TYR	-	expression tag	UNP Q8QL27
F	-19	HIS	-	expression tag	UNP Q8QL27
F	-18	HIS	-	expression tag	UNP Q8QL27
F	-17	HIS	-	expression tag	UNP Q8QL27
F	-16	HIS	-	expression tag	UNP Q8QL27
F	-15	HIS	-	expression tag	UNP Q8QL27
F	-14	HIS	-	expression tag	UNP Q8QL27
F	-13	ASP	-	expression tag	UNP Q8QL27
F	-12	TYR	-	expression tag	UNP Q8QL27
F	-11	ASP	-	expression tag	UNP Q8QL27
F	-10	ILE	-	expression tag	UNP Q8QL27



6S	CF

Chain	Residue	Modelled	Actual	Comment	Reference
F	-9	PRO	-	expression tag	UNP Q8QL27
F	-8	THR	_	expression tag	UNP Q8QL27
F	-7	THR	-	expression tag	UNP Q8QL27
F	-6	GLU	-	expression tag	UNP Q8QL27
F	-5	ASN	-	expression tag	UNP Q8QL27
F	-4	LEU	-	expression tag	UNP Q8QL27
F	-3	TYR	-	expression tag	UNP Q8QL27
F	-2	PHE	-	expression tag	UNP Q8QL27
F	-1	GLN	-	expression tag	UNP Q8QL27
F	0	GLY	-	expression tag	UNP Q8QL27
F	47	ALA	HIS	conflict	UNP Q8QL27
G	-23	MET	-	initiating methionine	UNP Q8QL27
G	-22	SER	-	expression tag	UNP Q8QL27
G	-21	TYR	-	expression tag	UNP Q8QL27
G	-20	TYR	-	expression tag	UNP Q8QL27
G	-19	HIS	-	expression tag	UNP Q8QL27
G	-18	HIS	-	expression tag	UNP Q8QL27
G	-17	HIS	-	expression tag	UNP Q8QL27
G	-16	HIS	-	expression tag	UNP Q8QL27
G	-15	HIS	-	expression tag	UNP Q8QL27
G	-14	HIS	-	expression tag	UNP Q8QL27
G	-13	ASP	-	expression tag	UNP Q8QL27
G	-12	TYR	-	expression tag	UNP Q8QL27
G	-11	ASP	-	expression tag	UNP Q8QL27
G	-10	ILE	-	expression tag	UNP Q8QL27
G	-9	PRO	-	expression tag	UNP Q8QL27
G	-8	THR	-	expression tag	UNP Q8QL27
G	-7	THR	-	expression tag	UNP Q8QL27
G	-6	GLU	-	expression tag	UNP Q8QL27
G	-5	ASN	-	expression tag	UNP Q8QL27
G	-4	LEU	-	expression tag	UNP Q8QL27
G	-3	TYR	-	expression tag	UNP Q8QL27
G	-2	PHE	-	expression tag	UNP Q8QL27
G	-1	GLN	-	expression tag	UNP Q8QL27
G	0	GLY	-	expression tag	UNP Q8QL27
G	47	ALA	HIS	conflict	UNP Q8QL27
Н	-23	MET	-	initiating methionine	UNP Q8QL27
Н	-22	SER	-	expression tag	UNP Q8QL27
Н	-21	TYR	-	expression tag	UNP Q8QL27
Н	-20	TYR	-	expression tag	$\overline{\text{UNP Q8QL27}}$
Н	-19	HIS	-	expression tag	UNP Q8QL27
Н	-18	HIS	-	expression tag	UNP Q8QL27



Chain	Residue	Modelled	Actual	Comment	Reference
Н	-17	HIS	-	expression tag	UNP Q8QL27
Н	-16	HIS	-	expression tag	UNP Q8QL27
Н	-15	HIS	-	expression tag	UNP Q8QL27
Н	-14	HIS	-	expression tag	UNP Q8QL27
Н	-13	ASP	-	expression tag	UNP Q8QL27
Н	-12	TYR	-	expression tag	UNP Q8QL27
Н	-11	ASP	-	expression tag	UNP Q8QL27
Н	-10	ILE	-	expression tag	UNP Q8QL27
Н	-9	PRO	-	expression tag	UNP Q8QL27
Н	-8	THR	-	expression tag	UNP Q8QL27
Н	-7	THR	-	expression tag	UNP Q8QL27
Н	-6	GLU	-	expression tag	UNP Q8QL27
Н	-5	ASN	-	expression tag	UNP Q8QL27
Н	-4	LEU	-	expression tag	UNP Q8QL27
Н	-3	TYR	-	expression tag	UNP Q8QL27
Н	-2	PHE	-	expression tag	UNP Q8QL27
Н	-1	GLN	-	expression tag	UNP Q8QL27
Н	0	GLY	-	expression tag	UNP Q8QL27
Н	47	ALA	HIS	conflict	UNP Q8QL27

• Molecule 2 is a RNA chain called cyclic oligoadenylate.

Mol	Chain	Residues		Atoms			ZeroOcc	AltConf	Trace	
9	K	4	Total	С	Ν	Ο	Р	0	0	0
	IX	4	88	40	20	24	4	0	0	0
9	Т	4	Total	С	Ν	Ο	Р	0	0	0
	1	4	88	40	20	24	4			
0	т	4	Total	С	Ν	Ο	Р	0	0	0
		4	88	40	20	24	4	0	0	U
0	9 M	ЛА	Total	С	Ν	Ο	Р	0	0	0
Z M	4	88	40	20	24	4	U	0	0	

• Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	80	Total O 80 80	0	0
3	В	77	Total O 78 78	0	1
3	С	66	Total O 66 66	0	0
3	D	51	$\begin{array}{cc} \text{Total} & \text{O} \\ 51 & 51 \end{array}$	0	0

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	Е	56	Total O 56 56	0	0
3	F	110	Total O 111 111	0	1
3	G	75	Total O 75 75	0	0
3	Н	41	Total O 41 41	0	0
3	К	10	Total O 10 10	0	0
3	Ι	8	Total O 8 8	0	0
3	L	9	Total O 10 10	0	1
3	М	9	Total O 9 9	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Uncharacterized protein

• Molecule 1: Uncharacterized protein



Chain F:	74%	9%	17%
MET SER TYR TYR HIS HIS HIS HIS	ASP TYR ASP ASP THR THR THR ASN ASN ASN ASN ASN CLU CLU CLU CLU CLU CLU CLU CLU CLU CLU	124 D25 K70 K70 8102 8102 8102 8102 8102 8102 8103 8103 81111 1111 11112 1114	
• Molecule 1:	Uncharacterized protein		
Chain G:	76%	5%	19%
MET SER TYR TYR HIS HIS HIS HIS HIS	ASP TYR TYR TILE TILE TILE CLU CLU CLU CLU CLU CLU CLU CLU CLU CLU	D37 N38 E39 E39 E39 E39 E46 I46 K76 I111 I111 I111 X76 K76 K76 K76 K76 K76 K76 K76 K76 K76 K	
• Molecule 1:	Uncharacterized protein		
Chain H:	75%	·	21%
MET SER TYR TYR HIS HIS HIS HIS	ASP TYR TYR THR THR THR ASN CLU CLU CLU MET MET CLN CLN CLN CLN CLN CLN CLN CLN CLN	136 D37 ASN ASN ASP ASP ASP ASP B10 B102 E102 E102	V113 ASP
• Molecule 2:	cyclic oligoadenylate		
Chain K:	25%	50%	25%
A1 A3 A4			
• Molecule 2:	cyclic oligoadenylate		
Chain I:	50%	25%	25%
A1 A2 A3 A4			
• Molecule 2:	cyclic oligoadenylate		
Chain L:	50%	25%	25%
A1 A2 A3 A4			
• Molecule 2:	cyclic oligoadenylate		
Chain M:	50%	50%	
A1 A3 A4			



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1	Depositor
Cell constants	49.83Å 51.73Å 85.61Å	Depositor
a, b, c, α , β , γ	80.22° 89.68° 83.38°	Depositor
Bosolution(A)	50.64 - 1.55	Depositor
Resolution (A)	50.63 - 1.55	EDS
% Data completeness	98.6 (50.64-1.55)	Depositor
(in resolution range)	98.6(50.63-1.55)	EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$2.11 (at 1.55 \text{\AA})$	Xtriage
Refinement program	REFMAC 5.8.0218	Depositor
P. P.	0.202 , 0.249	Depositor
n, n_{free}	0.210 , 0.255	DCC
R_{free} test set	5883 reflections (4.91%)	wwPDB-VP
Wilson B-factor $(Å^2)$	13.7	Xtriage
Anisotropy	0.250	Xtriage
Bulk solvent $k_{sol}(e/A^3)$, $B_{sol}(A^2)$	0.37, 45.8	EDS
L-test for twinning ²	$ < L >=0.49, < L^2>=0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8312	wwPDB-VP
Average B, all atoms $(Å^2)$	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 82.38 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.8279e-07. The detected translational NCS is most likely also responsible for the elevated intensity ratio.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bo	nd lengths	Bo	ond angles
	Ullaili	RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	0.65	0/925	0.78	0/1243
1	В	0.65	0/959	0.79	0/1288
1	С	0.67	0/937	0.76	0/1262
1	D	0.60	0/932	0.77	0/1256
1	Е	0.69	0/911	0.81	1/1226~(0.1%)
1	F	0.70	0/959	0.84	1/1291~(0.1%)
1	G	0.59	0/926	0.77	0/1246
1	Н	0.64	1/890~(0.1%)	0.72	0/1195
2	Ι	0.82	0/99	1.51	1/152~(0.7%)
2	Κ	0.72	0/99	1.53	1/152~(0.7%)
2	L	0.89	0/99	1.48	2/152~(1.3%)
2	М	0.95	0/99	1.50	1/152~(0.7%)
All	All	0.66	$1/\overline{7835}\ (0.0\%)$	0.84	7/10615~(0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	С	0	1
2	L	0	1
All	All	0	2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	$\operatorname{Ideal}(\operatorname{\AA})$
1	Н	110	GLU	CD-OE1	-5.71	1.19	1.25

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	М	2	A	O4'-C1'-N9	-6.73	102.81	108.20



Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	Κ	2	А	O4'-C1'-N9	-6.48	103.02	108.20
2	L	4	А	O4'-C1'-N9	-6.19	103.25	108.20
2	L	1	А	O5'-P-OP2	-6.04	100.26	105.70
1	F	70	LYS	CD-CE-NZ	5.49	124.33	111.70
2	Ι	3	А	O4'-C1'-N9	-5.33	103.94	108.20
1	Е	30	LEU	CA-CB-CG	5.23	127.32	115.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	С	10	PHE	Mainchain
2	L	4	A	Sidechain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	916	0	942	14	0
1	В	950	0	979	7	0
1	С	928	0	949	2	0
1	D	920	0	956	2	0
1	Е	902	0	919	7	0
1	F	950	0	972	12	0
1	G	917	0	947	5	0
1	Н	882	0	921	4	0
2	Ι	88	0	44	0	0
2	K	88	0	44	0	0
2	L	88	0	44	2	0
2	М	88	0	44	2	0
3	А	80	0	0	0	0
3	В	78	0	0	0	0
3	С	66	0	0	0	0
3	D	51	0	0	0	0
3	Е	56	0	0	1	0
3	F	111	0	0	4	0
3	G	75	0	0	2	0



0 0							
Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes	
3	Н	41	0	0	1	0	
3	Ι	8	0	0	0	0	
3	Κ	10	0	0	0	0	
3	L	10	0	0	0	0	
3	М	9	0	0	0	0	
All	All	8312	0	7761	41	0	

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:A:34:GLU:OE2	1:F:112:ILE:HD13	1.98	0.64
1:B:12:ILE:HD11	1:G:111:ILE:HD11	1.78	0.64
1:H:29:ARG:HE	1:H:57:LEU:HD22	1.68	0.58
1:A:31:GLU:CG	3:F:218:HOH:O	2.53	0.56
1:D:12:ILE:HD11	1:F:111:ILE:HD11	1.87	0.56
1:A:31:GLU:HG3	3:F:218:HOH:O	2.06	0.55
1:F:22[A]:VAL:HG12	1:F:113:VAL:HG12	1.89	0.55
1:G:29:ARG:HE	1:G:57:LEU:HD22	1.72	0.54
1:B:39:GLU:H	1:B:39:GLU:CD	2.12	0.53
1:G:2:ASN:N	3:G:201:HOH:O	2.42	0.53
1:C:111[A]:ILE:HD11	1:E:12:ILE:HD11	1.91	0.52
1:E:21:LYS:HG3	1:E:114:ASP:HA	1.92	0.51
1:A:42:ILE:CD1	1:E:103:SER:HB3	2.41	0.51
3:E:225:HOH:O	2:L:2:A:H1'	2.12	0.50
1:A:30:LEU:HD21	1:F:23:VAL:CG1	2.42	0.50
1:B:29:ARG:HD2	1:B:101:TYR:OH	2.13	0.49
1:C:76:LYS:HE3	1:C:110:GLU:OE2	2.13	0.48
1:A:30:LEU:HD21	1:F:23:VAL:HG12	1.96	0.48
1:B:29:ARG:HE	1:B:57:LEU:HD22	1.78	0.48
1:F:102:GLU:CD	3:F:201:HOH:O	2.52	0.47
1:A:28:ASP:OD2	1:F:25:ASP:OD2	2.32	0.46
1:A:39:GLU:H	1:A:39:GLU:CD	2.19	0.46
1:H:81:GLN:HA	3:H:212:HOH:O	2.16	0.45
2:L:2:A:H2'	2:L:2:A:N3	2.32	0.45
1:G:8:ASN:N	1:G:45[B]:ILE:HD13	2.31	0.44
1:A:34:GLU:CD	1:F:112:ILE:HG21	2.38	0.44
1:B:73:LYS:HG3	1:B:114:ASP:HB3	2.00	0.44
1:A:57:LEU:HD23	1:H:30:LEU:HD23	2.01	0.43



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:42:ILE:HG12	1:F:103[A]:SER:HB2	2.01	0.43
2:M:4:A:H2'	2:M:4:A:N3	2.34	0.43
1:D:83:SER:HA	1:F:13:ASN:OD1	2.18	0.43
3:F:212:HOH:O	2:M:4:A:H1'	2.19	0.43
1:A:39:GLU:HG3	1:E:85:ARG:O	2.18	0.43
1:G:76:LYS:HG3	3:G:245:HOH:O	2.18	0.42
1:E:29:ARG:HH22	1:E:30:LEU:HD23	1.85	0.42
1:A:73:LYS:HG3	1:A:114:ASP:HB3	2.02	0.41
1:B:42:ILE:HD11	1:F:103[A]:SER:HB3	2.02	0.41
1:E:29:ARG:NH2	1:E:30:LEU:HD23	2.36	0.41
1:A:42:ILE:HD11	1:E:103:SER:HB3	2.03	0.41
1:F:84:GLN:OE1	1:F:105:LYS:HD3	2.20	0.41
1:A:57:LEU:HD23	1:H:30:LEU:CD2	2.52	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	111/138 (80%)	110 (99%)	1 (1%)	0	100 100
1	В	115/138~(83%)	114 (99%)	1 (1%)	0	100 100
1	С	113/138~(82%)	112 (99%)	1 (1%)	0	100 100
1	D	114/138~(83%)	113~(99%)	1 (1%)	0	100 100
1	Е	111/138 (80%)	110 (99%)	1 (1%)	0	100 100
1	F	116/138~(84%)	112~(97%)	4(3%)	0	100 100
1	G	112/138~(81%)	110 (98%)	2(2%)	0	100 100
1	Н	105/138~(76%)	103 (98%)	2 (2%)	0	100 100
All	All	897/1104 (81%)	884 (99%)	13 (1%)	0	100 100



There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	А	106/130~(82%)	105~(99%)	1 (1%)	78	61
1	В	110/130~(85%)	110 (100%)	0	100	100
1	С	106/130~(82%)	106 (100%)	0	100	100
1	D	104/130~(80%)	104 (100%)	0	100	100
1	Ε	102/130~(78%)	102 (100%)	0	100	100
1	F	110/130~(85%)	110 (100%)	0	100	100
1	G	105/130~(81%)	105 (100%)	0	100	100
1	Н	102/130~(78%)	102 (100%)	0	100	100
All	All	845/1040 (81%)	844 (100%)	1 (0%)	93	86

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	29	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	А	13	ASN

5.3.3 RNA (i)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	Ι	3/4~(75%)	2~(66%)	0
2	Κ	3/4~(75%)	3~(100%)	0
2	L	3/4~(75%)	3 (100%)	0
2	М	3/4~(75%)	2(66%)	0



Continued from previous page...

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
All	All	12/16~(75%)	10 (83%)	0

All (10) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	Κ	2	А
2	Κ	3	А
2	Κ	4	А
2	Ι	2	А
2	Ι	3	А
2	L	2	А
2	L	3	А
2	L	4	А
2	М	2	А
2	М	4	А

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle RSRZ \rangle$	#RSRZ>2	$\mathbf{OWAB}(\mathbf{A}^2)$	Q<0.9
1	А	113/138 (81%)	-0.01	0 100 100	8, 16, 28, 38	0
1	В	113/138~(81%)	0.07	1 (0%) 84 87	10, 16, 29, 34	0
1	С	112/138~(81%)	0.32	4 (3%) 42 50	11, 21, 33, 43	0
1	D	112/138~(81%)	0.33	1 (0%) 84 87	11, 22, 36, 45	0
1	Ε	112/138~(81%)	0.26	4 (3%) 42 50	9, 19, 45, 56	0
1	F	114/138 (82%)	0.11	0 100 100	10, 16, 30, 44	0
1	G	112/138~(81%)	0.14	3 (2%) 54 62	10, 19, 32, 39	0
1	Н	109/138~(78%)	0.28	4 (3%) 41 48	12, 24, 37, 45	0
2	Ι	4/4~(100%)	-0.52	0 100 100	11, 12, 13, 13	0
2	Κ	4/4~(100%)	-0.43	0 100 100	13, 13, 15, 16	0
2	L	4/4~(100%)	-0.40	0 100 100	11, 14, 14, 16	0
2	М	4/4 (100%)	-0.51	0 100 100	12, 12, 13, 15	0
All	All	913/1120 (81%)	0.18	17 (1%) 66 73	8, 18, 33, 56	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	С	102	GLU	3.8
1	Е	74	GLU	3.8
1	С	101	TYR	3.4
1	G	37	ASP	3.2
1	Е	75	ASP	3.2
1	С	98	LEU	3.0
1	G	38	ASN	2.8
1	D	91[A]	ILE	2.6
1	G	40	ASP	2.5
1	Н	37	ASP	2.4
1	Е	40	ASP	2.4



Mol	Chain	Res	Type	RSRZ
1	Н	102	GLU	2.3
1	С	39	GLU	2.2
1	Н	98	LEU	2.2
1	В	38	ASN	2.1
1	Е	113	VAL	2.1
1	Н	36	ILE	2.0

Continued from previous page...

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

There are no ligands in this entry.

6.5 Other polymers (i)

There are no such residues in this entry.

