



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 15, 2021 – 03:02 PM GMT

PDB ID : 6SJH
Title : Structure of the PRY-SPRY domain of human Trim16L/Trim70
Authors : Mikkelsen, J.H.; Andersen, C.B.F.
Deposited on : 2019-08-13
Resolution : 1.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtrriage (Phenix) : 1.13
EDS : **FAILED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.17


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.




Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	3144 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Note EDS failed to run properly.

Mol	Chain	Length	Quality of chain
1	A	190	 87% 12% .
1	B	190	 87% 10% .
1	C	190	 92% 8% .

2 Entry composition i

There are 2 unique types of molecules in this entry. The entry contains 5410 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tripartite motif-containing protein 16-like protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	187	Total 1591	C 1033	N 265	O 285	S 8	0	6	0
1	B	185	Total 1584	C 1030	N 263	O 284	S 7	0	7	0
1	C	189	Total 1598	C 1035	N 265	O 292	S 6	0	4	0

There are 27 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	143	GLY	-	expression tag	UNP Q309B1
A	144	SER	-	expression tag	UNP Q309B1
A	145	HIS	-	expression tag	UNP Q309B1
A	146	MET	-	expression tag	UNP Q309B1
A	171	GLN	ARG	conflict	UNP Q309B1
A	235	ARG	GLN	conflict	UNP Q309B1
A	241	ASN	SER	conflict	UNP Q309B1
A	277	ARG	TRP	conflict	UNP Q309B1
A	299	THR	SER	conflict	UNP Q309B1
B	143	GLY	-	expression tag	UNP Q309B1
B	144	SER	-	expression tag	UNP Q309B1
B	145	HIS	-	expression tag	UNP Q309B1
B	146	MET	-	expression tag	UNP Q309B1
B	171	GLN	ARG	conflict	UNP Q309B1
B	235	ARG	GLN	conflict	UNP Q309B1
B	241	ASN	SER	conflict	UNP Q309B1
B	277	ARG	TRP	conflict	UNP Q309B1
B	299	THR	SER	conflict	UNP Q309B1
C	143	GLY	-	expression tag	UNP Q309B1
C	144	SER	-	expression tag	UNP Q309B1
C	145	HIS	-	expression tag	UNP Q309B1
C	146	MET	-	expression tag	UNP Q309B1
C	171	GLN	ARG	conflict	UNP Q309B1

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	235	ARG	GLN	conflict	UNP Q309B1
C	241	ASN	SER	conflict	UNP Q309B1
C	277	ARG	TRP	conflict	UNP Q309B1
C	299	THR	SER	conflict	UNP Q309B1

- Molecule 2 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	240	Total O 240 240	0	0
2	B	169	Total O 169 169	0	0
2	C	228	Total O 228 228	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


Note EDS failed to run properly.

- Molecule 1: Tripartite motif-containing protein 16-like protein

Chain A: 




- Molecule 1: Tripartite motif-containing protein 16-like protein

Chain B: 



- Molecule 1: Tripartite motif-containing protein 16-like protein

Chain C: 



4 Data and refinement statistics

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	72.78Å 95.77Å 116.09Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 1.50	Depositor
% Data completeness (in resolution range)	88.3 (30.00-1.50)	Depositor
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.96 (at 1.14Å)	Xtrriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.181 , 0.212	Depositor
Wilson B-factor (Å ²)	21.5	Xtrriage
Anisotropy	0.051	Xtrriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	5410	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.79% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.33	0/1661	0.58	0/2252
1	B	0.34	0/1656	0.57	0/2246
1	C	0.34	0/1662	0.57	0/2255
All	All	0.34	0/4979	0.58	0/6753

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1591	0	1530	17	0
1	B	1584	0	1529	17	0
1	C	1598	0	1521	17	0
2	A	240	0	0	8	2
2	B	169	0	0	5	1
2	C	228	0	0	7	1
All	All	5410	0	4580	51	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 51 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:331:ASP:OD1	2:A:401:HOH:O	1.91	0.87
1:B:230[B]:CYS:SG	2:B:442:HOH:O	2.34	0.86
1:A:298:ASP:OD2	2:A:402:HOH:O	1.92	0.86
1:A:331:ASP:O	2:A:404:HOH:O	2.01	0.79
1:A:311[B]:SER:OG	2:A:403:HOH:O	2.00	0.78

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:567:HOH:O	2:B:526:HOH:O[3_554]	1.79	0.41
2:A:620:HOH:O	2:C:581:HOH:O[3_554]	2.01	0.19

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

There are no protein backbone outliers to report in this entry.

5.3.2 Protein sidechains [i](#)

There are no protein residues with a non-rotameric sidechain to report in this entry.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS failed to run properly - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS failed to run properly - this section is therefore empty.

6.3 Carbohydrates

EDS failed to run properly - this section is therefore empty.

6.4 Ligands

EDS failed to run properly - this section is therefore empty.

6.5 Other polymers

EDS failed to run properly - this section is therefore empty.