



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 16, 2024 – 04:50 PM EDT

PDB ID : 8SJR
Title : [3T17] Self-assembling right-handed tensegrity triangle with 17 interjunction base pairs and R3 symmetry
Authors : Janowski, J.; Vecchioni, S.; Sha, R.; Ohayon, Y.P.
Deposited on : 2023-04-18
Resolution : 5.25 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtrriage (Phenix) : 1.13
EDS : 2.36.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 5.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1175 (6.72-3.80)
Clashscore	141614	1012 (6.64-3.84)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	19	
2	B	17	
3	C	14	
4	D	12	

2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 1265 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (5'-D(*CP*AP*GP*CP*AP*GP*CP*CP*TP*GP*AP*AP*TP*AP*CP*CP*GP*CP*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
1	A	19	384	183	75	108	18	0	0	0

- Molecule 2 is a DNA chain called DNA (5'-D(P*GP*CP*GP*GP*TP*AP*TP*TP*CP*AP*CP*CP*AP*CP*GP*AP*T)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	B	17	347	165	63	102	17	0	0	0

- Molecule 3 is a DNA chain called DNA (5'-D(*TP*GP*CP*GP*CP*TP*GP*TP*GP*GP*CP*TP*GP*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
3	C	14	285	136	50	86	13	0	0	0

- Molecule 4 is a DNA chain called DNA (5'-D(P*TP*CP*GP*TP*GP*GP*AP*CP*AP*GP*CP*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
4	D	12	249	117	48	72	12	0	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA (5'-D(*CP*AP*GP*CP*AP*GP*CP*CP*TP*GP*AP*AP*TP*AP*CP*CP*GP*CP*A)-3')

Chain A:  16% 84%



- Molecule 2: DNA (5'-D(P*GP*CP*GP*GP*TP*AP*TP*TP*CP*AP*CP*CP*AP*CP*GP*A P*T)-3')

Chain B:  35% 59% 6%





- Molecule 3: DNA (5'-D(*TP*GP*CP*GP*CP*TP*GP*TP*GP*GP*CP*TP*GP*C)-3')

Chain C:  50% 43% 7%



- Molecule 4: DNA (5'-D(P*TP*CP*GP*TP*GP*GP*AP*CP*AP*GP*CP*G)-3')

Chain D:  25% 75%



4 Data and refinement statistics i

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	170.21Å 170.21Å 80.25Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	38.72 – 5.25 85.11 – 5.25	Depositor EDS
% Data completeness (in resolution range)	55.9 (38.72-5.25) 48.3 (85.11-5.25)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.21 (at 5.12Å)	Xtrriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.209 , 0.250 0.205 , 0.242	Depositor DCC
R_{free} test set	105 reflections (5.63%)	wwPDB-VP
Wilson B-factor (Å ²)	306.9	Xtrriage
Anisotropy	1.166	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.07 , -50.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.065 for h,-h-k,-l	Xtrriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	1265	wwPDB-VP
Average B, all atoms (Å ²)	407.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.74% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.93	0/431	1.10	0/662
2	B	0.83	0/388	1.10	1/596 (0.2%)
3	C	0.95	0/318	1.15	2/490 (0.4%)
4	D	0.79	0/279	1.06	0/429
All	All	0.88	0/1416	1.10	3/2177 (0.1%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	105	DT	N3-C4-O4	5.18	123.01	119.90
3	C	112	DT	N3-C4-O4	5.08	122.95	119.90
3	C	108	DT	N3-C4-O4	5.05	122.93	119.90

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	384	0	213	17	0
2	B	347	0	192	15	0
3	C	285	0	160	10	0
4	D	249	0	135	9	0
All	All	1265	0	700	46	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 23.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:106:DT:H2''	3:C:107:DG:C8	2.02	0.94
1:A:103:DG:H2''	1:A:104:DC:H5''	1.57	0.85
2:B:103:DA:H2''	2:B:104:DT:H5''	1.59	0.83
1:A:110:DG:H2'	1:A:111:DA:C8	2.13	0.83
3:C:109:DG:H2'	3:C:110:DG:C8	2.19	0.77
2:B:111:DC:H2''	2:B:112:DG:H5''	1.77	0.67
3:C:110:DG:H2''	3:C:111:DC:H5'	1.76	0.66
2:B:108:DC:H2'	2:B:109:DC:C5	2.33	0.63
4:D:111:DC:H2''	4:D:112:DG:C8	2.34	0.63
2:B:109:DC:H2''	2:B:110:DA:C8	2.35	0.61
2:B:109:DC:H6	2:B:109:DC:H5''	1.67	0.60
1:A:102:DA:H1'	1:A:103:DG:OP2	2.05	0.57
1:A:104:DC:H2''	1:A:105:DA:H8	1.70	0.56
2:B:112:DG:H2''	2:B:113:DA:C8	2.40	0.56
4:D:117:DC:H2''	4:D:118:DA:H5''	1.87	0.56
1:A:104:DC:H2''	1:A:105:DA:H5'	1.91	0.52
1:A:106:DG:C2	1:A:107:DC:C2	2.98	0.52
4:D:115:DG:H2''	4:D:116:DA:O5'	2.12	0.50
1:A:104:DC:H2''	1:A:105:DA:C8	2.48	0.49
4:D:115:DG:H2''	4:D:116:DA:C5'	2.44	0.47
2:B:108:DC:C4'	3:C:108:DT:H2''	2.44	0.47
1:A:112:DA:H8	1:A:112:DA:OP1	1.98	0.46
1:A:110:DG:H2'	1:A:111:DA:N9	2.31	0.46
3:C:106:DT:C2'	3:C:107:DG:C8	2.88	0.46
2:B:108:DC:O4'	3:C:108:DT:H2''	2.16	0.46
4:D:115:DG:H4'	4:D:116:DA:OP1	2.16	0.46
2:B:102:DT:H2''	2:B:103:DA:O5'	2.16	0.45
1:A:112:DA:C8	1:A:113:DT:H72	2.52	0.45
2:B:109:DC:H5''	2:B:109:DC:C6	2.50	0.45
2:B:105:DT:H2''	2:B:106:DC:C6	2.52	0.45
4:D:115:DG:H2'	4:D:116:DA:C8	2.53	0.44
1:A:115:DC:H2''	1:A:116:DC:O5'	2.17	0.44
4:D:110:DT:H1'	4:D:111:DC:H5'	1.99	0.44
1:A:111:DA:H2''	1:A:112:DA:OP1	2.16	0.44
2:B:110:DA:C2	2:B:111:DC:C4	3.06	0.44
1:A:106:DG:H2''	1:A:107:DC:OP2	2.18	0.44
2:B:112:DG:H2''	2:B:113:DA:H8	1.82	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:109:DG:H2'	3:C:110:DG:H8	1.79	0.43
1:A:108:DC:C2	3:C:110:DG:N2	2.87	0.43
2:B:108:DC:H1'	3:C:108:DT:C2	2.53	0.42
1:A:117:DG:C4	1:A:118:DC:C5	3.07	0.42
4:D:113:DT:H2''	4:D:114:DG:H5''	2.01	0.42
1:A:115:DC:H4'	1:A:116:DC:OP1	2.19	0.41
4:D:116:DA:H2''	4:D:117:DC:C6	2.56	0.41
2:B:108:DC:C5	3:C:108:DT:C5	3.09	0.40
1:A:118:DC:H2''	1:A:119:DA:C8	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

5.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.