

# wwPDB X-ray Structure Validation Summary Report (i)

#### Oct 6, 2022 – 01:36 PM EDT

PDB ID : 5SKF

Title: Crystal Structure of human phosphodiesterase 10 in complex with 3-[5-[1-(2,2)]

- difluoro -1, 3-benzo dioxol -4-yl) -4-oxopyridazin -3-yl] pyrazol -1-yl] benzo nitrile

Authors: Joseph, C.; Benz, J.; Flohr, A.; Rodriguez-Sarmiento, R.M.; Rudolph, M.G.

Deposited on : 2022-02-01

Resolution : 2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.31.2buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$ 

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.31.2

### 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $X\text{-}RAY\ DIFFRACTION$ 

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \text{Whole archive} \\ (\#\text{Entries}) \end{array}$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries,\ resolution\ range(\mathring{\rm A})}) \end{array}$
$R_{free}$	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	343	80%	%	9%
1	В	343	79% 120	6	• 8%
1	С	343	78% 12%	•	9%
1	D	343	81%	6 •	9%



## 2 Entry composition (i)

There are 8 unique types of molecules in this entry. The entry contains 10893 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called cAMP and cAMP-inhibited cGMP 3',5'-cyclic phosphodiesterase 10A.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	Λ	313	Total	С	N	О	S	0	1	0
1	A	313	2549	1629	435	461	24	0	1	
1	В	315	Total	С	N	О	S	0	1	0
1	Б	319	2559	1635	437	463	24	0	1	U
1	C	212	Total	С	N	О	S	0	1	0
1		313	2549	1629	435	461	24	U	1	
1	D	313	Total	С	N	О	S	0	0	0
1	ש	313	2537	1622	433	458	24	0	0	

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	447	GLY	-	expression tag	UNP Q9Y233
A	448	SER	-	expression tag	UNP Q9Y233
В	447	GLY	-	expression tag	UNP Q9Y233
В	448	SER	-	expression tag	UNP Q9Y233
С	447	GLY	-	expression tag	UNP Q9Y233
С	448	SER	-	expression tag	UNP Q9Y233
D	447	GLY	-	expression tag	UNP Q9Y233
D	448	SER	-	expression tag	UNP Q9Y233

• Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Zn 1 1	0	0
2	В	1	Total Zn 1 1	0	0
2	С	1	Total Zn 1 1	0	0
2	D	1	Total Zn 1 1	0	0




• Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula:
---

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Mg 1 1	0	0
3	В	1	Total Mg 1 1	0	0
3	С	1	Total Mg 1 1	0	0
3	D	1	Total Mg 1 1	0	0

• Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Cl 1 1	0	0


• Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 6 3 3	0	0

• Molecule 6 is 3-{5-[1-(2,2-difluoro-2H-1,3-benzodioxol-4-yl)-4-oxo-1,4-dihydropyridazin-3-y l]-1H-pyrazol-1-yl}benzonitrile (three-letter code: KF9) (formula:  $C_{21}H_{11}F_2N_5O_3$ ) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
6	Λ	1	Total	С	F	N	О	0	0
	A	1	31	21	2	5	3	0	U
6	B	1	Total	С	F	N	О	0	0
	Б	1	31	21		5	3	0	U
6	С	1	Total	С	F	N	О	0	0
0		1	31	21	2	5	3	0	0
6	D	1	Total	С	F	N	О	0	0
	D	1	31	21	2	5	3	0	U

• Molecule 7 is PRASEODYMIUM ION (three-letter code: PR) (formula: Pr).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	В	5	Total Pr 5 5	0	0
7	С	3	Total Pr 3 3	0	0

• Molecule 8 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	141	Total O 141 141	0	0
8	В	151	Total O 151 151	0	0
8	С	156	Total O 156 156	0	0

Continued on next page...



Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	D	104	Total 104	O 104	0	0




### 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: cAMP and cAMP-inhibited cGMP 3',5'-cyclic phosphodiesterase 10A









# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	H 3	Depositor
Cell constants	135.05Å 135.05Å 234.40Å	Donositon
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.00^{\circ}$ $120.00^{\circ}$	Depositor
Resolution (Å)	39.07 - 2.10	Depositor
rtesolution (A)	39.07 - 2.10	EDS
% Data completeness	100.0 (39.07-2.10)	Depositor
(in resolution range)	99.1 (39.07-2.10)	EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.29 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
D D.	0.171 , 0.187	Depositor
$R, R_{free}$	0.177 , $0.189$	DCC
$R_{free}$ test set	4655 reflections $(5.00%)$	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	31.5	Xtriage
Anisotropy	0.099	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.35, 30.1	EDS
L-test for twinning <sup>2</sup>	$< L >=0.40, < L^2>=0.22$	Xtriage
Estimated twinning fraction	0.279 for h,-h-k,-l	Xtriage
Reported twinning fraction	0.701 for H, K, L	Depositor
Reported twinning fraction	0.299 for K, H, -L	Depositor
Outliers	0 of 93041 reflections	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	10893	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 2.65% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $<L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

### 5 Model quality (i)

### 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, PR, CME, ZN, KF9, MG, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
IVIOI	Chain	RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	A	0.63	0/2603	0.69	0/3521	
1	В	0.63	0/2613	0.70	0/3535	
1	С	0.62	0/2603	0.68	0/3521	
1	D	0.64	0/2588	0.68	0/3502	
All	All	0.63	0/10407	0.69	0/14079	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2549	0	2526	21	0
1	В	2559	0	2528	24	0
1	С	2549	0	2524	21	0
1	D	2537	0	2511	17	0
2	A	1	0	0	0	0
2	В	1	0	0	0	0
2	С	1	0	0	0	0
2	D	1	0	0	0	0
3	A	1	0	0	0	0

Continued on next page...



Continued from previous page...

Mol	Chain		H(model)	H(added)	Clashes	Symm-Clashes
3	В	1	0	0	0	0
3	С	1	0	0	0	0
3	D	1	0	0	0	0
4	A	1	0	0	0	0
5	A	6	0	8	0	0
6	A	31	0	0	0	0
6	В	31	0	0	0	0
6	С	31	0	0	0	0
6	D	31	0	0	0	0
7	В	5	0	0	0	0
7	С	3	0	0	0	0
8	A	141	0	0	0	0
8	В	151	0	0	1	0
8	С	156	0	0	2	0
8	D	104	0	0	1	0
All	All	10893	0	10097	80	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 80 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
1:C:616:SER:O	1:C:620:GLU:HG3	1.84	0.78
1:B:644:GLN:HE21	1:B:644:GLN:HA	1.55	0.71
1:A:461:GLN:NE2	1:A:461:GLN:HA	2.10	0.65
1:D:469:CYS:SG	8:D:999:HOH:O	2.55	0.63
1:B:727:LEU:HD12	1:B:766:ILE:HD12	1.81	0.61

There are no symmetry-related clashes.

#### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured Allowed		Outliers	Percentiles		
1	A	311/343~(91%)	304 (98%)	7 (2%)	0	100 100		
1	В	313/343~(91%)	304 (97%)	9 (3%)	0	100 100		
1	C	311/343~(91%)	303 (97%)	8 (3%)	0	100 100		
1	D	310/343~(90%)	302 (97%)	8 (3%)	0	100 100		
All	All	1245/1372~(91%)	1213 (97%)	32 (3%)	0	100 100		

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric Outliers		Percentiles		
1	A	282/305~(92%)	270 (96%)	12 (4%)	29 29		
1	В	282/305~(92%)	273 (97%)	9 (3%)	39 41		
1	$\mathbf{C}$	282/305~(92%)	271 (96%)	11 (4%)	32 33		
1	D	280/305~(92%)	267 (95%)	13 (5%)	27 26		
All	All	$1126/1220 \ (92\%)$	1081 (96%)	45 (4%)	31 32		

5 of 45 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	С	597	SER
1	D	504	GLU
1	С	605	LEU
1	D	457	GLN
1	D	517	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 26 such sidechains are listed below:

Mol	Chain	Res	Type
1	С	542	GLN
1	D	476	HIS
1	D	731	ASN

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type
1	С	743	GLN
1	D	484	ASN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Type		Chain Res	Res	Dan	Dan	Link	В	Bond lengths			Bond angles		
Mol   Type	LillK			Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2				
1	CME	D	509	1	8,9,10	0.49	0	5,9,11	0.30	0			
1	CME	В	509	1	8,9,10	0.34	0	5,9,11	0.23	0			
1	CME	A	509	1	8,9,10	0.47	0	5,9,11	0.40	0			
1	CME	С	509	1	8,9,10	0.50	0	5,9,11	0.39	0			

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CME	D	509	1	-	2/5/8/10	-
1	CME	В	509	1	-	2/5/8/10	-
1	CME	A	509	1	-	1/5/8/10	-
1	CME	С	509	1	-	2/5/8/10	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.



5 of 7 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	С	509	CME	CE-SD-SG-CB
1	В	509	CME	SD-CE-CZ-OH
1	С	509	CME	SD-CE-CZ-OH
1	A	509	CME	CZ-CE-SD-SG
1	В	509	CME	N-CA-CB-SG

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	С	509	CME	1	0

### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

#### 5.6 Ligand geometry (i)

Of 22 ligands modelled in this entry, 17 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Trmo	Clasia	Res	Link	Bond lengths				Bond angles		
Mol Type Chain	Chain	rtes	LILIK	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2		
5	GOL	A	804	-	5,5,5	0.11	0	5,5,5	0.32	0	
6	KF9	В	808	_	30,35,35	1.28	1 (3%)	34,52,52	2.13	9 (26%)	
6	KF9	A	805	-	30,35,35	1.31	1 (3%)	34,52,52	2.22	11 (32%)	
6	KF9	С	806	-	30,35,35	1.48	3 (10%)	34,52,52	2.12	8 (23%)	
6	KF9	D	803	-	30,35,35	1.35	2 (6%)	34,52,52	2.14	11 (32%)	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	A	804	-	-	1/4/4/4	-
6	KF9	В	808	-	-	0/9/24/24	0/5/5/5
6	KF9	A	805	-	-	0/9/24/24	0/5/5/5
6	KF9	С	806	-	-	0/9/24/24	0/5/5/5
6	KF9	D	803	-	-	0/9/24/24	0/5/5/5

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$Observed(\mathring{A})$	Ideal(Å)
6	D	803	KF9	N4-N2	-5.78	1.27	1.38
6	В	808	KF9	N4-N2	-5.77	1.27	1.38
6	С	806	KF9	N4-N2	-5.68	1.28	1.38
6	A	805	KF9	N4-N2	-5.44	1.28	1.38
6	С	806	KF9	C8-N4	-3.14	1.39	1.43

The worst 5 of 39 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
6	С	806	KF9	C20-N11-N5	8.34	109.77	103.70
6	В	808	KF9	C20-N11-N5	7.36	109.05	103.70
6	A	805	KF9	C20-N11-N5	7.17	108.91	103.70
6	D	803	KF9	C20-N11-N5	6.94	108.75	103.70
6	A	805	KF9	C14-C15-C12	-6.09	117.06	121.31

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	804	GOL	O1-C1-C2-C3


There are no ring outliers.

No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.




The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.
















## 5.7 Other polymers (i)

There are no such residues in this entry.

### 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



### 6 Fit of model and data (i)

#### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$\#\mathrm{RSRZ}{>}2$	$OWAB(A^2)$	Q<0.9
1	A	312/343~(90%)	-0.10	6 (1%) 66 71	24, 36, 58, 72	0
1	В	314/343~(91%)	-0.09	8 (2%) 57 62	24, 35, 56, 79	0
1	С	312/343~(90%)	-0.17	3 (0%) 82 85	23, 36, 53, 74	0
1	D	312/343~(90%)	-0.13	8 (2%) 56 61	27, 40, 58, 76	0
All	All	1250/1372~(91%)	-0.12	25 (2%) 65 69	23, 37, 57, 79	0

The worst 5 of 25 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	458	GLY	5.1
1	D	460	MET	4.5
1	С	770	GLU	4.3
1	D	456	TRP	4.0
1	A	460	MET	3.9

### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
1	CME	D	509	10/11	0.82	0.24	47,54,72,73	0
1	CME	С	509	10/11	0.85	0.14	39,45,62,67	0
1	CME	В	509	10/11	0.88	0.19	38,47,63,64	0
1	CME	A	509	10/11	0.90	0.17	39,46,61,62	0



#### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.


### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}({ ext{\AA}}^2)$	Q<0.9
7	PR	В	804	1/1	0.85	0.18	138,138,138,138	0
5	GOL	A	804	6/6	0.89	0.18	51,57,61,65	0
7	PR	С	805	1/1	0.91	0.04	92,92,92,92	1
7	PR	В	806	1/1	0.93	0.03	110,110,110,110	0
6	KF9	A	805	31/31	0.93	0.11	27,34,39,40	0
6	KF9	В	808	31/31	0.94	0.11	30,35,43,44	0
4	CL	A	803	1/1	0.95	0.06	68,68,68,68	0
7	PR	В	805	1/1	0.95	0.04	108,108,108,108	0
3	MG	D	802	1/1	0.95	0.09	29,29,29,29	0
7	PR	С	804	1/1	0.95	0.03	99,99,99,99	1
6	KF9	D	803	31/31	0.95	0.10	32,35,44,49	0
6	KF9	С	806	31/31	0.96	0.10	27,32,42,45	0
7	PR	В	807	1/1	0.97	0.07	107,107,107,107	0
3	MG	С	802	1/1	0.98	0.10	24,24,24,24	0
7	PR	С	803	1/1	0.98	0.04	93,93,93,93	1
3	MG	В	802	1/1	0.99	0.12	21,21,21,21	0
7	PR	В	803	1/1	0.99	0.06	78,78,78,78	0
3	MG	A	802	1/1	0.99	0.11	22,22,22,22	0
2	ZN	С	801	1/1	1.00	0.13	29,29,29,29	0
2	ZN	D	801	1/1	1.00	0.11	33,33,33,33	0
2	ZN	A	801	1/1	1.00	0.14	28,28,28,28	0
2	ZN	В	801	1/1	1.00	0.14	27,27,27,27	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



















## 6.5 Other polymers (i)

There are no such residues in this entry.

