



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 5, 2024 – 12:13 PM EST

PDB ID : 1SL2
Title : Ternary 5' complex of T7 DNA polymerase with a DNA primer/template containing a cis-syn thymine dimer on the template and an incoming nucleotide
Authors : Li, Y.; Dutta, S.; Doublie, S.; Bdour, H.M.; Taylor, J.S.; Ellenberger, T.
Deposited on : 2004-03-05
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	P	22	
2	T	25	
3	A	698	
4	B	108	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	DAD	A	4004	X	-	-	-

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 6558 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called 5'-D(*CP*GP*AP*AP*AP*AP*CP*GP*AP*CP*GP*GP*CP*CP*AP*GP*TP*GP*CP*CP*TP*(2DA))-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
1	P	9	182	87	33	53	9	0	0	0

- Molecule 2 is a DNA chain called 5'-D(*CP*CP*CP*(TTD)P*AP*GP*GP*CP*AP*CP*TP*GP*GP*CP*CP*GP*TP*CP*GP*TP*TP*TP*TP*CP*G)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	T	10	228	108	42	67	11	0	0	0

- Molecule 3 is a protein called DNA polymerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	A	667	5113	3248	892	950	23	0	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	?	-	LYS	deletion	UNP P00581
A	?	-	ARG	deletion	UNP P00581
A	?	-	PHE	deletion	UNP P00581
A	?	-	GLY	deletion	UNP P00581
A	?	-	SER	deletion	UNP P00581
A	?	-	HIS	deletion	UNP P00581


- Molecule 4 is a protein called Thioredoxin 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	B	105	780	502	125	150	3	0	0	0

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	3	Total	Mg	0	0
			3	3		

- Molecule 6 is 2',3'-DIDEOXYADENOSINE-5'-TRIPHOSPHATE (three-letter code: DAD) (formula: C₁₀H₁₆N₅O₁₁P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total	C	N	O	P	0	0
			29	10	5	11	3		



- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	P	1	Total	O	0	0
			1	1		
7	T	12	Total	O	0	0
			12	12		
7	A	194	Total	O	0	0
			194	194		
7	B	16	Total	O	0	0
			16	16		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: 5'-D(*CP*GP*AP*AP*AP*AP*CP*GP*AP*CP*GP*GP*CP*CP*AP*GP*TP*GP*CP*CP*TP*(2DA))-3'




- Molecule 2: 5'-D(*CP*CP*CP*(TTD)P*AP*GP*GP*CP*AP*CP*TP*GP*GP*CP*CP*GP*TP*CP*GP*TP*TP*TP*TP*CP*G)-3'




- Molecule 3: DNA polymerase





● Molecule 4: Thioredoxin 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	104.98Å 212.60Å 51.95Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.30 33.57 – 2.30	Depositor EDS
% Data completeness (in resolution range)	92.8 (50.00-2.30) 92.9 (33.57-2.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.84 (at 2.31Å)	Xtrriage
Refinement program	CNS	Depositor
R, R_{free}	0.246 , 0.279 0.234 , 0.268	Depositor DCC
R_{free} test set	2416 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å ²)	33.9	Xtrriage
Anisotropy	0.551	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 41.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	6558	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.75% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 2DA, MG, TTD, DAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	P	0.52	0/180	0.88	0/275
2	T	0.36	0/211	0.81	0/324
3	A	0.35	0/5240	0.57	0/7127
4	B	0.31	0/795	0.57	0/1084
All	All	0.35	0/6426	0.60	0/8810

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	P	0	1
2	T	0	1
All	All	0	2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	P	20	DC	Sidechain
2	T	7	DG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	P	182	0	102	7	0
2	T	228	0	125	3	0
3	A	5113	0	4719	90	0
4	B	780	0	747	13	0
5	A	3	0	0	0	0
6	A	29	0	12	3	0
7	A	194	0	0	4	0
7	B	16	0	0	2	0
7	P	1	0	0	0	0
7	T	12	0	0	0	0
All	All	6558	0	5705	109	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (109) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:570:GLN:HE22	3:A:606:PRO:HB3	1.34	0.92
3:A:94:LEU:HB3	3:A:185:LEU:HD13	1.57	0.85
3:A:173:GLN:O	3:A:176:VAL:HG22	1.82	0.80
3:A:343:GLN:HG3	3:A:362:PRO:HG3	1.73	0.70
3:A:391:MET:O	3:A:391:MET:HE2	1.92	0.69
3:A:35:ARG:HB3	3:A:36:PRO:HD2	1.73	0.69
3:A:343:GLN:HG3	3:A:362:PRO:CG	2.23	0.68
3:A:480:GLU:CD	6:A:4004:DAD:H2'1	2.13	0.68
4:B:32:CYS:SG	4:B:34:PRO:HD2	2.35	0.67
3:A:530:TYR:CE1	3:A:614:LEU:HD12	2.30	0.66
3:A:141:ASP:O	3:A:145:ARG:HG2	1.96	0.66
3:A:135:MET:HG3	3:A:174:ASP:OD1	1.97	0.65
1:P:16:DG:H2''	1:P:17:DT:C5'	2.28	0.64
4:B:47:ASP:O	4:B:50:GLN:HG3	1.97	0.63
1:P:16:DG:H2''	1:P:17:DT:H5'	1.81	0.63
3:A:530:TYR:HE1	3:A:614:LEU:HD12	1.63	0.63
3:A:66:VAL:HB	3:A:67:PRO:HD3	1.82	0.62

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:506:HIS:HA	3:A:509:ASN:HD22	1.64	0.62
3:A:704:HIS:HB3	7:A:5155:HOH:O	1.98	0.62
3:A:19:HIS:O	3:A:36:PRO:HD3	2.01	0.59
3:A:79:ARG:HD3	7:A:5071:HOH:O	2.02	0.59
3:A:421:ASN:HB3	3:A:431:THR:OG1	2.02	0.59
3:A:64:TYR:O	3:A:67:PRO:HD2	2.03	0.58
3:A:530:TYR:CE1	3:A:611:ASN:HA	2.39	0.58
3:A:478:GLY:HA2	6:A:4004:DAD:O2B	2.04	0.58
3:A:522:LYS:HE3	6:A:4004:DAD:O1B	2.03	0.58
3:A:326:TYR:HB3	4:B:92:GLY:HA2	1.87	0.57
4:B:39:ALA:HB3	4:B:40:PRO:HD3	1.87	0.56
2:T:9:DC:H3'	3:A:404:LYS:HG3	1.87	0.56
3:A:632:MET:O	3:A:635:GLU:HG2	2.06	0.56
3:A:91:THR:HB	3:A:181:LEU:HD13	1.87	0.55
3:A:85:ARG:HG3	3:A:222:TRP:CG	2.41	0.55
3:A:391:MET:HE3	3:A:395:ARG:HG3	1.89	0.55
3:A:233:PRO:HB2	3:A:456:GLY:O	2.07	0.54
3:A:49:GLU:HA	3:A:52:ARG:HH11	1.73	0.53
3:A:49:GLU:HA	3:A:52:ARG:NH1	2.24	0.52
3:A:391:MET:HE1	3:A:392:ILE:HA	1.90	0.52
3:A:678:ARG:NH1	3:A:691:ASP:OD1	2.36	0.52
1:P:20:DC:H6	1:P:20:DC:H5'	1.76	0.51
3:A:343:GLN:HG3	3:A:362:PRO:HG2	1.93	0.51
3:A:336:PRO:HB2	3:A:389:TYR:CD1	2.45	0.51
3:A:574:VAL:HG12	3:A:575:GLU:N	2.26	0.51
4:B:13:ASP:OD1	4:B:18:LYS:HE3	2.11	0.51
3:A:362:PRO:O	3:A:364:VAL:HG23	2.11	0.51
2:T:8:DG:H4'	3:A:432:HIS:O	2.11	0.50
3:A:237:LYS:O	3:A:241:GLU:HG3	2.10	0.50
3:A:632:MET:HA	3:A:635:GLU:HG2	1.94	0.49
4:B:13:ASP:HB3	7:B:5229:HOH:O	2.11	0.49
1:P:18:DG:H4'	3:A:339:ARG:NH2	2.28	0.48
3:A:638:LEU:HD12	3:A:638:LEU:N	2.27	0.48
1:P:19:DC:C2	1:P:20:DC:C5	3.02	0.48
3:A:176:VAL:HG23	3:A:177:VAL:N	2.29	0.48
3:A:391:MET:HE2	3:A:391:MET:C	2.34	0.48
3:A:131:ARG:O	3:A:135:MET:HG2	2.14	0.47
3:A:368:VAL:O	3:A:372:VAL:HG23	2.15	0.47
4:B:37:MET:O	4:B:40:PRO:HD2	2.15	0.47
3:A:173:GLN:O	3:A:176:VAL:CG2	2.59	0.47
3:A:484:LEU:O	3:A:488:MET:HG2	2.14	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:667:GLN:CG	3:A:696:MET:HE1	2.44	0.47
3:A:357:THR:OG1	3:A:361:ALA:HB3	2.14	0.47
3:A:446:PRO:O	3:A:447:TYR:HB2	2.15	0.46
3:A:142:ASP:O	3:A:146:MET:HG3	2.16	0.46
3:A:448:GLY:O	3:A:452:ARG:HB2	2.15	0.46
3:A:512:ALA:C	3:A:514:GLU:H	2.18	0.45
3:A:518:ARG:O	3:A:522:LYS:HG3	2.16	0.45
3:A:173:GLN:HA	3:A:176:VAL:HG22	1.98	0.45
3:A:323:GLY:O	3:A:325:PRO:HD3	2.17	0.45
3:A:429:ARG:HG3	3:A:619:ALA:HB2	1.99	0.45
3:A:678:ARG:NH1	3:A:690:LEU:O	2.50	0.45
3:A:130:TYR:CZ	3:A:134:GLU:HG3	2.52	0.45
3:A:57:VAL:HG22	3:A:89:ILE:HB	1.98	0.45
3:A:292:PRO:O	3:A:322:ALA:HA	2.17	0.44
3:A:512:ALA:O	3:A:514:GLU:N	2.51	0.44
3:A:422:PRO:HB2	7:A:5024:HOH:O	2.17	0.44
4:B:80:LEU:HD23	4:B:81:PHE:N	2.32	0.44
3:A:365:ASP:OD1	3:A:368:VAL:HG13	2.17	0.44
3:A:522:LYS:O	3:A:525:ILE:HG22	2.17	0.44
3:A:512:ALA:C	3:A:514:GLU:N	2.71	0.44
3:A:315:LEU:C	3:A:315:LEU:HD23	2.37	0.44
4:B:67:ALA:HB3	4:B:68:PRO:HD3	1.98	0.43
4:B:7:LEU:HD13	4:B:12:PHE:CD2	2.53	0.43
3:A:638:LEU:N	3:A:638:LEU:CD1	2.80	0.43
3:A:597:ASP:OD1	3:A:599:ARG:HD2	2.19	0.43
4:B:53:LEU:HG	4:B:54:THR:N	2.32	0.43
3:A:17:LYS:HB3	3:A:17:LYS:NZ	2.34	0.43
3:A:336:PRO:HB2	3:A:389:TYR:CE1	2.54	0.43
3:A:420:VAL:O	3:A:422:PRO:HD3	2.18	0.43
3:A:597:ASP:OD1	3:A:599:ARG:CD	2.67	0.43
3:A:500:ILE:C	3:A:502:ASN:H	2.23	0.42
4:B:14:THR:HG23	7:B:5229:HOH:O	2.18	0.42
3:A:140:LYS:HE3	3:A:154:TYR:OH	2.20	0.42
3:A:279:THR:CG2	3:A:281:LYS:HB2	2.49	0.42
3:A:589:LYS:HE3	3:A:589:LYS:HB2	1.90	0.42
3:A:442:GLY:O	3:A:448:GLY:HA3	2.20	0.42
3:A:79:ARG:HG2	3:A:80:GLU:N	2.35	0.42
3:A:53:GLY:HA2	7:A:5027:HOH:O	2.20	0.41
1:P:16:DG:H1'	1:P:17:DT:H5''	2.03	0.41
3:A:351:TRP:CH2	3:A:353:PRO:HG3	2.56	0.41
1:P:21:DT:H2'	1:P:22:2DA:C8	2.51	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:T:5:TTD:H72	2:T:5:TTD:H5A1	1.91	0.41
3:A:15:VAL:HG22	3:A:72:LEU:HD21	2.02	0.41
3:A:138:GLU:O	3:A:142:ASP:OD1	2.39	0.41
3:A:364:VAL:O	3:A:364:VAL:HG12	2.21	0.41
3:A:391:MET:CE	3:A:395:ARG:HG3	2.52	0.41
3:A:158:MET:HA	3:A:161:TRP:CE2	2.56	0.40
3:A:172:VAL:O	3:A:176:VAL:HG13	2.21	0.40
3:A:452:ARG:HG3	3:A:700:TRP:HB3	2.02	0.40
4:B:79:LEU:HD12	4:B:79:LEU:HA	1.92	0.40
3:A:276:HIS:HA	3:A:277:PRO:HD3	1.92	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	A	659/698 (94%)	637 (97%)	19 (3%)	3 (0%)	29	35
4	B	103/108 (95%)	100 (97%)	3 (3%)	0	100	100
All	All	762/806 (94%)	737 (97%)	22 (3%)	3 (0%)	34	42

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	A	359	LYS
3	A	653	HIS
3	A	513	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	A	475/579 (82%)	462 (97%)	13 (3%)	44	61
4	B	73/87 (84%)	71 (97%)	2 (3%)	44	61
All	All	548/666 (82%)	533 (97%)	15 (3%)	44	61

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	A	85	ARG
3	A	94	LEU
3	A	142	ASP
3	A	171	ASN
3	A	181	LEU
3	A	232	PHE
3	A	343	GLN
3	A	391	MET
3	A	599	ARG
3	A	624	LEU
3	A	686	PHE
3	A	693	GLU
3	A	700	TRP
4	B	89	THR
4	B	99	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
3	A	276	HIS
3	A	343	GLN
3	A	347	GLN
3	A	509	ASN
3	A	510	GLN
3	A	570	GLN
4	B	50	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	2DA	P	22	1	17,22,23	0.72	0	13,31,34	0.82	0
2	TTD	T	5	2	42,45,46	4.00	10 (23%)	62,74,77	3.12	20 (32%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	2DA	P	22	1	-	0/3/18/19	0/3/3/3
2	TTD	T	5	2	-	4/22/109/110	0/5/6/6

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	T	5	TTD	C5T-C6T	-20.31	1.31	1.55
2	T	5	TTD	C5-C6	-11.70	1.41	1.55
2	T	5	TTD	C1R-N1T	5.84	1.53	1.45
2	T	5	TTD	C5A-C5	4.30	1.62	1.53
2	T	5	TTD	C1'-N1	3.84	1.50	1.45
2	T	5	TTD	C5M-C5T	3.05	1.59	1.53
2	T	5	TTD	C2-N1	2.97	1.42	1.36
2	T	5	TTD	C2-N3	-2.63	1.33	1.38
2	T	5	TTD	C6T-C6	2.26	1.63	1.56
2	T	5	TTD	O4'-C1'	2.26	1.47	1.42

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	T	5	TTD	O4R-C1R-N1T	9.66	120.09	108.65
2	T	5	TTD	O4T-C4T-C5T	8.37	129.56	122.88
2	T	5	TTD	C5-C5T-C6T	7.88	98.18	88.38
2	T	5	TTD	C5T-C5-C4	7.78	138.85	113.21
2	T	5	TTD	O4'-C1'-N1	-6.27	101.23	108.65
2	T	5	TTD	C5T-C5-C6	-5.84	81.10	88.38
2	T	5	TTD	C5A-C5-C4	-5.83	98.17	108.22
2	T	5	TTD	C2R-C1R-N1T	-4.69	109.26	115.59
2	T	5	TTD	C5T-C6T-N1T	4.68	122.17	115.61
2	T	5	TTD	C5-C5T-C4T	-4.42	98.62	113.21
2	T	5	TTD	C4'-O4R-C1R	-4.40	98.82	109.45
2	T	5	TTD	C5T-C6T-C6	-4.28	82.26	89.28
2	T	5	TTD	O4T-C4T-N3T	-4.17	113.85	120.50
2	T	5	TTD	C2'-C1'-N1	3.16	119.85	115.59
2	T	5	TTD	O4-C4-C5	2.80	125.11	122.88
2	T	5	TTD	C5-C6-N1	2.64	119.31	115.61
2	T	5	TTD	C5'-C4R-C3R	-2.47	108.89	114.53
2	T	5	TTD	C3'-C2R-C1R	-2.18	97.07	102.54
2	T	5	TTD	O4'-C1'-C2'	-2.15	102.19	106.25
2	T	5	TTD	C5A-C5-C5T	-2.08	110.36	116.39

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	T	5	TTD	C2'-C1'-N1-C6
2	T	5	TTD	C2'-C1'-N1-C2
2	T	5	TTD	O4'-C1'-N1-C6
2	T	5	TTD	O4'-C1'-N1-C2

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	P	22	2DA	1	0
2	T	5	TTD	1	0

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry

Of 4 ligands modelled in this entry, 3 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	DAD	A	4004	5	25,31,31	0.77	0	26,48,48	1.07	2 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	DAD	A	4004	5	2/2/5/5	3/18/31/31	0/3/3/3

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	4004	DAD	O2G-PG-O1G	2.34	119.86	110.68
6	A	4004	DAD	C5-C6-N6	2.26	123.78	120.35

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
6	A	4004	DAD	C1'
6	A	4004	DAD	C4'

All (3) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
6	A	4004	DAD	PG-O3B-PB-O1B
6	A	4004	DAD	PG-O3B-PB-O2B
6	A	4004	DAD	C4'-C5'-O5'-PA

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	4004	DAD	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	P	8/22 (36%)	0.44	1 (12%) 3 5	34, 45, 60, 67	0
2	T	9/25 (36%)	-0.24	1 (11%) 5 7	31, 37, 47, 61	0
3	A	667/698 (95%)	0.35	54 (8%) 12 16	18, 31, 55, 61	0
4	B	105/108 (97%)	0.06	1 (0%) 82 86	28, 38, 50, 55	0
All	All	789/853 (92%)	0.31	57 (7%) 15 20	18, 32, 55, 67	0

All (57) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	A	551	LEU	9.6
3	A	503	GLY	7.8
3	A	360	GLY	7.4
3	A	541	VAL	7.0
3	A	548	GLY	5.7
3	A	540	ILE	5.6
3	A	516	PRO	5.5
3	A	542	GLY	5.5
3	A	114	LYS	5.3
3	A	543	ALA	5.3
3	A	523	THR	5.1
3	A	555	PHE	5.1
3	A	294	VAL	5.1
3	A	539	GLN	5.1
3	A	358	ASP	4.6
3	A	535	GLU	4.2
3	A	552	LYS	4.2
3	A	145	ARG	4.2
3	A	501	LEU	4.1
3	A	511	ILE	4.0
3	A	558	ASN	4.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	A	509	ASN	3.9
3	A	550	GLU	3.8
3	A	547	ARG	3.6
3	A	515	LEU	3.5
3	A	512	ALA	3.3
3	A	505	ILE	3.2
3	A	322	ALA	3.1
3	A	361	ALA	3.0
3	A	656	ILE	3.0
3	A	520	ASN	2.9
3	A	113	GLY	2.9
3	A	517	THR	2.8
3	A	544	GLY	2.8
3	A	502	ASN	2.8
3	A	354	THR	2.7
3	A	57	VAL	2.6
3	A	194	PRO	2.6
3	A	91	THR	2.6
3	A	549	LYS	2.6
3	A	534	ASP	2.5
3	A	295	GLY	2.4
3	A	524	PHE	2.4
3	A	521	ALA	2.3
3	A	704	HIS	2.3
3	A	148	GLU	2.3
3	A	4	SER	2.2
3	A	317	THR	2.2
3	A	526	TYR	2.2
2	T	14	DG	2.2
4	B	50	GLN	2.2
3	A	269	GLY	2.1
3	A	89	ILE	2.1
3	A	553	LYS	2.1
3	A	538	GLY	2.1
1	P	14	DC	2.0
3	A	528	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column

labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	TTD	T	5	40/41	0.70	0.28	47,60,67,67	0
1	2DA	P	22	20/21	0.95	0.15	34,42,46,46	0

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	MG	A	4003	1/1	-0.12	0.40	47,47,47,47	0
6	DAD	A	4004	29/29	0.81	0.22	50,52,56,58	0
5	MG	A	4002	1/1	0.89	0.08	36,36,36,36	0
5	MG	A	4001	1/1	0.90	0.15	39,39,39,39	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.