



# wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 22, 2020 – 02:37 PM BST

PDB ID : 6SLI  
Title : Structure of the RagAB peptide transporter  
Authors : Madej, M.; Ranson, N.A.; White, J.B.R.  
Deposited on : 2019-08-19  
Resolution : 3.38 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.13.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.38 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1691 (3.46-3.30)
Clashscore	141614	1762 (3.46-3.30)
Ramachandran outliers	138981	1732 (3.46-3.30)
Sidechain outliers	138945	1731 (3.46-3.30)
RSRZ outliers	127900	1635 (3.46-3.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	488	<p>74% 24% ..</p>
1	C	488	<p>77% 21% ..</p>
1	F	488	<p>78% 20% .</p>
1	I	488	<p>78% 21% .</p>
2	B	997	<p>54% 33% . 10%</p>
2	D	997	<p>53% 34% . 10%</p>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	G	997	<p>9% 56% 32% 10%</p>
2	J	997	<p>6% 56% 31% 10%</p>
3	H	13	<p>8% 69% 23% 8%</p>
3	P	13	<p>8% 62% 31% 8%</p>
4	E	13	<p>8% 38% 38% 23%</p>
4	K	13	<p>8% 69% 23% 8%</p>

## 2 Entry composition i

There are 6 unique types of molecules in this entry. The entry contains 43926 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Lipoprotein RagB.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	482	3842	2440	656	737	9	0	0	0
1	C	482	3842	2440	656	737	9	0	0	0
1	F	482	3842	2440	656	737	9	0	0	0
1	I	482	3842	2440	656	737	9	0	0	0

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	502	HIS	-	expression tag	UNP F5H948
A	503	HIS	-	expression tag	UNP F5H948
A	504	HIS	-	expression tag	UNP F5H948
A	505	HIS	-	expression tag	UNP F5H948
A	506	HIS	-	expression tag	UNP F5H948
A	507	HIS	-	expression tag	UNP F5H948
C	502	HIS	-	expression tag	UNP F5H948
C	503	HIS	-	expression tag	UNP F5H948
C	504	HIS	-	expression tag	UNP F5H948
C	505	HIS	-	expression tag	UNP F5H948
C	506	HIS	-	expression tag	UNP F5H948
C	507	HIS	-	expression tag	UNP F5H948
F	502	HIS	-	expression tag	UNP F5H948
F	503	HIS	-	expression tag	UNP F5H948
F	504	HIS	-	expression tag	UNP F5H948
F	505	HIS	-	expression tag	UNP F5H948
F	506	HIS	-	expression tag	UNP F5H948
F	507	HIS	-	expression tag	UNP F5H948
I	502	HIS	-	expression tag	UNP F5H948
I	503	HIS	-	expression tag	UNP F5H948
I	504	HIS	-	expression tag	UNP F5H948

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
I	505	HIS	-	expression tag	UNP F5H948
I	506	HIS	-	expression tag	UNP F5H948
I	507	HIS	-	expression tag	UNP F5H948

- Molecule 2 is a protein called RagA protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	900	Total	C	N	O	S	0	0	0
			7048	4466	1179	1371	32			
2	D	900	Total	C	N	O	S	0	0	0
			7048	4466	1179	1371	32			
2	G	900	Total	C	N	O	S	0	0	0
			7048	4466	1179	1371	32			
2	J	900	Total	C	N	O	S	0	0	0
			7048	4466	1179	1371	32			


- Molecule 3 is a protein called ALA-SER-THR-THR-GLY-GLY-ASN-SER-GLN-ARG-GLY-SER-GLY.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	P	13	Total	C	N	O	0	0	0
			81	43	18	20			
3	H	13	Total	C	N	O	0	0	0
			81	43	18	20			

- Molecule 4 is a protein called ASTTGGNSQRGGG.


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	E	13	Total	C	N	O	0	0	0
			79	42	18	19			
4	K	13	Total	C	N	O	0	0	0
			79	42	18	19			

- Molecule 5 is (1R,4S,6R)-6-([2-(ACETYLAMINO)-2-DEOXY-ALPHA-D-GLUCOPYRANOSYL]OXY)METHYL)-4-HYDROXY-1-[[[(15-METHYLHEXADECANOYL)OXY]METHYL]-4-OXIDO-7-OXO-3,5-DIOXA-8-AZA-4-PHOSPHAHEPTACOS-1-YL 15-METHYLHEXADECANOATE (three-letter code: 5PL) (formula: C<sub>67</sub>H<sub>129</sub>N<sub>2</sub>O<sub>15</sub>P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
5	B	1	20	16	4	0	0

- Molecule 6 is (HYDROXYETHYLOXY)TRI(ETHYLOXY)OCTANE (three-letter code: C8E) (formula:  $C_{16}H_{34}O_5$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
6	B	1	13	11	2	0	0
6	B	1	13	11	2	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: Lipoprotein RagB




#### • Molecule 1: Lipoprotein RagB




#### • Molecule 1: Lipoprotein RagB






• Molecule 1: Lipoprotein RagB




• Molecule 2: RagA protein













• Molecule 2: RagA protein





• Molecule 2: RagA protein






- Molecule 3: ALA-SER-THR-THR-GLY-GLY-ASN-SER-GLN-ARG-GLY-SER-GLY




- Molecule 3: ALA-SER-THR-THR-GLY-GLY-ASN-SER-GLN-ARG-GLY-SER-GLY



- Molecule 4: ASTTGGNSQRGGG



- Molecule 4: ASTTGGNSQRGGG



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	190.56Å 376.87Å 369.24Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	83.92 – 3.38 83.92 – 3.38	Depositor EDS
% Data completeness (in resolution range)	99.7 (83.92-3.38) 99.9 (83.92-3.38)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.35 (at 3.41Å)	Xtrriage
Refinement program	PHENIX 1.15.2_3472, PHENIX 1.15.2_3472	Depositor
R, $R_{free}$	0.204 , 0.251 0.204 , 0.251	Depositor DCC
$R_{free}$ test set	9223 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	83.0	Xtrriage
Anisotropy	0.759	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 77.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.43$ , $\langle L^2 \rangle = 0.26$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	43926	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	101.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.71% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: 5PL, C8E

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.64	0/3928	0.84	5/5331 (0.1%)
1	C	0.65	0/3928	0.85	4/5331 (0.1%)
1	F	0.59	0/3928	0.79	3/5331 (0.1%)
1	I	0.47	0/3928	0.66	1/5331 (0.0%)
2	B	0.64	4/7211 (0.1%)	0.88	9/9753 (0.1%)
2	D	0.61	4/7211 (0.1%)	0.84	5/9753 (0.1%)
2	G	0.51	1/7211 (0.0%)	0.76	5/9753 (0.1%)
2	J	0.50	0/7211	0.75	2/9753 (0.0%)
3	H	0.45	0/80	0.74	0/105
3	P	0.53	0/80	0.93	0/105
4	E	0.38	0/78	0.72	0/102
4	K	0.49	0/78	0.70	0/102
All	All	0.58	9/44872 (0.0%)	0.80	34/60750 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
2	B	0	1
2	D	0	1
2	J	0	1
All	All	0	4

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	498	ASN	C-N	8.83	1.54	1.34
2	B	498	ASN	C-N	6.64	1.49	1.34

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	G	498	ASN	C-N	5.95	1.47	1.34
2	B	552	VAL	CB-CG2	-5.62	1.41	1.52
2	D	440	THR	C-N	-5.53	1.21	1.34

The worst 5 of 34 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	498	ASN	C-N-CA	-10.40	95.70	121.70
2	B	498	ASN	C-N-CA	-10.38	95.76	121.70
2	G	498	ASN	C-N-CA	-9.89	96.97	121.70
2	J	498	ASN	C-N-CA	-9.16	98.81	121.70
1	C	188	LEU	CB-CG-CD2	-7.85	97.65	111.00

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	94	ASP	Peptide
2	B	292	ALA	Peptide
2	D	292	ALA	Peptide
2	J	809	TYR	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3842	0	3739	76	0
1	C	3842	0	3740	71	0
1	F	3842	0	3740	78	0
1	I	3842	0	3740	71	0
2	B	7048	0	6792	289	1
2	D	7048	0	6791	272	1
2	G	7048	0	6792	243	0
2	J	7048	0	6791	247	0
3	H	81	0	75	5	0
3	P	81	0	75	4	0
4	E	79	0	73	10	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	K	79	0	73	3	0
5	B	20	0	21	4	0
6	B	26	0	42	1	0
All	All	43926	0	42484	1303	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

The worst 5 of 1303 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:142:LEU:HD21	2:B:203:MET:HE3	1.35	1.08
1:I:465:ASN:OD1	2:J:276:GLY:N	1.88	1.05
2:B:142:LEU:CD1	2:B:203:MET:HE1	1.87	1.04
2:B:142:LEU:HD21	2:B:203:MET:CE	1.87	1.04
2:B:754:ASN:HB3	2:B:782:ASN:HD21	1.25	0.98

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:342:SER:OG	2:D:342:SER:OG[3_655]	1.97	0.23
2:B:342:SER:OG	2:B:342:SER:OG[3_755]	2.09	0.11

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	480/488 (98%)	439 (92%)	38 (8%)	3 (1%)	25 59
1	C	480/488 (98%)	444 (92%)	33 (7%)	3 (1%)	25 59

*Continued on next page...*



Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	F	480/488 (98%)	444 (92%)	33 (7%)	3 (1%)	25	59
1	I	480/488 (98%)	444 (92%)	33 (7%)	3 (1%)	25	59
2	B	896/997 (90%)	760 (85%)	118 (13%)	18 (2%)	7	33
2	D	896/997 (90%)	760 (85%)	117 (13%)	19 (2%)	7	32
2	G	896/997 (90%)	768 (86%)	111 (12%)	17 (2%)	8	34
2	J	896/997 (90%)	759 (85%)	118 (13%)	19 (2%)	7	32
3	H	11/13 (85%)	7 (64%)	4 (36%)	0	100	100
3	P	11/13 (85%)	7 (64%)	4 (36%)	0	100	100
4	E	11/13 (85%)	7 (64%)	3 (27%)	1 (9%)	1	5
4	K	11/13 (85%)	6 (54%)	4 (36%)	1 (9%)	1	5
All	All	5548/5992 (93%)	4845 (87%)	616 (11%)	87 (2%)	9	37

5 of 87 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	642	GLN
2	B	712	PHE
2	B	809	TYR
2	B	874	TRP
2	D	532	ASP

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	402/408 (98%)	399 (99%)	3 (1%)	84	92
1	C	402/408 (98%)	398 (99%)	4 (1%)	76	87
1	F	402/408 (98%)	399 (99%)	3 (1%)	84	92
1	I	402/408 (98%)	398 (99%)	4 (1%)	76	87
2	B	752/832 (90%)	710 (94%)	42 (6%)	21	52
2	D	752/832 (90%)	715 (95%)	37 (5%)	25	57

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	G	752/832 (90%)	714 (95%)	38 (5%)	24	55
2	J	752/832 (90%)	710 (94%)	42 (6%)	21	52
3	H	8/8 (100%)	7 (88%)	1 (12%)	4	18
3	P	8/8 (100%)	5 (62%)	3 (38%)	0	0
4	E	7/7 (100%)	5 (71%)	2 (29%)	0	1
4	K	7/7 (100%)	6 (86%)	1 (14%)	3	14
All	All	4646/4990 (93%)	4466 (96%)	180 (4%)	32	62

5 of 180 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	D	905	LEU
2	G	263	ASP
2	J	608	PHE
2	D	956	TYR
1	F	182	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 25 such sidechains are listed below:

Mol	Chain	Res	Type
2	D	754	ASN
2	D	932	GLN
2	J	754	ASN
2	D	782	ASN
2	D	1014	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
6	C8E	B	1102	-	12,12,20	0.43	0	11,11,19	0.56	0
5	5PL	B	1101	1	19,19,85	1.80	5 (26%)	21,21,101	1.81	4 (19%)
6	C8E	B	1103	-	12,12,20	0.42	0	11,11,19	0.70	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	C8E	B	1102	-	-	5/10/10/18	-
5	5PL	B	1101	1	-	9/20/20/107	-
6	C8E	B	1103	-	-	6/10/10/18	-

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	1101	5PL	OCL-CBL	4.13	1.45	1.33
5	B	1101	5PL	OCK-CBH	-3.41	1.41	1.47
5	B	1101	5PL	OCK-CBU	3.12	1.43	1.34
5	B	1101	5PL	CBM-CBL	2.33	1.57	1.50
5	B	1101	5PL	CBI-CBH	2.25	1.55	1.50

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	1101	5PL	OCK-CBU-CBV	4.36	120.91	111.50
5	B	1101	5PL	OCL-CBL-CBM	3.59	123.17	111.91
5	B	1101	5PL	CBH-OCK-CBU	-3.12	113.86	117.88
5	B	1101	5PL	OCK-CBH-CBI	2.59	112.13	106.13

There are no chirality outliers.

5 of 20 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	B	1101	5PL	CAW-CBH-CBI-OCL
5	B	1101	5PL	OCK-CBH-CBI-OCL
5	B	1101	5PL	CBV-CBU-OCK-CBH
5	B	1101	5PL	OCN-CBU-OCK-CBH
5	B	1101	5PL	CBM-CBL-OCL-CBI


There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	B	1102	C8E	1	0
5	B	1101	5PL	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	482/488 (98%)	0.13	4 (0%) 86 89	40, 63, 94, 133	0
1	C	482/488 (98%)	0.00	1 (0%) 95 97	44, 67, 100, 132	0
1	F	482/488 (98%)	0.04	3 (0%) 89 92	48, 76, 105, 146	0
1	I	482/488 (98%)	0.53	44 (9%) 9 12	68, 115, 152, 186	0
2	B	900/997 (90%)	0.26	19 (2%) 63 67	45, 84, 132, 214	0
2	D	900/997 (90%)	0.63	94 (10%) 6 8	56, 115, 180, 265	0
2	G	900/997 (90%)	0.59	87 (9%) 7 10	53, 115, 182, 253	0
2	J	900/997 (90%)	0.47	58 (6%) 19 23	56, 112, 170, 233	0
3	H	13/13 (100%)	0.64	1 (7%) 13 16	94, 107, 140, 141	0
3	P	13/13 (100%)	0.37	1 (7%) 13 16	74, 93, 108, 121	0
4	E	13/13 (100%)	0.44	1 (7%) 13 16	88, 100, 137, 152	0
4	K	13/13 (100%)	0.73	1 (7%) 13 16	102, 120, 133, 153	0
All	All	5580/5992 (93%)	0.38	314 (5%) 24 28	40, 98, 161, 265	0

The worst 5 of 314 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	J	315	PHE	6.2
1	I	236	TYR	6.0
2	D	132	GLU	5.7
2	D	116	SER	5.3
2	D	536	ASP	5.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
6	C8E	B	1102	13/21	0.84	0.31	66,87,110,110	0
5	5PL	B	1101	20/85	0.87	0.26	87,90,99,100	0
6	C8E	B	1103	13/21	0.93	0.29	36,53,89,92	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around 5PL B 1101:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around C8E B 1103:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers

There are no such residues in this entry.