



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 21, 2020 – 08:52 PM BST

PDB ID : 6SLM  
Title : Crystal structure of full-length HPV31 E6 oncoprotein in complex with LXXLL peptide of ubiquitin ligase E6AP  
Authors : Conrady, M.; Gogl, G.; Cousido-Siah, A.; Mitschler, A.; Trave, G.; Simon, C.  
Deposited on : 2019-08-20  
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.14.6  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.14.6


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	551	 3% 82% 16% ..
2	B	2	 100%

## 2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 4281 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Maltose/maltodextrin-binding periplasmic protein, Protein E6, Ubiquitin-protein ligase E3A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	539	4213	2695	709	794	15	0	2	0

There are 29 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	630	MET	ALA	conflict	UNP P0AEX9
A	712	ALA	ASP	conflict	UNP P0AEX9
A	713	ALA	LYS	conflict	UNP P0AEX9
A	869	ALA	LYS	conflict	UNP P0AEX9
A	989	ALA	GLU	conflict	UNP P0AEX9
A	992	ALA	LYS	conflict	UNP P0AEX9
A	993	ALA	ASP	conflict	UNP P0AEX9
A	997	ASN	ARG	conflict	UNP P0AEX9
A	998	ALA	ILE	conflict	UNP P0AEX9
A	999	ALA	THR	conflict	UNP P0AEX9
A	1000	ALA	LYS	conflict	UNP P0AEX9
A	1097	SER	CYS	conflict	UNP P17386
A	1111	SER	CYS	conflict	UNP P17386
A	1150	GLY	-	linker	UNP P17386
A	1151	SER	-	linker	UNP P17386
A	1152	SER	-	linker	UNP P17386
A	1153	GLY	-	linker	UNP P17386
A	1154	SER	-	linker	UNP P17386
A	1155	GLY	-	linker	UNP P17386
A	1156	SER	-	linker	UNP P17386
A	1157	GLY	-	linker	UNP P17386
A	1158	SER	-	linker	UNP P17386
A	1159	GLY	-	linker	UNP P17386
A	1160	SER	-	linker	UNP P17386
A	1161	GLY	-	linker	UNP P17386
A	1162	SER	-	linker	UNP P17386

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	1163	ALA	-	linker	UNP P17386
A	1164	ALA	-	linker	UNP P17386
A	1165	ALA	-	linker	UNP P17386

- Molecule 2 is an oligosaccharide called alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose.




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
			Total	C	O			
2	B	2	23	12	11	0	0	0

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn) (labeled as "Ligand of Interest" by author).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Zn	0	0
			2	2		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
4	A	1	6	3	3	0	0

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	31	Total	O	0	0
			31	31		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Maltose/maltodextrin-binding periplasmic protein, Protein E6, Ubiquitin-protein ligase E3A



- Molecule 2: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	113.64Å 113.64Å 185.97Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	49.21 – 2.80 49.21 – 2.80	Depositor EDS
% Data completeness (in resolution range)	99.4 (49.21-2.80) 92.7 (49.21-2.80)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.44 (at 2.81Å)	Xtrriage
Refinement program	PHENIX 1.14_3260	Depositor
R, $R_{free}$	0.230 , 0.268 0.231 , 0.270	Depositor DCC
$R_{free}$ test set	894 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	56.6	Xtrriage
Anisotropy	0.274	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 58.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	4281	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	85.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.10% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, ZN, GLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.25	0/4312	0.41	0/5850

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4213	0	4141	55	0
2	B	23	0	21	2	0
3	A	2	0	0	0	0
4	A	12	0	16	2	0
5	A	31	0	0	3	0
All	All	4281	0	4178	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1144:ARG:HG2	1:A:1144:ARG:HH21	1.43	0.84
1:A:637:LEU:HB2	1:A:665:VAL:HG12	1.62	0.79
2:B:1:GLC:O3	2:B:2:GLC:O2	1.60	0.78
1:A:775:SER:HB2	1:A:852:THR:HG22	1.74	0.68
1:A:752:LEU:HD21	1:A:756:PRO:HD3	1.77	0.67
1:A:860:TRP:HB3	5:A:1303:HOH:O	1.95	0.67
1:A:689:ILE:HD11	1:A:906:ALA:HB1	1.77	0.65
1:A:711:PRO:HG2	1:A:716:GLN:OE1	1.98	0.64
1:A:814:ASP:O	1:A:819:LYS:NZ	2.32	0.63
1:A:902:ASN:HB3	1:A:905:LEU:HB3	1.80	0.63
1:A:749:LYS:NZ	5:A:1305:HOH:O	2.31	0.63
1:A:1005:PRO:O	1:A:1010:ARG:NH2	2.32	0.63
1:A:748:ASN:HD22	1:A:870:VAL:HG13	1.65	0.61
1:A:916:THR:HG23	1:A:919:GLY:H	1.66	0.60
1:A:1144:ARG:CG	1:A:1144:ARG:HH21	2.11	0.59
1:A:1072:LYS:NZ	1:A:1148:GLN:CG	2.65	0.59
1:A:778:MET:HB2	1:A:852:THR:HG21	1.85	0.59
1:A:900:SER:O	1:A:903:LYS:NZ	2.36	0.58
1:A:1008:ARG:HG2	1:A:1054:TYR:CZ	2.41	0.56
1:A:774:LYS:HD2	1:A:851:GLU:HA	1.87	0.56
1:A:1102:ARG:HD3	1:A:1107:GLN:HG3	1.90	0.54
1:A:752:LEU:HD11	1:A:765:LEU:HD11	1.90	0.53
1:A:1011:LYS:HE2	1:A:1051:THR:HG22	1.92	0.52
1:A:779:PHE:HB2	5:A:1306:HOH:O	2.10	0.52
1:A:1144:ARG:NH1	4:A:1204:GOL:O2	2.43	0.51
1:A:642:ASN:ND2	1:A:644:ASP:OD1	2.45	0.50
1:A:739:ALA:HA	1:A:932:VAL:HA	1.94	0.50
1:A:797:TYR:HB2	4:A:1203:GOL:H11	1.94	0.50
1:A:1072:LYS:HZ2	1:A:1148:GLN:CG	2.25	0.49
1:A:794:ASP:HB3	1:A:817:GLY:HA2	1.95	0.48
1:A:944:ASP:HB3	1:A:947:ILE:HB	1.94	0.48
1:A:698:GLY:HA3	1:A:962:ASN:O	2.13	0.48
1:A:884:PRO:HB3	1:A:956:LYS:HD2	1.96	0.47
1:A:673:LEU:H	1:A:673:LEU:HD23	1.80	0.47
1:A:689:ILE:HD12	1:A:910:LEU:HD11	1.96	0.47
1:A:695:ASP:OD2	2:B:2:GLC:O3	2.27	0.45
1:A:859:PRO:HA	1:A:862:TRP:CE2	2.51	0.45
1:A:1021:ILE:HG13	1:A:1022:PRO:HD2	1.99	0.45
1:A:709:ILE:HG13	1:A:711:PRO:HD3	2.00	0.44
1:A:727:VAL:HG23	1:A:733:LEU:HD12	2.00	0.44
1:A:634:GLU:HA	1:A:902:ASN:HD21	1.83	0.43
1:A:1144:ARG:HG2	1:A:1144:ARG:NH2	2.21	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:921:GLU:HG2	1:A:925:LYS:HE3	2.00	0.42
1:A:959:ILE:H	1:A:959:ILE:HD12	1.85	0.42
1:A:1144:ARG:CG	1:A:1144:ARG:NH2	2.73	0.42
1:A:1114:GLU:HG2	1:A:1137:ILE:HG13	2.01	0.42
1:A:663:ILE:HG12	1:A:905:LEU:HD21	2.01	0.42
1:A:789:PRO:HG3	1:A:887:PRO:HB3	2.02	0.42
1:A:862:TRP:HB2	1:A:928:PRO:HG2	2.01	0.42
1:A:979:ASN:HB3	1:A:985:GLN:HG2	2.00	0.42
1:A:976:ALA:HB2	1:A:994:ALA:HB2	2.00	0.42
1:A:1142:ARG:O	1:A:1142:ARG:HG3	2.20	0.42
1:A:736:TYR:HB2	1:A:894:ALA:HB3	2.01	0.42
1:A:1012:LEU:HD11	1:A:1042:VAL:HG13	2.03	0.41
1:A:1144:ARG:HG2	1:A:1144:ARG:H	1.46	0.40
1:A:719:LEU:HB2	1:A:724:TRP:HE1	1.87	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	536/551 (97%)	515 (96%)	21 (4%)	0	<a href="#">100</a>   <a href="#">100</a>

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	435/448 (97%)	425 (98%)	10 (2%)	50 82

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	645	LYS
1	A	716	GLN
1	A	757	LYS
1	A	888	PHE
1	A	903	LYS
1	A	1083	VAL
1	A	1105	THR
1	A	1144	ARG
1	A	1145	THR
1	A	1147	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	GLC	B	1	2	12,12,12	1.50	3 (25%)	17,17,17	2.05	3 (17%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	GLC	B	2	2	11,11,12	1.63	4 (36%)	15,15,17	1.70	4 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	B	1	2	-	2/2/22/22	0/1/1/1
2	GLC	B	2	2	-	0/2/19/22	0/1/1/1

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	2	GLC	O5-C5	-2.73	1.37	1.43
2	B	2	GLC	O5-C1	-2.72	1.39	1.43
2	B	1	GLC	O5-C5	-2.59	1.38	1.44
2	B	2	GLC	C1-C2	-2.14	1.47	1.52
2	B	1	GLC	C3-C2	-2.04	1.47	1.52
2	B	2	GLC	C2-C3	-2.04	1.49	1.52
2	B	1	GLC	C4-C5	-2.04	1.48	1.53

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1	GLC	C4-C3-C2	-6.98	98.64	110.82
2	B	2	GLC	O3-C3-C2	-3.94	102.44	109.99
2	B	2	GLC	O5-C1-C2	-2.46	106.98	110.77
2	B	2	GLC	O2-C2-C1	-2.29	104.47	109.15
2	B	1	GLC	O1-C1-O5	-2.26	103.58	110.38
2	B	2	GLC	C6-C5-C4	2.13	117.98	113.00
2	B	1	GLC	C6-C5-C4	-2.11	108.06	113.00

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	1	GLC	O5-C5-C6-O6
2	B	1	GLC	C4-C5-C6-O6

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1	GLC	1	0
2	B	2	GLC	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



## 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	GOL	A	1204	-	5,5,5	0.09	0	5,5,5	0.49	0
4	GOL	A	1203	-	5,5,5	0.93	0	5,5,5	0.99	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	1204	-	-	4/4/4/4	-
4	GOL	A	1203	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	1204	GOL	C1-C2-C3-O3
4	A	1204	GOL	O1-C1-C2-C3
4	A	1203	GOL	C1-C2-C3-O3
4	A	1203	GOL	O2-C2-C3-O3
4	A	1204	GOL	O2-C2-C3-O3
4	A	1204	GOL	O1-C1-C2-O2

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1204	GOL	1	0
4	A	1203	GOL	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	539/551 (97%)	0.17	18 (3%) 46 36	34, 83, 138, 185	0

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1151	SER	4.5
1	A	712	ALA	4.0
1	A	910	LEU	3.6
1	A	739	ALA	3.0
1	A	934	LEU	2.8
1	A	731	GLY	2.7
1	A	1164	ALA	2.6
1	A	722	PHE	2.6
1	A	1143	PRO	2.6
1	A	773	GLY	2.5
1	A	637	LEU	2.4
1	A	914	LEU	2.3
1	A	662	GLY	2.3
1	A	751	LEU	2.2
1	A	913	TYR	2.2
1	A	870	VAL	2.1
1	A	869	ALA	2.0
1	A	1148	GLN	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.




### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	GLC	B	1	12/12	0.92	0.17	61,66,70,76	0
2	GLC	B	2	11/12	0.94	0.15	42,53,61,63	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,




median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	GOL	A	1203	6/6	0.47	0.33	92,94,98,99	0
4	GOL	A	1204	6/6	0.90	0.20	55,69,71,76	0
3	ZN	A	1202	1/1	0.97	0.20	53,53,53,53	0
3	ZN	A	1201	1/1	0.99	0.14	42,42,42,42	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around ZN A 1202:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.