



## wwPDB EM Validation Summary Report ⓘ

Apr 20, 2024 – 12:29 PM EDT

PDB ID : 8SXB  
EMDB ID : EMD-40830  
Title : Bovine multidrug resistance protein 4 (MRP4) bound to prostaglandin E2 in MSP lipid nanodisc  
Authors : Pourmal, S.; Stroud, R.M.  
Deposited on : 2023-05-20  
Resolution : 2.90 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev92  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
MolProbity : 4.02b-467  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
MapQ : 1.9.13  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*ELECTRON MICROSCOPY*

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion  $< 40\%$ ). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1325	

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 18635 atoms, of which 9436 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Multidrug resistance-associated protein 4 isoform X1.

Mol	Chain	Residues	Atoms					AltConf	Trace	
			Total	C	H	N	O			S
1	A	1152	18576	5951	9403	1544	1643	35	0	0

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1375	HIS	UNK	conflict	UNP A0A6P5CNW8
A	1864	ASN	UNK	conflict	UNP A0A6P5CNW8
A	2517	ILE	UNK	conflict	UNP A0A6P5CNW8
A	2600	GLU	UNK	conflict	UNP A0A6P5CNW8
A	2638	PHE	LEU	conflict	UNP A0A6P5CNW8

- Molecule 2 is (Z)-7-[(1R,2R,3R)-3-hydroxy-2-[(E,3S)-3-hydroxyoct-1-enyl]-5-oxo-cyclopentyl]hept-5-enoic acid (three-letter code: P2E) (formula: C<sub>20</sub>H<sub>32</sub>O<sub>5</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				AltConf
			Total	C	H	O	
2	A	1	56	20	31	5	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms			AltConf
			Total	H	O	
3	A	1	3	2	1	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Multidrug resistance-associated protein 4 isoform X1



H2380	A2469	I2535	Q2595	H2381	L2470	Q2536	L2596	H2382	L2471	Q2537	L2597	H2383	E2472	K2538	K2598	H2384	V2473	I2539	A2599	H2385	Q2474	R2540	E2600	K2386	L2475	E2541	A2601	S2387	K2476	K2542	A2602	K2388	E2477	F2543	A2603	E2389	A2478	A2544	L2604	K2390	I2479	H2545	T2605	V2391	E2480	C2546	E2606	G2392	D2481	T2547	T2607	I2393	L2482	V2548	A2608	V2394	L2483	L2549	K2609	G2395	P2483	T2550	Q2610	R2396	G2484	I2551	V2611	T2397	K2485	A2552	Y2612	E2413	M2486	H2553	F2613	F2414	D2487	R2554	F2614	E2415	T2488	L2555	K2615	G2416	E2489	N2556	R2616	K2417	L2490	T2557	Y2617	I2418	A2491	I2558	P2618	D2421	E2492	I2559	ASP	K2422	E2493	D2560	ILE	E2427	G2494	S2561	THR	D2432	S2495	D2562	HIS	D2437	N2496	K2563	ASN	Q2442	F2497	I2564	GLY	E2443	S2498	H2565	HIS	L2446	V2499	M2566	VAL	F2447	G2500	V2566	VAL	T2448	Q2501	L2567	MET	G2449	R2502	D2568	ASN	R2452	Q2503	S2569	ALA	R2453	L2504	SER	SER	N2459	V2505	K2570	GLY	E2460	C2506	R2571	GLN	H2461	A2510	L2572	PRO	S2462	R2513	K2573	SER	D2463	K2514	E2574	ALA	E2464	I2517	Y2575	PHE	E2465	R2517	D2576	THR	L2466	D2521	E2577	ILE	W2467	E2522	P2578	PHE	N2468	A2523	Y2579	GLU	E2469	T2524	V2580	THR	S2470	T2525	L2581	ALA	S2471	A2526	L2582	LEU	E2472	N2526	Q2583		Q2474	V2527	N2584		L2475	D2528	R2585		K2476	P2529	D2586		E2477	R2530	S2587		A2478	T2531	L2588		I2479	D2532	F2589		E2480	E2533	Y2590		D2481	L2534	K2591		L2482	T2535	M2592		P2483	L2536	V2593		G2484	L2537	Q2594		K2485	L2538			M2486	L2539			D2487	L2540			T2488	L2541			E2489	L2542			L2490	L2543			A2491	L2544			E2492	L2545			S2493	L2546			G2494	L2547			S2495	L2548			N2496	L2549			F2497	L2550			S2498	L2551			V2499	L2552			G2500	L2553			Q2501	L2554			R2502	L2555			Q2503	L2556			L2504	L2557			V2505	L2558			C2506	L2559			A2510	L2560			R2513	L2561			K2514	L2562			I2517	L2563			D2521	L2564			E2522	L2565			A2523	L2566			T2524	L2567			A2525	L2568			N2526	L2569			V2527	L2570			D2528	L2571			P2529	L2572			R2530	L2573			T2531	L2574			D2532	L2575			E2533	L2576			L2534	L2577			L2535	L2578			L2536	L2579			L2537	L2580			L2538	L2581			L2539	L2582			L2540	L2583			L2541	L2584			L2542	L2585			L2543	L2586			L2544	L2587			L2545	L2588			L2546	L2589			L2547	L2590			L2548	L2591			L2549	L2592			L2550	L2593			L2551	L2594		
-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-----	-------	-------	-------	-----	-------	-------	-------	-----	-------	-------	-------	-----	-------	-------	-------	-----	-------	-------	-------	-----	-------	-------	-------	-----	-------	-------	-------	-----	-------	-------	-------	-----	-------	-------	-------	-----	-------	-------	-------	-----	-------	-------	-------	-----	-------	-------	-----	-----	-------	-------	-------	-----	-------	-------	-------	-----	-------	-------	-------	-----	-------	-------	-------	-----	-------	-------	-------	-----	-------	-------	-------	-----	-------	-------	-------	-----	-------	-------	-------	-----	-------	-------	-------	-----	-------	-------	-------	-----	-------	-------	-------	-----	-------	-------	-------	-----	-------	-------	-------	-----	-------	-------	-------	--	-------	-------	-------	--	-------	-------	-------	--	-------	-------	-------	--	-------	-------	-------	--	-------	-------	-------	--	-------	-------	-------	--	-------	-------	-------	--	-------	-------	-------	--	-------	-------	-------	--	-------	-------	-------	--	-------	-------	-------	--	-------	-------	--	--	-------	-------	--	--	-------	-------	--	--	-------	-------	--	--	-------	-------	--	--	-------	-------	--	--	-------	-------	--	--	-------	-------	--	--	-------	-------	--	--	-------	-------	--	--	-------	-------	--	--	-------	-------	--	--	-------	-------	--	--	-------	-------	--	--	-------	-------	--	--	-------	-------	--	--	-------	-------	--	--	-------	-------	--	--	-------	-------	--	--	-------	-------	--	--	-------	-------	--	--	-------	-------	--	--	-------	-------	--	--	-------	-------	--	--	-------	-------	--	--	-------	-------	--	--	-------	-------	--	--	-------	-------	--	--	-------	-------	--	--	-------	-------	--	--	-------	-------	--	--	-------	-------	--	--	-------	-------	--	--	-------	-------	--	--	-------	-------	--	--	-------	-------	--	--	-------	-------	--	--	-------	-------	--	--	-------	-------	--	--	-------	-------	--	--	-------	-------	--	--	-------	-------	--	--	-------	-------	--	--	-------	-------	--	--	-------	-------	--	--	-------	-------	--	--	-------	-------	--	--	-------	-------	--	--	-------	-------	--	--	-------	-------	--	--	-------	-------	--	--	-------	-------	--	--	-------	-------	--	--	-------	-------	--	--	-------	-------	--	--	-------	-------	--	--	-------	-------	--	--

## 4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	187047	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	66	Depositor
Minimum defocus (nm)	500	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	105000	Depositor
Image detector	GATAN K3 BIOQUANTUM (6k x 4k)	Depositor
Maximum map value	3.115	Depositor
Minimum map value	-2.620	Depositor
Average map value	0.002	Depositor
Map value standard deviation	0.066	Depositor
Recommended contour level	0.55	Depositor
Map size (Å)	250.5, 250.5, 250.5	wwPDB
Map dimensions	300, 300, 300	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.835, 0.835, 0.835	Depositor

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: P2E

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.32	0/9365	0.56	0/12693

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	9173	9403	9398	17	0
2	A	25	31	31	0	0
3	A	1	2	0	0	0
All	All	9199	9436	9429	17	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

The worst 5 of 17 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:2088:TYR:O	1:A:2092:THR:HG23	1.92	0.70

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:2127:ARG:NH2	1:A:2342:GLU:O	2.29	0.64
1:A:2522:GLU:HG2	1:A:2552:ALA:HA	1.89	0.54
1:A:1423:GLU:OE1	1:A:1537:ALA:HB1	2.09	0.52
1:A:1698:GLU:O	1:A:1701:VAL:HG22	2.13	0.48

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	1144/1325 (86%)	1121 (98%)	23 (2%)	0	100   100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	999/1156 (86%)	999 (100%)	0	100   100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1571	GLN
1	A	2254	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	P2E	A	2701	-	25,25,25	0.71	1 (4%)	26,31,31	1.10	3 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	P2E	A	2701	-	-	4/19/35/35	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	2701	P2E	O2-C1	-2.69	1.21	1.30

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	2701	P2E	C16-C15-C14	-2.52	107.18	111.60
2	A	2701	P2E	O2-C1-C2	2.18	121.03	114.03
2	A	2701	P2E	C11-C10-C9	-2.15	102.65	105.56

There are no chirality outliers.


All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	2701	P2E	C13-C14-C15-O5
2	A	2701	P2E	C13-C14-C15-C16
2	A	2701	P2E	C16-C17-C18-C19
2	A	2701	P2E	O2-C1-C2-C3

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.


## 6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-40830. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

### 6.1 Orthogonal projections [i](#)


#### 6.1.1 Primary map



X




Y




Z


#### 6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

## 6.2 Central slices [i](#)


### 6.2.1 Primary map



X Index: 150




Y Index: 150




Z Index: 150


### 6.2.2 Raw map



X Index: 150



Y Index: 150




Z Index: 150


The images above show central slices of the map in three orthogonal directions.

## 6.3 Largest variance slices [i](#)


### 6.3.1 Primary map



X Index: 149




Y Index: 142




Z Index: 140


### 6.3.2 Raw map



X Index: 149



Y Index: 143




Z Index: 152


The images above show the largest variance slices of the map in three orthogonal directions.

## 6.4 Orthogonal standard-deviation projections (False-color) [i](#)


### 6.4.1 Primary map



X




Y




Z


### 6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.



## 6.5 Orthogonal surface views [i](#)

### 6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.55. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

### 6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


## 6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

## 7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

### 7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


## 7.2 Volume estimate [i](#)



The volume at the recommended contour level is 31 nm<sup>3</sup>; this corresponds to an approximate mass of 28 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

### 7.3 Rotationally averaged power spectrum [i](#)




\*Reported resolution corresponds to spatial frequency of 0.345 Å<sup>-1</sup>

## 8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

### 8.1 FSC [i](#)



\*Reported resolution corresponds to spatial frequency of  $0.345 \text{\AA}^{-1}$

## 8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	2.90	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	3.44	4.01	3.52

\*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.44 differs from the reported value 2.9 by more than 10 %

## 9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-40830 and PDB model 8SXB. Per-residue inclusion information can be found in section 3 on page 5.

### 9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.55 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

## 9.2 Q-score mapped to coordinate model [\(i\)](#)



The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.


## 9.3 Atom inclusion mapped to coordinate model [\(i\)](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.55).







## 9.4 Atom inclusion [i](#)



At the recommended contour level, 53% of all backbone atoms, 52% of all non-hydrogen atoms, are inside the map.

## 9.5 Map-model fit summary [i](#)

The table lists the average atom inclusion at the recommended contour level (0.55) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.5220	 0.4350
A	 0.5190	 0.4350

