

wwPDB X-ray Structure Validation Summary Report (i)

Sep 23, 2023 – 07:55 PM EDT

PDB ID : 5SYT

Title: Crystal Structure of ZMPSTE24

Authors: Clark, K.; Jenkins, J.L.; Fedoriw, N.; Dumont, M.E.; Membrane Protein Struc-

tural Biology Consortium (MPSBC)

Deposited on : 2016-08-11

Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.35.1

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# \mathrm{Entries})$	$(\# ext{Entries}, ext{ resolution range}(ext{Å}))$
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
			3%	
1	A	480	79%	12% • 8%



2 Entry composition (i)

There are 10 unique types of molecules in this entry. The entry contains 7938 atoms, of which 3932 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


• Molecule 1 is a protein called CAAX prenyl protease 1 homolog.

Mol	Chain	Residues			Atom	ıs			ZeroOcc	AltConf	Trace
1	A	444	Total 7123	C 2380	H 3554	N 556	O 621	S 12	0	1	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	475	SER	-	expression tag	UNP O75844
A	476	GLY	-	expression tag	UNP O75844
A	477	LEU	-	expression tag	UNP O75844
A	478	GLU	-	expression tag	UNP O75844
A	479	VAL	-	expression tag	UNP O75844
A	480	LEU	-	expression tag	UNP O75844


• Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0


 \bullet Molecule 3 is 1,2-DIACYL-SN-GLYCERO-3-PHOSPHOCHOLINE (three-letter code: PC1) (formula: C44H88NO8P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
3	A	1	Total	C 21	H 40	N	0	P	0	0


• Molecule 4 is PENTAETHYLENE GLYCOL MONODECYL ETHER (three-letter code: CXE) (formula: $C_{20}H_{42}O_6$).





Mol	Chain	Residues	A	A ton	ns		ZeroOcc	AltConf
1	٨	1	Total	С	Н	О	0	0
4	А	1	61	18	37	6	0	U

• Molecule 5 is (HYDROXYETHYLOXY)TRI(ETHYLOXY)OCTANE (three-letter code: C8E) (formula: $C_{16}H_{34}O_5$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
5	Λ	1	Total	С	Н	О	0	0	
9	A	1	55	16	34	5	0	U	
5	Λ	1	Total	С	Н	О	0	0	
0	A	1	30	8	17	5	U	U	

Continued on next page...



 $Continued\ from\ previous\ page...$


Mol	Chain	Residues	A	ton	ns		ZeroOcc	AltConf
5	A	1	Total	С	Н	О	0	0
	Λ	1	17	7	6	4	U	U
5	A	1	Total	\mathbf{C}	Η	O	0	0
	11	1	55	16	34	5	0	U
5	A	1	Total	С	Η	O	0	0
	71	1	26	7	15	4	0	0
5	A	1	Total	\mathbf{C}	Η	O	0	0
	71	1	30	8	17	5	0	0
5	A	1	Total	С	Η	Ο	0	0
	71	1	39	11	23	5	Ŭ .	
5	A	1	Total	\mathbf{C}	Η	Ο	0	0
		1	50	15	31	4	0	
5	A	1	Total	\mathbf{C}	Η	O	0	0
		1	26	7	15	4	0	
5	A	1	Total	С	Η	Ο	0	0
	11	1	29	8	17	4	0	
5	A	1	Total	С	Η	Ο	0	0
	11	1	25	7	14	4	Ü	V
5	A	1	Total	С	Η	Ο	0	0
	11	1	55	16	34	5		

• Molecule 6 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total 1	Zn 1	0	0


• Molecule 7 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).





Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1	Total C H (0	0
7	A	1	Total C H (O	0	0


 $\bullet \ \, \text{Molecule 8 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: $C_4H_{10}O_3$)}. \\$



Mol	Chain	Residues	A	${f Atoms}$				AltConf
Q	Λ	1	Total	С	Н	О	0	0
0	Λ	1	17	4	10	3	0	

 \bullet Molecule 9 is DIMETHYL SULFOXIDE (three-letter code: DMS) (formula: $\mathrm{C_2H_6OS}).$





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf		
0	Λ	1	Total	С	Н	О	S	0	0
9	A	1	10	2	6	1	1	0	U
0	Λ	1	Total	С	Н	О	S	0	0
9	A	1	10	2	6	1	1	0	U
0	Λ	1	Total	С	Н	О	S	0	0
9	A	1	10	2	6	1	1	0	0

• Molecule 10 is water.


\mathbf{Mol}	Chain	Residues	${f Atoms}$	ZeroOcc	AltConf
10	A	158	Total O 160 160	0	2



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: CAAX prenyl protease 1 homolog





4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants	149.46Å 84.56Å 76.89Å	Domositon
a, b, c, α , β , γ	90.00° 119.07° 90.00°	Depositor
Resolution (Å)	33.60 - 2.00	Depositor
Resolution (A)	33.60 - 2.00	EDS
% Data completeness	99.7 (33.60-2.00)	Depositor
(in resolution range)	93.7 (33.60-2.00)	EDS
R_{merge}	0.44	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.66 (at 2.00Å)	Xtriage
Refinement program	PHENIX 1.10.1_2155	Depositor
D.D.	0.218 , 0.249	Depositor
R, R_{free}	0.226 , 0.256	DCC
R_{free} test set	1997 reflections (3.54%)	wwPDB-VP
Wilson B-factor (Å ²)	19.7	Xtriage
Anisotropy	0.683	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.41,62.4	EDS
L-test for twinning ²	$< L > = 0.47, < L^2> = 0.30$	Xtriage
Estimated twinning fraction	0.037 for -h-2*l,-k,l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	7938	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 5.97% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, DMS, SO4, PC1, GOL, CXE, C8E, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond	lengths	Bond angles	
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.58	0/3671	0.68	1/4983 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}(^{o})$
1	A	330	LEU	CA-CB-CG	5.79	128.63	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3569	3554	3549	46	1
2	A	10	0	0	0	0
3	A	31	40	36	4	1
4	A	24	37	35	2	0
5	A	180	257	252	6	0
6	A	1	0	0	0	0
7	A	12	16	16	4	0
8	A	7	10	10	0	0
9	A	12	18	18	0	0
10	A	160	0	0	7	0

Continued on next page...



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	4006	3932	3916	48	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 48 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
1:A:475:SER:OG	1:A:476:GLY:N	2.13	0.81
1:A:265:ASN:ND2	10:A:602:HOH:O	2.20	0.73
1:A:314:LYS:HA	1:A:315:ALA:HB3	1.74	0.68
1:A:43:ARG:NH2	3:A:503:PC1:O12	2.29	0.65
1:A:397:SER:HB2	1:A:398:PRO:HD3	1.84	0.60

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	$\begin{array}{c} \text{Clash} \\ \text{overlap } (\text{\AA}) \end{array}$
1:A:121:THR:OG1	3:A:503:PC1:O14[4_346]	1.97	0.23

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mo	l Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	439/480 (92%)	422 (96%)	15 (3%)	2 (0%)	29 23

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	313	ILE
1	A	318	LYS



5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	378/424 (89%)	361 (96%)	17 (4%)	27 24

5 of 17 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	380	ASP
1	A	477	LEU
1	A	236	GLU
1	A	250	THR
1	A	313	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 23 ligands modelled in this entry, 1 is monoatomic - leaving 22 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The



Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Во	ond leng	ths	В	ond ang	gles
MIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	GOL	A	519	-	5,5,5	0.45	0	5,5,5	0.51	0
5	C8E	A	506	-	12,12,20	0.46	0	11,11,19	0.50	0
5	C8E	A	510	-	12,12,20	0.51	0	11,11,19	0.49	0
5	C8E	A	512	_	18,18,20	0.33	0	17,17,19	0.71	0
5	C8E	A	511	-	15,15,20	0.59	0	14,14,19	0.46	0
5	C8E	A	513	_	10,10,20	0.45	0	9,9,19	0.39	0
3	PC1	A	503	-	30,30,53	1.27	2 (6%)	36,38,61	1.54	5 (13%)
9	DMS	A	522	-	3,3,3	0.75	0	3,3,3	0.99	0
9	DMS	A	521	-	3,3,3	0.61	0	3,3,3	0.69	0
5	C8E	A	516	-	20,20,20	0.36	0	19,19,19	0.81	1 (5%)
5	C8E	A	509	-	10,10,20	0.37	0	9,9,19	0.47	0
5	C8E	A	505	-	20,20,20	0.46	0	19,19,19	0.44	0
7	GOL	A	518	-	5,5,5	0.38	0	5,5,5	0.87	0
5	C8E	A	515	-	10,10,20	0.47	0	9,9,19	0.31	0
8	PEG	A	520	_	6,6,6	0.56	0	5,5,5	0.82	0
2	SO4	A	502	-	4,4,4	0.20	0	6,6,6	0.38	0
2	SO4	A	501	-	4,4,4	0.20	0	6,6,6	0.41	0
5	C8E	A	507	-	10,10,20	0.41	0	9,9,19	0.33	0
5	C8E	A	508	-	20,20,20	0.44	0	19,19,19	0.49	0
5	C8E	A	514	-	11,11,20	0.48	0	10,10,19	0.58	0
4	CXE	A	504	-	23,23,25	0.51	0	22,22,24	0.52	0
9	DMS	A	523	_	3,3,3	0.67	0	3,3,3	1.03	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	C8E	A	516	-	-	9/18/18/18	-
5	C8E	A	509	-	-	4/8/8/18	-
5	C8E	A	505	-	-	12/18/18/18	-
5	C8E	A	508	-	-	9/18/18/18	-
5	C8E	A	515	-	-	1/8/8/18	-
5	C8E	A	514	-	-	8/9/9/18	_
7	GOL	A	518	-	-	3/4/4/4	-

Continued on next page...



Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	GOL	A	519	-	-	4/4/4/4	-
5	C8E	A	506	-	-	4/10/10/18	-
8	PEG	A	520	-	-	2/4/4/4	-
4	CXE	A	504	-	-	11/21/21/23	-
5	C8E	A	510	-	-	6/10/10/18	-
5	C8E	A	512	-	-	8/16/16/18	-
5	C8E	A	511	-	-	4/13/13/18	-
5	C8E	A	513	-	-	3/8/8/18	-
3	PC1	A	503	-	-	11/34/34/57	-
5	C8E	A	507	-	-	5/8/8/18	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(A)	$\operatorname{Ideal}(ext{\AA})$
3	A	503	PC1	O31-C31	3.42	1.43	1.33
3	A	503	PC1	O21-C21	3.19	1.43	1.34

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
3	A	503	PC1	O21-C21-C22	4.49	121.19	111.50
3	A	503	PC1	O31-C31-C32	3.79	121.31	111.38
3	A	503	PC1	C23-C22-C21	-3.39	101.28	113.62
5	A	516	C8E	O12-C11-C10	2.31	120.82	110.39
3	A	503	PC1	C15-N-C13	2.17	114.55	108.97

There are no chirality outliers.

5 of 104 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	503	PC1	O13-C11-C12-N
7	A	518	GOL	O1-C1-C2-C3
5	A	516	C8E	C10-C11-O12-C13
5	A	506	C8E	C13-C14-O15-C16
5	A	507	C8E	C16-C17-O18-C19


There are no ring outliers.

7 monomers are involved in 17 short contacts:




Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	A	519	GOL	2	0
5	A	510	C8E	3	0
5	A	511	C8E	2	0
3	A	503	PC1	4	1
5	A	516	C8E	1	0
7	A	518	GOL	2	0
4	A	504	CXE	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











5.7 Other polymers (i)

There are no such residues in this entry.



5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mo	ol Chain	nain Analysed	<RSRZ $>$	# RSRZ > 2		$OWAB(A^2)$	Q < 0.9
1	A	A 444/480 (92%)	0.05	16 (3%) 42	42	19, 32, 68, 96	0

The worst 5 of 16 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	11	TRP	4.9
1	A	317	VAL	4.4
1	A	319	ASN	4.4
1	A	475	SER	4.0
1	A	480	LEU	3.9

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
5	C8E	A	511	16/21	0.77	0.33	28,55,69,79	0
5	C8E	A	514	12/21	0.77	0.22	44,69,83,85	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}(\mathring{\mathbf{A}}^2)$	Q<0.9
7	GOL	A	519	6/6	0.77	0.24	68,90,105,117	0
5	C8E	A	506	13/21	0.80	0.16	41,57,66,69	0
3	PC1	A	503	31/54	0.81	0.18	30,55,74,78	0
5	C8E	A	505	21/21	0.82	0.15	32,56,84,84	0
9	DMS	A	522	4/4	0.82	0.20	32,40,53,82	0
4	CXE	A	504	24/26	0.83	0.15	33,54,70,85	0
8	PEG	A	520	7/7	0.83	0.17	46,55,65,67	0
5	C8E	A	508	21/21	0.83	0.18	42,56,70,72	0
5	C8E	A	509	11/21	0.86	0.14	36,54,71,74	0
5	C8E	A	510	13/21	0.87	0.14	34,44,54,56	0
5	C8E	A	516	21/21	0.87	0.12	30,41,55,63	0
7	GOL	A	518	6/6	0.87	0.14	35,42,48,50	0
5	C8E	A	513	11/21	0.89	0.21	43,58,76,79	0
5	C8E	A	507	11/21	0.89	0.17	41,49,53,62	0
5	C8E	A	515	11/21	0.89	0.12	41,53,74,80	0
5	C8E	A	512	19/21	0.89	0.12	31,44,67,68	0
9	DMS	A	523	4/4	0.89	0.14	47,61,96,96	0
2	SO4	A	502	5/5	0.92	0.24	47,58,71,83	0
2	SO4	A	501	5/5	0.92	0.30	62,73,88,92	0
9	DMS	A	521	4/4	0.93	0.25	46,56,92,92	0
6	ZN	A	517	1/1	0.99	0.11	30,30,30,30	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around C8E A 514:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around C8E A 506:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around PC1 A 503:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around C8E A 505: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c \ (\mathrm{at}\ 0.7\ \mathrm{rmsd}) \ \mathrm{in}\ \mathrm{gray}$

 ${
m mF}_o{
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around CXE A 504:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)











Electron density around C8E A 509: 2mF_o-DF_c (at 0.7 rmsd) in gray mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)




Electron density around C8E A 510:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around C8E A 516:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around C8E A 513:

 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around C8E A 507:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around C8E A 515: 2mF_o-DF_c (at 0.7 rmsd) in gray mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)

Electron density around C8E A 512:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









6.5 Other polymers (i)

There are no such residues in this entry.

