



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 20, 2020 – 06:23 PM BST

PDB ID : 5T05
Title : Crystal structure of heparan sulfate 6-O-sulfotransferase with bound PAP and IdoA2S containing hexasaccharide substrate
Authors : Pedersen, L.C.; Moon, A.F.; krahn, J.M.; Liu, J.
Deposited on : 2016-08-15
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2580 (1.96-1.96)
Clashscore	141614	2705 (1.96-1.96)
Ramachandran outliers	138981	2678 (1.96-1.96)
Sidechain outliers	138945	2678 (1.96-1.96)
RSRZ outliers	127900	2539 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	692	
1	B	692	
2	C	6	
2	E	6	
3	D	4	
3	F	4	

2 Entry composition

There are 10 unique types of molecules in this entry. The entry contains 12096 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called maltose binding protein - heparan sulfate 6-O-sulfotransferase isoform 3 fusion protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	670	5248	3342	892	988	26	0	2	0
1	B	677	5306	3380	900	999	27	0	7	0

There are 26 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	initiating methionine	UNP P0AEY0
A	82	ALA	ASP	engineered mutation	UNP P0AEY0
A	83	ALA	LYS	engineered mutation	UNP P0AEY0
A	172	ALA	GLU	engineered mutation	UNP P0AEY0
A	173	ALA	ASN	engineered mutation	UNP P0AEY0
A	239	ALA	LYS	engineered mutation	UNP P0AEY0
A	359	ALA	GLU	engineered mutation	UNP P0AEY0
A	362	ALA	LYS	engineered mutation	UNP P0AEY0
A	363	ALA	ASP	engineered mutation	UNP P0AEY0
A	367	ASN	-	linker	UNP P0AEY0
A	368	ALA	-	linker	UNP P0AEY0
A	369	ALA	-	linker	UNP P0AEY0
A	370	ALA	-	linker	UNP P0AEY0
B	0	MET	-	initiating methionine	UNP P0AEY0
B	82	ALA	ASP	engineered mutation	UNP P0AEY0
B	83	ALA	LYS	engineered mutation	UNP P0AEY0
B	172	ALA	GLU	engineered mutation	UNP P0AEY0
B	173	ALA	ASN	engineered mutation	UNP P0AEY0
B	239	ALA	LYS	engineered mutation	UNP P0AEY0
B	359	ALA	GLU	engineered mutation	UNP P0AEY0
B	362	ALA	LYS	engineered mutation	UNP P0AEY0
B	363	ALA	ASP	engineered mutation	UNP P0AEY0
B	367	ASN	-	linker	UNP P0AEY0
B	368	ALA	-	linker	UNP P0AEY0

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	369	ALA	-	linker	UNP P0AEY0
B	370	ALA	-	linker	UNP P0AEY0

- Molecule 2 is an oligosaccharide called 2-deoxy-2-(sulfoamino)-alpha-D-glucopyranose-(1-4)-beta-D-glucopyranuronic acid-(1-4)-2-deoxy-2-(sulfoamino)-alpha-D-glucopyranose-(1-4)-2-O-sulfo-alpha-L-idopyranuronic acid-(1-4)-2-deoxy-2-(sulfoamino)-alpha-D-glucopyranose-(1-4)-beta-D-glucopyranuronic acid.




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	6	Total	C	N	O	S	0	0	0
			85	36	3	42	4			
2	E	6	Total	C	N	O	S	0	4	0
			100	42	4	49	5			

- Molecule 3 is an oligosaccharide called alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose.




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	Trace	
3	D	4	Total	C	O	0	0	0
			45	24	21			
3	F	4	Total	C	O	0	0	0
			45	24	21			

- Molecule 4 is ADENOSINE-3'-5'-DIPHOSPHATE (three-letter code: A3P) (formula: C₁₀H₁₅N₅O₁₀P₂).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	N	O			P
4	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
4	B	1	Total	C	N	O	P	0	0
			27	10	5	10	2		

- Molecule 5 is P-NITROPHENOL (three-letter code: NPO) (formula: $C_6H_5NO_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
5	A	1	Total	C	N	O	0	0
			10	6	1	3		
5	B	1	Total	C	N	O	0	1
			8	6	1	1		

- Molecule 6 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	B	1	Total C O 4 2 2	0	0
6	B	1	Total C O 4 2 2	0	0
6	B	1	Total C O 4 2 2	0	0
6	B	1	Total C O 4 2 2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	C	O	0	0
			4	2	2		
6	B	1	Total	C	O	0	0
			4	2	2		


- Molecule 7 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	B	1	Total	Cl	0	0
			1	1		
7	A	3	Total	Cl	0	0
			3	3		

- Molecule 8 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	B	2	Total	Na	0	0
			2	2		
8	A	5	Total	Na	0	0
			5	5		

- Molecule 9 is L(+)-TARTARIC ACID (three-letter code: TLA) (formula: C₄H₆O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	A	1	Total	C	O	0	0
			10	4	6		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	A	1	Total C O 10 4 6	0	0
9	B	1	Total C O 10 4 6	0	0
9	B	1	Total C O 10 4 6	0	0


- Molecule 10 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
10	A	553	Total O 563 563	0	10
10	B	514	Total O 521 521	0	8


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: maltose binding protein - heparan sulfate 6-O-sulfotransferase isoform 3 fusion protein



- Molecule 1: maltose binding protein - heparan sulfate 6-O-sulfotransferase isoform 3 fusion protein



- Molecule 2: 2-deoxy-2-(sulfoamino)-alpha-D-glucopyranose-(1-4)-beta-D-glucopyranuronic acid-(1-4)-2-deoxy-2-(sulfoamino)-alpha-D-glucopyranose-(1-4)-2-O-sulfo-alpha-L-idopyranuronic acid-(1-4)-2-deoxy-2-(sulfoamino)-alpha-D-glucopyranose-(1-4)-beta-D-glucopyranuronic acid



- Molecule 2: 2-deoxy-2-(sulfoamino)-alpha-D-glucopyranose-(1-4)-beta-D-glucopyranuronic acid-(1-4)-2-deoxy-2-(sulfoamino)-alpha-D-glucopyranose-(1-4)-2-O-sulfo-alpha-L-idopyranuronic acid-(1-4)-2-deoxy-2-(sulfoamino)-alpha-D-glucopyranose-(1-4)-beta-D-glucopyranuronic acid



BDP1
GNS2
IDS3
GNS4
BDP5
GNS6

- Molecule 3: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose

Chain D:  50% 50%

GLC1
GLC2
GLC3
GLC4

- Molecule 3: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose

Chain F:  50% 50%

GLC1
GLC2
GLC3
GLC4

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	75.78Å 128.16Å 178.71Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.49 – 1.95 48.93 – 1.95	Depositor EDS
% Data completeness (in resolution range)	99.7 (38.49-1.95) 95.4 (48.93-1.95)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.50 (at 1.95Å)	Xtrriage
Refinement program	PHENIX 1.8.2_1309	Depositor
R, R_{free}	0.178 , 0.212 0.178 , 0.212	Depositor DCC
R_{free} test set	2811 reflections (2.22%)	wwPDB-VP
Wilson B-factor (Å ²)	34.0	Xtrriage
Anisotropy	0.232	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 46.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	12096	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.06% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GNS, NPO, A3P, CL, NA, GLC, IDS, EDO, TLA, BDP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.58	0/5371	0.64	0/7291
1	B	0.58	0/5431	0.63	0/7378
All	All	0.58	0/10802	0.64	0/14669

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5248	0	5034	7	0
1	B	5306	0	5040	6	0
2	C	85	0	39	0	0
2	E	100	0	46	0	0
3	D	45	0	39	0	0
3	F	45	0	39	0	0
4	A	27	0	11	0	0
4	B	27	0	11	0	0
5	A	10	0	4	0	0
5	B	8	0	4	0	0
6	A	36	0	54	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	24	0	36	1	0
7	A	3	0	0	1	0
7	B	1	0	0	0	0
8	A	5	0	0	0	0
8	B	2	0	0	0	0
9	A	20	0	8	1	0
9	B	20	0	8	0	0
10	A	563	0	0	3	0
10	B	521	0	0	1	0
All	All	12096	0	10373	15	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (15) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:53:THR:OG1	1:A:54:GLY:N	2.38	0.55
1:B:1342:ARG:NE	10:B:1517:HOH:O	2.42	0.53
1:A:1299:LEU:HD11	1:A:1358:LYS:HA	1.92	0.51
1:A:1092:LYS:O	1:A:1140:LYS:NZ	2.45	0.49
1:B:1299:LEU:HD11	1:B:1358:LYS:HA	1.94	0.49
9:A:1431:TLA:O1	10:A:1501:HOH:O	2.20	0.48
1:B:68:GLY:HA3	1:B:332:ASN:O	2.14	0.48
1:A:1235:ASP:OD2	10:A:1502:HOH:O	2.20	0.47
1:A:229:PRO:HA	1:A:232:TRP:CE2	2.50	0.47
1:A:68:GLY:HA3	1:A:332:ASN:O	2.16	0.46
7:A:1423:CL:CL	10:A:1972:HOH:O	2.58	0.46
1:B:307:TYR:CE2	1:B:311:LEU:HD11	2.53	0.43
1:B:1292:LYS:HE2	6:B:1416:EDO:O1	2.20	0.41
1:A:1229:THR:HG22	1:A:1231:TYR:H	1.87	0.40
1:B:89:LEU:HD12	1:B:304:LEU:HA	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	668/692 (96%)	656 (98%)	11 (2%)	1 (0%)	51	43
1	B	679/692 (98%)	669 (98%)	10 (2%)	0	100	100
All	All	1347/1384 (97%)	1325 (98%)	21 (2%)	1 (0%)	51	43

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	53	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	536/582 (92%)	532 (99%)	4 (1%)	84	82
1	B	536/582 (92%)	532 (99%)	4 (1%)	84	82
All	All	1072/1164 (92%)	1064 (99%)	8 (1%)	81	82

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	258	PHE
1	A	1192	ASP
1	A	1246	ASP
1	A	1339	MET
1	B	55	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	258	PHE
1	B	1192	ASP
1	B	1338	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

21 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	BDP	C	1	2,5	9,12,13	0.98	1 (11%)	12,17,19	0.82	1 (8%)
2	GNS	C	2	2	14,15,16	0.63	0	17,22,24	1.27	2 (11%)
2	IDS	C	3	8,2	13,16,17	0.89	0	15,24,26	0.75	1 (6%)
2	GNS	C	4	8,2	14,15,16	0.52	0	17,22,24	1.13	1 (5%)
2	BDP	C	5	8,2	9,12,13	0.89	0	12,17,19	0.90	0
2	GNS	C	6	2	14,15,16	0.50	0	17,22,24	1.49	2 (11%)
3	GLC	D	1	3	12,12,12	0.45	0	17,17,17	0.63	0
3	GLC	D	2	3	11,11,12	0.58	0	15,15,17	0.84	0
3	GLC	D	3	3	11,11,12	0.71	0	15,15,17	1.01	1 (6%)
3	GLC	D	4	3	11,11,12	0.52	0	15,15,17	1.34	1 (6%)
2	BDP	E	1[A]	2,5	9,12,13	0.96	1 (11%)	12,17,19	0.99	1 (8%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GNS	E	2[A]	2	14,15,16	0.64	0	17,22,24	1.44	2 (11%)
2	IDS	E	3[A]	2	13,16,17	0.90	0	15,24,26	1.13	1 (6%)
2	GNS	E	4[A]	2	14,15,16	0.67	0	17,22,24	1.71	5 (29%)
2	GNS	E	4[B]	2	14,15,16	0.61	0	17,22,24	1.26	1 (5%)
2	BDP	E	5	2	9,12,13	0.94	1 (11%)	12,17,19	1.44	1 (8%)
2	GNS	E	6	2	14,15,16	0.58	0	17,22,24	1.86	3 (17%)
3	GLC	F	1	3	12,12,12	0.50	0	17,17,17	0.60	0
3	GLC	F	2	3	11,11,12	0.52	0	15,15,17	0.85	0
3	GLC	F	3	3	11,11,12	0.68	0	15,15,17	0.93	1 (6%)
3	GLC	F	4	3	11,11,12	0.54	0	15,15,17	1.17	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BDP	C	1	2,5	-	0/0/21/24	0/1/1/1
2	GNS	C	2	2	-	0/7/24/27	0/1/1/1
2	IDS	C	3	8,2	-	0/5/26/29	0/1/1/1
2	GNS	C	4	8,2	-	0/7/24/27	0/1/1/1
2	BDP	C	5	8,2	-	0/0/21/24	0/1/1/1
2	GNS	C	6	2	-	0/7/24/27	0/1/1/1
3	GLC	D	1	3	-	0/2/22/22	0/1/1/1
3	GLC	D	2	3	-	0/2/19/22	0/1/1/1
3	GLC	D	3	3	-	0/2/19/22	0/1/1/1
3	GLC	D	4	3	-	0/2/19/22	0/1/1/1
2	BDP	E	1[A]	2,5	-	0/0/21/24	0/1/1/1
2	GNS	E	2[A]	2	-	1/7/24/27	0/1/1/1
2	IDS	E	3[A]	2	-	0/5/26/29	0/1/1/1
2	GNS	E	4[A]	2	-	1/7/24/27	0/1/1/1
2	GNS	E	4[B]	2	-	0/7/24/27	0/1/1/1
2	BDP	E	5	2	-	0/0/21/24	0/1/1/1
2	GNS	E	6	2	-	0/7/24/27	0/1/1/1
3	GLC	F	1	3	-	0/2/22/22	0/1/1/1
3	GLC	F	2	3	-	0/2/19/22	0/1/1/1
3	GLC	F	3	3	-	0/2/19/22	0/1/1/1
3	GLC	F	4	3	-	0/2/19/22	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	1[A]	BDP	O5-C5	-2.13	1.41	1.43
2	E	5	BDP	O5-C5	-2.11	1.41	1.43
2	C	1	BDP	O5-C5	-2.11	1.41	1.43

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	6	GNS	O3S-S1-O2S	-5.39	107.42	120.16
2	E	4[B]	GNS	O3S-S1-O2S	-4.50	109.53	120.16
2	E	2[A]	GNS	O3S-S1-O2S	-4.34	109.89	120.16
3	D	4	GLC	C1-O5-C5	4.31	118.03	112.19
2	C	6	GNS	O3S-S1-O2S	-4.30	110.00	120.16
2	E	4[A]	GNS	O3S-S1-O2S	-4.20	110.22	120.16
2	C	2	GNS	O3S-S1-O2S	-4.03	110.64	120.16
2	E	5	BDP	C1-C2-C3	3.65	114.15	109.67
2	C	4	GNS	O3S-S1-O2S	-3.18	112.65	120.16
2	E	6	GNS	C1-O5-C5	2.88	116.09	112.19
2	E	4[A]	GNS	O5-C1-C2	-2.85	106.79	111.29
2	E	4[A]	GNS	C3-C4-C5	-2.74	105.34	110.24
2	C	2	GNS	O5-C1-C2	-2.64	107.12	111.29
2	E	1[A]	BDP	C1-C2-C3	2.57	112.83	109.67
3	F	4	GLC	C1-O5-C5	2.57	115.67	112.19
2	C	6	GNS	C1-O5-C5	2.43	115.49	112.19
3	D	3	GLC	O5-C1-C2	-2.34	107.16	110.77
2	E	3[A]	IDS	C1-C2-C3	2.32	112.87	109.40
2	E	4[A]	GNS	O3-C3-C2	2.31	114.25	109.47
2	E	2[A]	GNS	O5-C1-C2	-2.31	107.65	111.29
2	C	1	BDP	C1-C2-C3	2.28	112.47	109.67
2	E	6	GNS	C4-C3-C2	-2.16	107.86	111.02
3	F	4	GLC	O5-C5-C6	2.15	110.58	107.20
2	E	4[A]	GNS	C1-C2-N2	-2.08	106.70	110.27
3	F	3	GLC	O5-C1-C2	-2.06	107.58	110.77
2	C	3	IDS	C1-C2-C3	2.00	112.39	109.40

There are no chirality outliers.


All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	E	4[A]	GNS	C2-N2-S1-O3S
2	E	2[A]	GNS	C2-N2-S1-O2S


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.







5.6 Ligand geometry [i](#)

Of 34 ligands modelled in this entry, 11 are monoatomic - leaving 23 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	EDO	B	1415	-	3,3,3	0.38	0	2,2,2	0.69	0
6	EDO	A	1416	-	3,3,3	0.60	0	2,2,2	0.51	0
5	NPO	B	1408[A]	2	8,8,10	1.14	1 (12%)	10,10,13	0.88	0
6	EDO	A	1414	-	3,3,3	0.46	0	2,2,2	0.35	0
5	NPO	A	1408	2	9,10,10	0.71	0	11,13,13	0.92	0
6	EDO	A	1419	-	3,3,3	0.48	0	2,2,2	0.45	0
6	EDO	A	1417	-	3,3,3	0.41	0	2,2,2	0.66	0
6	EDO	A	1420	-	3,3,3	0.71	0	2,2,2	0.43	0
4	A3P	B	1401	-	26,29,29	0.61	0	31,45,45	0.83	1 (3%)
6	EDO	A	1421	-	3,3,3	0.18	0	2,2,2	0.55	0
6	EDO	A	1415	-	3,3,3	0.41	0	2,2,2	0.25	0
9	TLA	A	1430	8	3,9,9	0.43	0	6,12,12	1.22	1 (16%)
6	EDO	B	1414	-	3,3,3	0.62	0	2,2,2	0.43	0
9	TLA	B	1422	-	3,9,9	0.33	0	6,12,12	1.28	1 (16%)
4	A3P	A	1401	-	26,29,29	0.64	0	31,45,45	0.79	1 (3%)
6	EDO	A	1413	-	3,3,3	0.44	0	2,2,2	0.64	0
6	EDO	B	1416	-	3,3,3	0.65	0	2,2,2	0.54	0
6	EDO	B	1417	-	3,3,3	0.57	0	2,2,2	0.24	0
6	EDO	B	1413	-	3,3,3	0.84	0	2,2,2	0.37	0
6	EDO	A	1418	-	3,3,3	0.50	0	2,2,2	0.54	0
6	EDO	B	1418	-	3,3,3	0.28	0	2,2,2	0.40	0
9	TLA	A	1431	-	3,9,9	0.41	0	6,12,12	1.23	1 (16%)
9	TLA	B	1423	-	3,9,9	0.25	0	6,12,12	1.14	1 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	EDO	B	1415	-	-	0/1/1/1	-
6	EDO	A	1416	-	-	0/1/1/1	-
5	NPO	B	1408[A]	2	-	-	0/1/1/1
6	EDO	A	1414	-	-	0/1/1/1	-
5	NPO	A	1408	2	-	0/2/4/4	0/1/1/1
6	EDO	A	1419	-	-	0/1/1/1	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	EDO	A	1417	-	-	0/1/1/1	-
6	EDO	A	1420	-	-	0/1/1/1	-
4	A3P	B	1401	-	-	0/11/31/31	0/3/3/3
6	EDO	A	1421	-	-	0/1/1/1	-
6	EDO	A	1415	-	-	0/1/1/1	-
9	TLA	A	1430	8	-	0/4/12/12	-
6	EDO	B	1414	-	-	0/1/1/1	-
9	TLA	B	1422	-	-	0/4/12/12	-
4	A3P	A	1401	-	-	0/11/31/31	0/3/3/3
6	EDO	A	1413	-	-	0/1/1/1	-
6	EDO	B	1416	-	-	0/1/1/1	-
6	EDO	B	1417	-	-	1/1/1/1	-
6	EDO	B	1413	-	-	0/1/1/1	-
6	EDO	A	1418	-	-	0/1/1/1	-
6	EDO	B	1418	-	-	1/1/1/1	-
9	TLA	A	1431	-	-	0/4/12/12	-
9	TLA	B	1423	-	-	0/4/12/12	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	1408[A]	NPO	C1-N1	2.39	1.46	1.38

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
9	A	1430	TLA	C1-C2-C3	-2.67	107.36	113.11
9	A	1431	TLA	C1-C2-C3	-2.47	107.79	113.11
9	B	1423	TLA	C4-C3-C2	-2.42	107.91	113.11
4	B	1401	A3P	C5-C6-N6	2.20	123.69	120.35
4	A	1401	A3P	C5-C6-N6	2.17	123.66	120.35
9	B	1422	TLA	C4-C3-C2	-2.12	108.54	113.11

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
6	B	1418	EDO	O1-C1-C2-O2
6	B	1417	EDO	O1-C1-C2-O2


There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	B	1416	EDO	1	0
9	A	1431	TLA	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	670/692 (96%)	0.07	10 (1%) 73 81	17, 31, 53, 83	0
1	B	677/692 (97%)	0.21	27 (3%) 38 48	16, 35, 54, 66	11 (1%)
All	All	1347/1384 (97%)	0.14	37 (2%) 54 63	16, 33, 54, 83	11 (0%)

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1386	TRP	6.3
1	B	1387	ILE	5.1
1	A	52	ALA	4.4
1	B	139	LEU	3.9
1	A	51	ALA	3.8
1	B	173	ALA	3.8
1	A	2	ILE	3.7
1	B	135	LEU	3.4
1	B	186	ALA	3.4
1	A	92	PHE	3.4
1	B	1390	ARG	3.3
1	B	134	ALA	3.2
1	B	33	ILE	3.2
1	B	195	LEU	3.2
1	B	1385	ARG	3.1
1	B	141	ALA	3.0
1	B	4	GLU	2.8
1	B	1377	ARG	2.7
1	A	53	THR	2.6
1	B	1088	ASP	2.6
1	B	146	ALA	2.6
1	A	1170	VAL	2.5
1	A	1171	ILE	2.4
1	B	160	LEU	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	351	ALA	2.4
1	B	172	ALA	2.3
1	A	33	ILE	2.3
1	B	51	ALA	2.3
1	B	353	GLY	2.2
1	B	31	THR	2.2
1	B	92	PHE	2.2
1	A	4	GLU	2.2
1	B	1388	ARG	2.2
1	B	23	VAL	2.2
1	B	1378	LEU	2.1
1	A	1088	ASP	2.1
1	B	169	PHE	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GLC	F	4	11/12	0.82	0.13	39,46,50,51	11
3	GLC	D	4	11/12	0.82	0.18	27,35,40,40	11
2	BDP	E	1[A]	12/13	0.89	0.11	52,55,60,61	12
2	GNS	E	2[A]	15/16	0.90	0.10	45,50,63,68	15
2	BDP	C	1	12/13	0.93	0.10	41,43,47,49	0
3	GLC	F	3	11/12	0.94	0.09	32,35,37,41	0
2	GNS	E	4[A]	15/16	0.95	0.14	29,34,42,45	15
2	GNS	C	2	15/16	0.95	0.08	39,44,63,69	0
2	GNS	E	4[B]	15/16	0.95	0.14	32,35,37,38	15
3	GLC	F	1	12/12	0.95	0.08	28,31,35,37	0
2	BDP	C	5	12/13	0.96	0.11	23,25,30,31	0
3	GLC	F	2	11/12	0.96	0.07	27,28,31,32	0
2	BDP	E	5	12/13	0.96	0.12	23,24,29,29	0
2	IDS	E	3[A]	16/17	0.96	0.08	37,40,46,47	16
2	IDS	C	3	16/17	0.97	0.07	32,39,44,45	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	GLC	D	3	11/12	0.97	0.10	24,27,30,30	0
3	GLC	D	2	11/12	0.98	0.15	19,23,24,24	0
2	GNS	C	4	15/16	0.98	0.07	28,31,38,38	0
2	GNS	E	6	15/16	0.98	0.11	20,22,26,30	0
2	GNS	C	6	15/16	0.98	0.10	22,24,27,28	0
3	GLC	D	1	12/12	0.99	0.12	21,24,26,27	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around Chain E:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain D:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	EDO	A	1421	4/4	0.73	0.35	34,38,42,42	4
6	EDO	B	1417	4/4	0.77	0.16	54,57,59,64	0
8	NA	A	1427	1/1	0.79	0.23	36,36,36,36	1
6	EDO	A	1420	4/4	0.82	0.16	29,30,39,45	0
6	EDO	B	1416	4/4	0.85	0.15	32,37,40,45	4
5	NPO	B	1408[A]	8/10	0.85	0.13	48,53,55,56	8
6	EDO	B	1418	4/4	0.85	0.17	27,32,33,37	4
8	NA	B	1421	1/1	0.85	0.16	53,53,53,53	1
9	TLA	B	1423	10/10	0.85	0.25	26,32,40,40	10
8	NA	A	1428	1/1	0.87	0.17	35,35,35,35	1
6	EDO	A	1419	4/4	0.88	0.22	32,33,37,39	0
6	EDO	A	1413	4/4	0.88	0.23	38,41,45,48	4
8	NA	A	1426	1/1	0.89	0.14	50,50,50,50	0
8	NA	A	1429	1/1	0.90	0.22	43,43,43,43	1

Continued on next page...




Continued from previous page...


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
9	TLA	A	1430	10/10	0.90	0.14	26,33,38,39	10
8	NA	A	1425	1/1	0.91	0.08	47,47,47,47	0
6	EDO	A	1416	4/4	0.91	0.13	26,28,32,38	0
6	EDO	B	1413	4/4	0.92	0.12	25,27,28,33	0
5	NPO	A	1408	10/10	0.92	0.18	35,43,49,52	0
9	TLA	A	1431	10/10	0.93	0.27	39,47,48,53	10
6	EDO	A	1418	4/4	0.93	0.12	33,33,36,43	0
8	NA	B	1420	1/1	0.93	0.18	44,44,44,44	0
6	EDO	B	1414	4/4	0.94	0.16	33,33,34,41	0
9	TLA	B	1422	10/10	0.95	0.12	25,31,35,39	10
6	EDO	A	1414	4/4	0.96	0.09	35,36,36,39	0
7	CL	B	1419	1/1	0.96	0.07	43,43,43,43	0
6	EDO	A	1415	4/4	0.97	0.12	32,36,38,40	0
7	CL	A	1424	1/1	0.98	0.16	47,47,47,47	0
4	A3P	A	1401	27/27	0.98	0.12	19,22,25,26	0
6	EDO	B	1415	4/4	0.98	0.12	23,24,24,28	0
6	EDO	A	1417	4/4	0.98	0.09	31,32,33,35	0
4	A3P	B	1401	27/27	0.99	0.12	18,20,23,24	0
7	CL	A	1423	1/1	0.99	0.06	41,41,41,41	0
7	CL	A	1422	1/1	1.00	0.06	31,31,31,31	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around A3P A 1401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.