



Full wwPDB X-ray Structure Validation Report ⓘ

May 13, 2020 – 03:21 am BST

PDB ID : 3T5Y
Title : Crystal structure of CerJ from *Streptomyces tendae* - malonic acid covalently linked to the catalytic Cystein C116
Authors : Zocher, G.; Bretschneider, T.; Hertweck, C.; Stehle, T.
Deposited on : 2011-07-28
Resolution : 2.12 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.12 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6241 (2.14-2.10)
Clashscore	141614	6778 (2.14-2.10)
Ramachandran outliers	138981	6705 (2.14-2.10)
Sidechain outliers	138945	6706 (2.14-2.10)
RSRZ outliers	127900	6112 (2.14-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	357	 83% 11% . .
2	B	357	 83% 10% . .

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 5362 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called CerJ.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	342	2538	1592	457	481	8	0	3	0

- Molecule 2 is a protein called CerJ.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	342	2519	1577	455	479	8	0	0	0

- Molecule 3 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	1	4	2	0	0
3	A	1	4	2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			4	2	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	181	Total	O	0	0
			181	181		
4	B	112	Total	O	0	0
			112	112		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: CerJ



- Molecule 2: CerJ



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	106.04Å 106.04Å 121.69Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.41 – 2.12 29.41 – 2.12	Depositor EDS
% Data completeness (in resolution range)	100.0 (29.41-2.12) 99.8 (29.41-2.12)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.14 (at 2.12Å)	Xtrriage
Refinement program	REFMAC - Rigid Body	Depositor
R, R_{free}	0.168 , 0.208 0.176 , 0.216	Depositor DCC
R_{free} test set	1994 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	31.7	Xtrriage
Anisotropy	0.532	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 47.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.28$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5362	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.54% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MCS, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.72	0/2604	0.81	5/3556 (0.1%)
2	B	0.64	0/2561	0.79	2/3496 (0.1%)
All	All	0.68	0/5165	0.80	7/7052 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	5
2	B	0	6
All	All	0	11

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	291	ARG	NE-CZ-NH2	-6.61	117.00	120.30
1	A	209	GLU	C-N-CD	-6.04	107.31	120.60
2	B	329	TYR	N-CA-CB	5.94	121.30	110.60
1	A	267	ARG	NE-CZ-NH1	5.59	123.10	120.30
1	A	329	TYR	N-CA-CB	5.22	120.00	110.60
2	B	198	ARG	NE-CZ-NH1	5.19	122.89	120.30
1	A	291	ARG	NE-CZ-NH1	5.16	122.88	120.30

There are no chirality outliers.

All (11) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	150	GLY	Mainchain,Peptide
1	A	209	GLU	Peptide
1	A	328	GLY	Mainchain,Peptide
2	B	116	MCS	Mainchain
2	B	150	GLY	Mainchain,Peptide
2	B	209	GLU	Peptide
2	B	328	GLY	Mainchain,Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2538	0	2490	24	0
2	B	2519	0	2466	18	0
3	A	8	0	6	2	0
3	B	4	0	3	0	0
4	A	181	0	0	2	0
4	B	112	0	0	0	0
All	All	5362	0	4965	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:114:GLN:NE2	1:A:117:ASN:HD22	1.70	0.90
2:B:43:VAL:HG22	2:B:294:GLY:HA3	1.70	0.72
1:A:114:GLN:HE21	1:A:117:ASN:HD22	1.41	0.69
2:B:229:LEU:HD12	2:B:232:LEU:CD1	2.24	0.67
1:A:277:LEU:O	1:A:277:LEU:HD23	1.94	0.66
1:A:308:LEU:HD13	4:A:477:HOH:O	2.00	0.61
2:B:229:LEU:HD12	2:B:232:LEU:HD13	1.81	0.61
1:A:116:CYS:SG	3:A:358:ACT:H2	2.44	0.58
1:A:4:GLU:N	3:A:359:ACT:O	2.39	0.55
1:A:89:ILE:HD13	2:B:198:ARG:HB2	1.89	0.55
1:A:277:LEU:C	1:A:277:LEU:HD23	2.27	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:244:LYS:HG2	2:B:248:LEU:HD22	1.89	0.53
1:A:114:GLN:NE2	1:A:117:ASN:ND2	2.51	0.50
1:A:241:VAL:CG2	1:A:277:LEU:HD21	2.41	0.50
1:A:92:TRP:CD1	1:A:93:PRO:HD2	2.47	0.50
2:B:239:LEU:C	2:B:239:LEU:HD12	2.33	0.49
1:A:260[A]:HIS:ND1	1:A:308:LEU:HD21	2.28	0.48
1:A:196:LEU:HD11	1:A:232:LEU:HD11	1.94	0.48
4:A:463:HOH:O	2:B:210:PRO:HD3	2.12	0.48
2:B:260:HIS:CD2	2:B:308:LEU:HD11	2.49	0.48
2:B:277:LEU:C	2:B:277:LEU:HD23	2.36	0.46
2:B:265:VAL:HG22	2:B:290:GLY:HA3	1.99	0.45
2:B:229:LEU:HD12	2:B:232:LEU:HD12	1.97	0.45
1:A:58:ALA:HA	1:A:168:ALA:HB1	1.99	0.44
1:A:241:VAL:HG22	1:A:277:LEU:HD21	2.00	0.44
1:A:277:LEU:C	1:A:277:LEU:CD2	2.86	0.44
2:B:84:LEU:HD22	2:B:297:GLY:HA2	1.99	0.44
2:B:21:ALA:O	2:B:25:VAL:HG23	2.18	0.43
1:A:265:VAL:HG22	1:A:290:GLY:HA3	1.99	0.43
2:B:115:ARG:HB3	2:B:116:MCS:HAG2	2.01	0.43
1:A:16:PRO:HB2	1:A:17:PRO:CD	2.50	0.42
2:B:195:ILE:HG21	2:B:227:ALA:HB2	2.02	0.41
1:A:196:LEU:HD12	1:A:236[B]:TYR:OH	2.20	0.41
1:A:161:VAL:HG11	1:A:268:ARG:HH21	1.85	0.41
1:A:148:PHE:HB3	1:A:153:ILE:HG21	2.02	0.41
1:A:139:ALA:HA	1:A:173:LEU:O	2.21	0.41
1:A:26:MET:HA	1:A:26:MET:CE	2.50	0.41
2:B:148:PHE:HB3	2:B:153:ILE:HG21	2.03	0.41
1:A:245:THR:O	1:A:249:GLU:HG3	2.21	0.40
2:B:316:PRO:HA	2:B:338:ILE:HG12	2.03	0.40
2:B:92:TRP:CD1	2:B:93:PRO:HD2	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	343/357 (96%)	330 (96%)	10 (3%)	3 (1%)	17	12
2	B	339/357 (95%)	327 (96%)	10 (3%)	2 (1%)	25	20
All	All	682/714 (96%)	657 (96%)	20 (3%)	5 (1%)	22	17

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	210	PRO
2	B	210	PRO
1	A	115	ARG
1	A	151	PRO
2	B	151	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	252/262 (96%)	242 (96%)	10 (4%)	31	31
2	B	247/261 (95%)	232 (94%)	15 (6%)	18	15
All	All	499/523 (95%)	474 (95%)	25 (5%)	24	22

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	68	ARG
1	A	70	ASP
1	A	125	LEU
1	A	142	LEU
1	A	221	VAL
1	A	244	LYS
1	A	268	ARG
1	A	280	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	291	ARG
1	A	308	LEU
2	B	43	VAL
2	B	125	LEU
2	B	141	LEU
2	B	142	LEU
2	B	171	LEU
2	B	176	ARG
2	B	181	ARG
2	B	233	THR
2	B	237	ILE
2	B	239	LEU
2	B	248	LEU
2	B	268	ARG
2	B	280	LEU
2	B	338	ILE
2	B	341	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	5	ASN
1	A	114	GLN
1	A	301	GLN
1	A	307	HIS
2	B	260	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	MCS	B	116	2	6,11,12	1.43	1 (16%)	4,13,15	5.85	4 (100%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MCS	B	116	2	-	3/6/11/13	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	116	MCS	CAK-SAI	2.51	1.82	1.76

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	116	MCS	CB-SAI-CAK	9.21	113.71	100.84
2	B	116	MCS	CAG-CAK-SAI	6.31	121.55	113.69
2	B	116	MCS	OAC-CAK-CAG	-2.82	118.47	123.35
2	B	116	MCS	OAC-CAK-SAI	-2.17	119.80	122.61

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	116	MCS	OAC-CAK-SAI-CB
2	B	116	MCS	CAG-CAK-SAI-CB
2	B	116	MCS	CAJ-CAG-CAK-OAC

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	116	MCS	1	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	ACT	B	358	-	1,3,3	1.63	0	0,3,3	0.00	-
3	ACT	A	358	-	1,3,3	2.28	1 (100%)	0,3,3	0.00	-
3	ACT	A	359	-	1,3,3	1.86	0	0,3,3	0.00	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	358	ACT	CH3-C	2.28	1.51	1.48

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	358	ACT	1	0
3	A	359	ACT	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	342/357 (95%)	-0.07	3 (0%) 84 86	34, 44, 62, 78	0
2	B	341/357 (95%)	0.25	18 (5%) 26 31	33, 52, 76, 88	1 (0%)
All	All	683/714 (95%)	0.09	21 (3%) 49 55	33, 47, 71, 88	1 (0%)

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	342	PRO	4.1
1	A	70	ASP	3.8
2	B	230	PRO	3.7
2	B	269	GLY	3.6
2	B	70	ASP	3.6
2	B	313	LEU	3.3
2	B	232	LEU	3.0
2	B	312	ALA	2.7
2	B	234	HIS	2.6
2	B	221	VAL	2.6
2	B	71	VAL	2.5
2	B	229	LEU	2.5
1	A	230	PRO	2.4
2	B	268	ARG	2.3
2	B	37	LEU	2.3
2	B	226	LEU	2.2
2	B	270	THR	2.2
2	B	112	LEU	2.2
2	B	73	PRO	2.2
1	A	236[A]	TYR	2.1
2	B	74	ALA	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	MCS	B	116	12/13	0.93	0.17	41,48,60,61	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	ACT	B	358	4/4	0.86	0.27	42,45,45,46	4
3	ACT	A	359	4/4	0.91	0.25	34,37,40,45	4
3	ACT	A	358	4/4	0.97	0.12	48,49,49,52	0

6.5 Other polymers [i](#)

There are no such residues in this entry.