

Full wwPDB X-ray Structure Validation Report (i)

May 15, 2020 – 02:26 am BST

PDB ID : 6T73

Title: New antiparallel dimer of aureochrome 1a LOV domain mutants from Phaeo-

dactylum tricornutum

Authors : Essen, L.O.; Hepp, S.

Deposited on : 2019-10-21

Resolution : 3.44 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul: 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.11

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac: 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.11

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 3.44 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \text{Whole archive} \\ (\#\text{Entries}) \end{array}$	$\begin{array}{c} {\rm Similar \; resolution} \\ (\#{\rm Entries, \; resolution \; range(\AA)}) \end{array}$
R_{free}	130704	1278 (3.50-3.38)
Clashscore	141614	1361 (3.50-3.38)
Ramachandran outliers	138981	1327 (3.50-3.38)
Sidechain outliers	138945	1328 (3.50-3.38)
RSRZ outliers	127900	1192 (3.50-3.38)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	162	70%	14%	16%
1	В	162	70%	14%	16%
1	С	162	70%	14%	16%



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 3232 atoms, of which 0 are hydrogens and 0 are deuteriums.


In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Ptaureo1a lov2 domain.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	Λ	136	Total	С	N	О	S	0	0	0
1	Α	150	1052	659	180	207	6	0	0	
1	D	136	Total	С	N	О	S	0	0	0
1	Б	150	1048	657	180	205	6	0	0	
1	С	136	Total	С	N	О	S	0	0	0
1		130	1036	651	179	200	6	0	0	U

There are 63 discrepancies between the modelled and reference sequences:

A	21 =		Actual	${f Comment}$	${f Reference}$
	217	MET	-	initiating methionine	UNP A0A140UHJ0
A	218	GLY	-	expression tag	UNP A0A140UHJ0
A	219	SER	-	expression tag	UNP A0A140UHJ0
A	220	SER	-	expression tag	UNP A0A140UHJ0
A	221	HIS	_	expression tag	UNP A0A140UHJ0
A	222	HIS	-	expression tag	UNP A0A140UHJ0
A	223	HIS	-	expression tag	UNP A0A140UHJ0
A	224	HIS	-	expression tag	UNP A0A140UHJ0
A	225	HIS	-	expression tag	UNP A0A140UHJ0
A	226	HIS	-	expression tag	UNP A0A140UHJ0
A	227	SER	-	expression tag	UNP A0A140UHJ0
A	228	SER	_	expression tag	UNP A0A140UHJ0
A	229	GLY	-	expression tag	UNP A0A140UHJ0
A	230	LEU	-	expression tag	UNP A0A140UHJ0
A	231	VAL	-	expression tag	UNP A0A140UHJ0
A	232	PRO	-	expression tag	UNP A0A140UHJ0
A	233	ARG	-	expression tag	UNP A0A140UHJ0
A	234	GLY	=	expression tag	UNP A0A140UHJ0
A	235	SER	-	expression tag	UNP A0A140UHJ0
A	236	HIS	-	expression tag	UNP A0A140UHJ0
A	254	MET	VAL	conflict	UNP A0A140UHJ0
В	217	MET	-	initiating methionine	UNP A0A140UHJ0
В	218	GLY	-	expression tag	UNP A0A140UHJ0






Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
В	219	SER	_	expression tag	UNP A0A140UHJ0
В	220	SER	-	expression tag	UNP A0A140UHJ0
В	221	HIS	-	expression tag	UNP A0A140UHJ0
В	222	HIS	_	expression tag	UNP A0A140UHJ0
В	223	HIS	-	expression tag	UNP A0A140UHJ0
В	224	HIS	-	expression tag	UNP A0A140UHJ0
В	225	HIS	-	expression tag	UNP A0A140UHJ0
В	226	HIS	-	expression tag	UNP A0A140UHJ0
В	227	SER	=	expression tag	UNP A0A140UHJ0
В	228	SER	-	expression tag	UNP A0A140UHJ0
В	229	GLY	=	expression tag	UNP A0A140UHJ0
В	230	LEU	=	expression tag	UNP A0A140UHJ0
В	231	VAL	=	expression tag	UNP A0A140UHJ0
В	232	PRO	-	expression tag	UNP A0A140UHJ0
В	233	ARG	-	expression tag	UNP A0A140UHJ0
В	234	GLY	-	expression tag	UNP A0A140UHJ0
В	235	SER	_	expression tag	UNP A0A140UHJ0
В	236	HIS	=	expression tag	UNP A0A140UHJ0
В	254	MET	VAL	conflict	UNP A0A140UHJ0
С	217	MET	_	initiating methionine	UNP A0A140UHJ0
С	218	GLY	-	expression tag	UNP A0A140UHJ0
С	219	SER	_	expression tag	UNP A0A140UHJ0
С	220	SER	ı	expression tag	UNP A0A140UHJ0
С	221	HIS	-	expression tag	UNP A0A140UHJ0
С	222	HIS	ı	expression tag	UNP A0A140UHJ0
С	223	HIS	-	expression tag	UNP A0A140UHJ0
С	224	HIS	I	expression tag	UNP A0A140UHJ0
С	225	HIS	-	expression tag	UNP A0A140UHJ0
С	226	HIS	-	expression tag	UNP A0A140UHJ0
С	227	SER	-	expression tag	UNP A0A140UHJ0
С	228	SER	-	expression tag	UNP A0A140UHJ0
С	229	GLY	-	expression tag	UNP A0A140UHJ0
С	230	LEU	-	expression tag	UNP A0A140UHJ0
С	231	VAL	-	expression tag	UNP A0A140UHJ0
С	232	PRO	-	expression tag	UNP A0A140UHJ0
С	233	ARG	-	expression tag	UNP A0A140UHJ0
С	234	GLY	-	expression tag	UNP A0A140UHJ0
С	235	SER	-	expression tag	UNP A0A140UHJ0
С	236	HIS	-	expression tag	UNP A0A140UHJ0
С	254	MET	VAL	conflict	UNP A0A140UHJ0

 $\bullet \ \ Molecule\ 2\ is\ FLAVIN\ MONONUCLEOTIDE\ (three-letter\ code:\ FMN)\ (formula:\ C_{17}H_{21}N_4O_9P).$





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf		
2	Λ	1	Total	С	N	О	Р	0	0
	Α	1	31	17	4	9	1	U	0
2	D	1	Total	С	N	О	Р	0	0
	Ъ	1	31	17	4	9	1	U	0
2	С	1	Total	С	N	О	Р	0	0
		1	31	17	4	9	1	U	0

 \bullet Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

\mathbf{Mol}	Chain	Residues	${f Atoms}$	ZeroOcc	AltConf
3	В	3	Total Cl 3 3	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Ptaureo1a lov2 domain Chain A: 14% 16% ALA ALA ALA ASN ASP ASP ASP ASP ASP • Molecule 1: Ptaureo1a lov2 domain Chain B: 70% 14% 16% ALA ALA ASN ASP ASP GLU ASP • Molecule 1: Ptaureo1a lov2 domain Chain C: 70% 14% 16%



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants	168.37Å 168.37Å 95.71Å	Danagitan
a, b, c, α , β , γ	90.00° 90.00° 120.00°	Depositor
Resolution (Å)	48.60 - 3.44	Depositor
Resolution (A)	48.61 - 3.43	EDS
% Data completeness	98.8 (48.60-3.44)	Depositor
(in resolution range)	98.8 (48.61-3.43)	EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	0.92 (at 3.40Å)	Xtriage
Refinement program	PHENIX 1.16_3549	Depositor
D D	0.249 , 0.278	Depositor
R, R_{free}	0.249 , 0.278	DCC
R_{free} test set	578 reflections $(5.26%)$	wwPDB-VP
Wilson B-factor (Å ²)	107.3	Xtriage
Anisotropy	0.517	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.34 , 73.2	EDS
L-test for twinning ²	$ < L >=0.46, < L^2>=0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	3232	wwPDB-VP
Average B, all atoms $(Å^2)$	115.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 5.02% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $< L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: FMN, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Chain		Bond	lengths	Bond angles		
IVIOI	Chain	RMSZ	# Z >5	RMSZ	# Z > 5	
1	A	0.26	0/1072	0.44	0/1457	
1	В	0.26	0/1068	0.44	0/1452	
1	С	0.26	0/1056	0.44	0/1437	
All	All	0.26	0/3196	0.44	0/4346	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1052	0	987	21	0
1	В	1048	0	983	17	0
1	С	1036	0	969	15	1
2	A	31	0	19	1	0
2	В	31	0	19	1	0
2	С	31	0	19	1	0
3	В	3	0	0	0	0
All	All	3232	0	2996	44	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.



All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

A 4 1	A 4 0	Interatomic	Clash
Atom-1	Atom-2	${f distance}({f \AA})$	$-$ overlap (\AA)
1:B:318:LEU:HD11	1:B:326:THR:HB	1.70	0.74
1:A:318:LEU:HD11	1:A:326:THR:HB	1.70	0.74
1:C:318:LEU:HD11	1:C:326:THR:HB	1.70	0.72
1:A:238:ASP:OD2	1:B:302:ARG:NH1	2.15	0.71
1:A:262:ASN:HD21	1:A:346:PHE:HD2	1.41	0.68
1:C:262:ASN:HD21	1:C:346:PHE:HD2	1.41	0.67
1:B:262:ASN:HD21	1:B:346:PHE:HD2	1.41	0.67
1:A:261:ASP:OD1	1:A:288:ARG:NH2	2.30	0.64
1:C:261:ASP:OD1	1:C:288:ARG:NH2	2.30	0.63
1:A:364:GLN:OE1	1:B:243:LYS:NZ	2.28	0.62
1:B:261:ASP:OD1	1:B:288:ARG:NH2	2.30	0.62
1:A:243:LYS:NZ	1:B:364:GLN:OE1	2.27	0.61
1:A:247:THR:HG21	1:C:280:ASP:HB3	1.86	0.58
1:A:288:ARG:NH1	2:A:500:FMN:O3P	2.37	0.58
1:C:288:ARG:NH1	2:C:500:FMN:O3P	2.37	0.57
1:B:288:ARG:NH1	2:B:401:FMN:O3P	2.37	0.55
1:A:247:THR:CG2	1:C:280:ASP:HB3	2.37	0.55
1:C:248:ALA:O	1:C:268:SER:OG	2.24	0.53
1:A:301:GLU:HA	1:A:304:ARG:HG2	1.91	0.53
1:B:301:GLU:HA	1:B:304:ARG:HG2	1.91	0.52
1:C:301:GLU:HA	1:C:304:ARG:HG2	1.91	0.52
1:A:248:ALA:O	1:A:268:SER:OG	2.24	0.52
1:C:302:ARG:HG3	1:C:313:MET:HE1	1.93	0.51
1:A:302:ARG:HG3	1:A:313:MET:HE1	1.94	0.50
1:B:250:GLN:HB2	1:B:252:PHE:CE2	2.47	0.50
1:C:250:GLN:HB2	1:C:252:PHE:CE2	2.47	0.50
1:B:302:ARG:HG3	1:B:313:MET:HE1	1.93	0.49
1:B:248:ALA:O	1:B:268:SER:OG	2.24	0.49
1:A:247:THR:CG2	1:B:357:TYR:HE1	2.26	0.49
1:A:250:GLN:HB2	1:A:252:PHE:CE2	2.47	0.49
1:A:280:ASP:HB3	1:C:247:THR:HG21	1.99	0.44
1:A:250:GLN:O	1:A:268:SER:OG	2.36	0.43
1:B:250:GLN:O	1:B:268:SER:OG	2.36	0.42
1:B:358:ALA:O	1:B:362:THR:OG1	2.25	0.42
1:C:250:GLN:O	1:C:268:SER:OG	2.36	0.42
1:C:286:ASN:OD1	1:C:288:ARG:HG3	2.20	0.41
1:A:247:THR:HG21	1:B:357:TYR:HE1	1.85	0.41
1:C:303:ILE:HA	1:C:313:MET:SD	2.61	0.41
1:A:303:ILE:HA	1:A:313:MET:SD	2.61	0.41
1:B:286:ASN:OD1	1:B:288:ARG:HG3	2.20	0.41

Continued on next page...



Continued from previous page...

Atom-1	Atom-2	$egin{array}{c} ext{Interatomic} \ ext{distance} \ (ext{Å}) \end{array}$	$egin{aligned} ext{Clash} \ ext{overlap} \ (ext{\AA}) \end{aligned}$
1:B:303:ILE:HA	1:B:313:MET:SD	2.61	0.41
1:A:286:ASN:OD1	1:A:288:ARG:HG3	2.20	0.40
1:A:243:LYS:HG3	1:C:283:LEU:HB2	2.03	0.40
1:A:358:ALA:O	1:A:362:THR:OG1	2.25	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	$egin{aligned} ext{Interatomic} \ ext{distance} & (ext{Å}) \end{aligned}$	$egin{aligned} ext{Clash} \ ext{overlap } (ext{Å}) \end{aligned}$
1:C:243:LYS:NZ	1:C:364:GLN:OE1[11_555]	2.19	0.01

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	A	134/162 (83%)	129 (96%)	5 (4%)	0	100	100
1	В	134/162 (83%)	129 (96%)	4 (3%)	1 (1%)	22	60
1	С	134/162 (83%)	129 (96%)	4 (3%)	1 (1%)	22	60
All	All	402/486 (83%)	387 (96%)	13 (3%)	2 (0%)	29	66

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type	
1	В	293	PRO	
1	С	293	PRO	

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar



resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	A	112/137 (82%)	112 (100%)	0	100	100
1	В	111/137 (81%)	111 (100%)	0	100	100
1	С	108/137 (79%)	108 (100%)	0	100	100
All	All	331/411 (80%)	331 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

Of 6 ligands modelled in this entry, 3 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Tuno	Chain	Pos	Link	Bo	ond leng	ths	В	ond ang	les
MIGI	Type	Chain	res	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	FMN	В	401	-	31,33,33	1.40	4 (12%)	40,50,50	1.46	5 (12%)



Mol	T	Chain	Dog	Link	Bond lengths			Bond angles		
MIOI	Type	Chain	Res	Link	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	FMN	С	500	-	31,33,33	1.40	4 (12%)	40,50,50	1.47	5 (12%)
2	FMN	A	500	-	31,33,33	1.39	4 (12%)	40,50,50	1.46	5 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FMN	В	401	_	-	1/18/18/18	0/3/3/3
2	FMN	С	500	_	-	1/18/18/18	0/3/3/3
2	FMN	A	500	-	-	1/18/18/18	0/3/3/3

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\operatorname{\AA})$	$\operatorname{Ideal}(\text{\AA})$
2	A	500	FMN	C10-N1	4.06	1.38	1.33
2	В	401	FMN	C10-N1	4.06	1.38	1.33
2	С	500	FMN	C10-N1	4.01	1.38	1.33
2	С	500	FMN	C4A-N5	3.59	1.38	1.33
2	В	401	FMN	C4A-N5	3.59	1.38	1.33
2	A	500	FMN	C4A-N5	3.56	1.38	1.33
2	В	401	FMN	C4-N3	3.11	1.38	1.33
2	С	500	FMN	C4-N3	3.06	1.38	1.33
2	A	500	FMN	C4-N3	3.04	1.38	1.33
2	A	500	FMN	C1'-N10	2.14	1.50	1.48
2	С	500	FMN	C1'-N10	2.13	1.50	1.48
2	В	401	FMN	C1'-N10	2.08	1.50	1.48

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^o)$
2	С	500	FMN	C4-N3-C2	5.87	120.10	115.14
2	A	500	FMN	C4-N3-C2	5.78	120.02	115.14
2	В	401	FMN	C4-N3-C2	5.75	120.00	115.14
2	В	401	FMN	C4A-N5-C5A	3.24	120.01	116.77
2	A	500	FMN	C4A-N5-C5A	3.23	120.00	116.77
2	С	500	FMN	C4A-N5-C5A	3.20	119.97	116.77
2	С	500	FMN	C5A-C9A-N10	3.08	119.95	117.72
2	A	500	FMN	C5A-C9A-N10	3.06	119.93	117.72
2	В	401	FMN	C5A-C9A-N10	3.04	119.92	117.72
2	С	500	FMN	C4A-C4-N3	-2.57	119.91	123.43

 $Continued\ on\ next\ page...$



$\alpha \cdots \tau$	e	•	
Continued	trom	mraniaone	maaa
-	110116	predidus	puyc

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^o)$
2	В	401	FMN	C4A-C4-N3	-2.55	119.94	123.43
2	A	500	FMN	C4A-C4-N3	-2.53	119.97	123.43
2	С	500	FMN	C1'-N10-C9A	2.18	120.01	118.29
2	В	401	FMN	C1'-N10-C9A	2.18	120.01	118.29
2	A	500	FMN	C1'-N10-C9A	2.13	119.97	118.29

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	С	500	FMN	C4'-C5'-O5'-P
2	A	500	FMN	C4'-C5'-O5'-P
2	В	401	FMN	C4'-C5'-O5'-P


There are no ring outliers.

3 monomers are involved in 3 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	В	401	FMN	1	0
2	С	500	FMN	1	0
2	A	500	FMN	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.














5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\#\mathrm{RSRZ}{>}2$		>2	$\mathbf{OWAB}(\mathrm{\AA}^2)$	Q < 0.9
1	A	$136/162 \ (83\%)$	0.41	4 (2%)	51	50	104, 115, 138, 151	0
1	В	136/162 (83%)	0.26	1 (0%)	87	85	93, 108, 126, 131	0
1	С	136/162 (83%)	0.39	6 (4%)	34	33	106, 116, 139, 153	0
All	All	408/486 (83%)	0.35	11 (2%)	54	52	93, 114, 138, 153	0

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	339	ALA	4.0
1	С	328	TRP	2.7
1	С	346	PHE	2.5
1	С	318	LEU	2.4
1	A	346	PHE	2.3
1	A	318	LEU	2.2
1	A	333	ILE	2.1
1	С	316	CYS	2.1
1	С	350	GLN	2.1
1	В	346	PHE	2.1
1	С	235	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.




6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	${f B\text{-factors}}({f \AA}^2)$	Q < 0.9
3	CL	В	402	1/1	0.69	0.33	102,102,102,102	0
3	CL	В	404	1/1	0.81	0.37	96,96,96,96	0
2	FMN	A	500	31/31	0.89	0.28	112,112,112,112	0
2	FMN	С	500	31/31	0.92	0.26	111,111,111,111	0
2	FMN	В	401	31/31	0.93	0.29	100,100,100,100	0
3	CL	В	403	1/1	0.94	0.17	103,103,103,103	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.














6.5 Other polymers (i)

There are no such residues in this entry.

