



wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 24, 2024 – 09:22 AM EDT

PDB ID : 6T76
Title : PII-like protein CutA from Nostoc sp. PCC 7120 in apo form
Authors : Selim, K.A.; Albrecht, R.; Forchhammer, K.; Hartmann, M.D.
Deposited on : 2019-10-21
Resolution : 1.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 2.37.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	114	 2% 81% 11% 9%
1	B	114	 % 78% 12% 9%
1	C	114	 3% 82% 8% 10%
1	D	114	 3% 77% 11% 11%
1	E	114	 4% 82% 9% 9%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	114	 <p>A horizontal bar chart representing the quality of chain. The bar is divided into four segments: a small red segment at the beginning labeled '2%', a large green segment labeled '79%', a yellow segment labeled '10%', and a grey segment at the end labeled '11%'.</p>

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 9955 atoms, of which 4917 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Periplasmic divalent cation tolerance protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	104	1659	534	829	136	154	6	21	0	0
1	B	104	1665	535	835	136	153	6	21	0	0
1	C	103	1629	522	815	133	153	6	19	0	0
1	D	102	1624	518	814	136	150	6	20	0	0
1	E	104	1652	529	827	135	155	6	19	0	0
1	F	101	1589	509	797	130	147	6	22	0	0

There are 60 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	105	SER	-	expression tag	UNP Q8YL42
A	106	ALA	-	expression tag	UNP Q8YL42
A	107	TRP	-	expression tag	UNP Q8YL42
A	108	SER	-	expression tag	UNP Q8YL42
A	109	HIS	-	expression tag	UNP Q8YL42
A	110	PRO	-	expression tag	UNP Q8YL42
A	111	GLN	-	expression tag	UNP Q8YL42
A	112	PHE	-	expression tag	UNP Q8YL42
A	113	GLU	-	expression tag	UNP Q8YL42
A	114	LYS	-	expression tag	UNP Q8YL42
B	105	SER	-	expression tag	UNP Q8YL42
B	106	ALA	-	expression tag	UNP Q8YL42
B	107	TRP	-	expression tag	UNP Q8YL42
B	108	SER	-	expression tag	UNP Q8YL42
B	109	HIS	-	expression tag	UNP Q8YL42
B	110	PRO	-	expression tag	UNP Q8YL42
B	111	GLN	-	expression tag	UNP Q8YL42

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	112	PHE	-	expression tag	UNP Q8YL42
B	113	GLU	-	expression tag	UNP Q8YL42
B	114	LYS	-	expression tag	UNP Q8YL42
C	105	SER	-	expression tag	UNP Q8YL42
C	106	ALA	-	expression tag	UNP Q8YL42
C	107	TRP	-	expression tag	UNP Q8YL42
C	108	SER	-	expression tag	UNP Q8YL42
C	109	HIS	-	expression tag	UNP Q8YL42
C	110	PRO	-	expression tag	UNP Q8YL42
C	111	GLN	-	expression tag	UNP Q8YL42
C	112	PHE	-	expression tag	UNP Q8YL42
C	113	GLU	-	expression tag	UNP Q8YL42
C	114	LYS	-	expression tag	UNP Q8YL42
D	105	SER	-	expression tag	UNP Q8YL42
D	106	ALA	-	expression tag	UNP Q8YL42
D	107	TRP	-	expression tag	UNP Q8YL42
D	108	SER	-	expression tag	UNP Q8YL42
D	109	HIS	-	expression tag	UNP Q8YL42
D	110	PRO	-	expression tag	UNP Q8YL42
D	111	GLN	-	expression tag	UNP Q8YL42
D	112	PHE	-	expression tag	UNP Q8YL42
D	113	GLU	-	expression tag	UNP Q8YL42
D	114	LYS	-	expression tag	UNP Q8YL42
E	105	SER	-	expression tag	UNP Q8YL42
E	106	ALA	-	expression tag	UNP Q8YL42
E	107	TRP	-	expression tag	UNP Q8YL42
E	108	SER	-	expression tag	UNP Q8YL42
E	109	HIS	-	expression tag	UNP Q8YL42
E	110	PRO	-	expression tag	UNP Q8YL42
E	111	GLN	-	expression tag	UNP Q8YL42
E	112	PHE	-	expression tag	UNP Q8YL42
E	113	GLU	-	expression tag	UNP Q8YL42
E	114	LYS	-	expression tag	UNP Q8YL42
F	105	SER	-	expression tag	UNP Q8YL42
F	106	ALA	-	expression tag	UNP Q8YL42
F	107	TRP	-	expression tag	UNP Q8YL42
F	108	SER	-	expression tag	UNP Q8YL42
F	109	HIS	-	expression tag	UNP Q8YL42
F	110	PRO	-	expression tag	UNP Q8YL42
F	111	GLN	-	expression tag	UNP Q8YL42
F	112	PHE	-	expression tag	UNP Q8YL42
F	113	GLU	-	expression tag	UNP Q8YL42

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	114	LYS	-	expression tag	UNP Q8YL42

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
2	A	1	5	4	1	0	0
2	A	1	5	4	1	0	0
2	C	1	5	4	1	0	0
2	C	1	5	4	1	0	0
2	D	1	5	4	1	0	0
2	E	1	5	4	1	0	0
2	F	1	5	4	1	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	26	26	26	0	0
3	B	21	21	21	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	C	7	Total O 7 7	0	0
3	D	24	Total O 24 24	0	0
3	E	14	Total O 14 14	0	0
3	F	10	Total O 10 10	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Periplasmic divalent cation tolerance protein





- Molecule 1: Periplasmic divalent cation tolerance protein





- Molecule 1: Periplasmic divalent cation tolerance protein




- Molecule 1: Periplasmic divalent cation tolerance protein



- Molecule 1: Periplasmic divalent cation tolerance protein



- Molecule 1: Periplasmic divalent cation tolerance protein



4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	62.13Å 90.69Å 93.71Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.85 – 1.90 46.85 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.6 (46.85-1.90) 99.6 (46.85-1.90)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.01 (at 1.90Å)	Xtrriage
Refinement program	REFMAC 5.8.0049 2013/06/30	Depositor
R, R_{free}	0.199 , 0.226 0.208 , 0.230	Depositor DCC
R_{free} test set	2121 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	30.9	Xtrriage
Anisotropy	0.507	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.45 , 44.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.008 for -h,l,k	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	9955	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.28% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.85	0/848	0.90	2/1153 (0.2%)
1	B	0.96	3/848 (0.4%)	0.93	3/1152 (0.3%)
1	C	0.84	0/831	0.84	0/1130
1	D	0.88	0/827	0.94	2/1124 (0.2%)
1	E	0.84	0/842	0.87	1/1144 (0.1%)
1	F	0.73	0/808	0.86	1/1099 (0.1%)
All	All	0.85	3/5004 (0.1%)	0.89	9/6802 (0.1%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	11	GLU	CG-CD	6.23	1.61	1.51
1	B	11	GLU	CD-OE1	6.22	1.32	1.25
1	B	11	GLU	CD-OE2	5.87	1.32	1.25

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	91	ARG	NE-CZ-NH1	6.86	123.73	120.30
1	A	91	ARG	NE-CZ-NH2	-6.59	117.01	120.30
1	D	15	ARG	NE-CZ-NH1	5.78	123.19	120.30
1	B	91	ARG	NE-CZ-NH1	5.35	122.97	120.30
1	B	53	MET	CG-SD-CE	5.28	108.64	100.20

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	830	829	816	8	1
1	B	830	835	822	10	0
1	C	814	815	802	8	1
1	D	810	814	801	9	0
1	E	825	827	814	9	2
1	F	792	797	780	8	2
2	A	10	0	0	0	0
2	C	10	0	0	0	0
2	D	5	0	0	0	0
2	E	5	0	0	0	0
2	F	5	0	0	0	0
3	A	26	0	0	0	0
3	B	21	0	0	0	0
3	C	7	0	0	0	0
3	D	24	0	0	0	0
3	E	14	0	0	0	0
3	F	10	0	0	0	0
All	All	5038	4917	4835	36	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 36 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:1:MET:O	1:F:2:LYS:HD2	1.84	0.77
1:A:30:ASN:HB3	1:B:34:VAL:HG21	1.69	0.74
1:C:92:GLU:HG2	1:E:76:LEU:HD13	1.69	0.73
1:B:30:ASN:HB3	1:D:34:VAL:HG21	1.71	0.72
1:E:30:ASN:HB3	1:F:34:VAL:HG21	1.71	0.71

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:18:ARG:HH22	1:C:92:GLU:OE2[2_555]	1.46	0.14
1:E:75:GLU:OE1	1:F:12:HIS:HE1[2_554]	1.47	0.13
1:E:75:GLU:OE1	1:F:12:HIS:CE1[2_554]	2.19	0.01

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	102/114 (90%)	102 (100%)	0	0	100	100
1	B	102/114 (90%)	102 (100%)	0	0	100	100
1	C	101/114 (89%)	101 (100%)	0	0	100	100
1	D	100/114 (88%)	100 (100%)	0	0	100	100
1	E	102/114 (90%)	102 (100%)	0	0	100	100
1	F	99/114 (87%)	99 (100%)	0	0	100	100
All	All	606/684 (89%)	606 (100%)	0	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	91/104 (88%)	91 (100%)	0	100	100
1	B	90/104 (86%)	89 (99%)	1 (1%)	73	73
1	C	89/104 (86%)	89 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	D	89/104 (86%)	87 (98%)	2 (2%)	52	47
1	E	90/104 (86%)	88 (98%)	2 (2%)	52	47
1	F	85/104 (82%)	85 (100%)	0	100	100
All	All	534/624 (86%)	529 (99%)	5 (1%)	78	79

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	76	LEU
1	D	91	ARG
1	D	102	HIS
1	E	59	GLN
1	E	76	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	A	201	-	4,4,4	0.37	0	6,6,6	1.09	1 (16%)
2	SO4	F	200	-	4,4,4	0.43	0	6,6,6	0.77	0
2	SO4	C	200	-	4,4,4	0.64	0	6,6,6	0.54	0
2	SO4	E	200	-	4,4,4	0.67	0	6,6,6	0.55	0
2	SO4	D	200	-	4,4,4	0.59	0	6,6,6	0.87	0
2	SO4	A	202	-	4,4,4	0.60	0	6,6,6	0.44	0
2	SO4	C	201	-	4,4,4	0.51	0	6,6,6	0.43	0

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	201	SO4	O3-S-O1	-2.24	97.86	109.56

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	104/114 (91%)	0.17	2 (1%) 66 69	23, 33, 63, 71	0
1	B	104/114 (91%)	0.16	1 (0%) 82 84	22, 32, 50, 62	0
1	C	103/114 (90%)	0.31	3 (2%) 51 54	24, 39, 66, 82	0
1	D	102/114 (89%)	0.24	3 (2%) 51 54	22, 32, 54, 76	0
1	E	104/114 (91%)	0.34	4 (3%) 40 43	25, 38, 56, 80	0
1	F	101/114 (88%)	0.28	2 (1%) 65 68	26, 41, 60, 71	0
All	All	618/684 (90%)	0.25	15 (2%) 59 62	22, 35, 61, 82	0

The worst 5 of 15 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	42	GLY	5.7
1	F	42	GLY	4.5
1	E	41	LYS	3.9
1	C	103	LEU	3.9
1	A	103	LEU	3.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	SO4	E	200	5/5	0.89	0.19	52,68,71,76	0
2	SO4	F	200	5/5	0.90	0.11	58,59,70,77	0
2	SO4	C	200	5/5	0.91	0.12	59,69,75,80	0
2	SO4	A	201	5/5	0.95	0.10	49,53,62,68	0
2	SO4	A	202	5/5	0.96	0.28	51,53,65,67	0
2	SO4	D	200	5/5	0.96	0.13	44,52,60,65	0
2	SO4	C	201	5/5	0.98	0.11	42,45,46,49	0

6.5 Other polymers [i](#)

There are no such residues in this entry.