



wwPDB EM Validation Summary Report ⓘ

Feb 22, 2024 – 01:59 PM EST

PDB ID : 8T8P
EMDB ID : EMD-41101
Title : 33-mer FliF MS-ring from Salmonella
Authors : Singh, P.K.; Iverson, T.M.
Deposited on : 2023-06-23
Resolution : 3.40 Å (reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev70
MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




























Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	560	
1	AA	560	
1	B	560	
1	BA	560	
1	C	560	
1	CA	560	
1	D	560	
1	DA	560	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	E	560	
1	EA	560	
1	F	560	
1	FA	560	
1	G	560	
1	GA	560	
1	H	560	
1	HA	560	
1	I	560	
1	J	560	
1	K	560	
1	L	560	
1	M	560	
1	N	560	
1	O	560	
1	P	560	
1	Q	560	
1	R	560	
1	S	560	
1	T	560	
1	V	560	
1	W	560	
1	X	560	
1	Y	560	
1	Z	560	

2 Entry composition [i](#)

There is only 1 type of molecule in this entry. The entry contains 64845 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Flagellar M-ring protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	254	1945	1190	358	393	4	0	0
1	B	311	2005	1209	387	406	3	0	0
1	C	254	1945	1190	358	393	4	0	0
1	D	254	1945	1190	358	393	4	0	0
1	E	311	2005	1209	387	406	3	0	0
1	F	254	1945	1190	358	393	4	0	0
1	G	254	1945	1190	358	393	4	0	0
1	H	311	2005	1209	387	406	3	0	0
1	I	254	1945	1190	358	393	4	0	0
1	J	254	1945	1190	358	393	4	0	0
1	K	311	2005	1209	387	406	3	0	0
1	L	254	1945	1190	358	393	4	0	0
1	M	254	1945	1190	358	393	4	0	0
1	N	311	2005	1209	387	406	3	0	0
1	O	254	1945	1190	358	393	4	0	0
1	P	254	1945	1190	358	393	4	0	0
1	Q	311	2005	1209	387	406	3	0	0


Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms					AltConf	Trace
1	R	254	Total 1945	C 1190	N 358	O 393	S 4	0	0
1	S	254	Total 1945	C 1190	N 358	O 393	S 4	0	0
1	T	311	Total 2005	C 1209	N 387	O 406	S 3	0	0
1	V	254	Total 1945	C 1190	N 358	O 393	S 4	0	0
1	W	254	Total 1945	C 1190	N 358	O 393	S 4	0	0
1	X	311	Total 2005	C 1209	N 387	O 406	S 3	0	0
1	Y	254	Total 1945	C 1190	N 358	O 393	S 4	0	0
1	Z	254	Total 1945	C 1190	N 358	O 393	S 4	0	0
1	AA	311	Total 2005	C 1209	N 387	O 406	S 3	0	0
1	BA	254	Total 1945	C 1190	N 358	O 393	S 4	0	0
1	CA	254	Total 1945	C 1190	N 358	O 393	S 4	0	0
1	DA	311	Total 2005	C 1209	N 387	O 406	S 3	0	0
1	EA	254	Total 1945	C 1190	N 358	O 393	S 4	0	0
1	FA	254	Total 1945	C 1190	N 358	O 393	S 4	0	0
1	GA	311	Total 2005	C 1209	N 387	O 406	S 3	0	0
1	HA	254	Total 1945	C 1190	N 358	O 393	S 4	0	0




• Molecule 1: Flagellar M-ring protein







• Molecule 1: Flagellar M-ring protein





• Molecule 1: Flagellar M-ring protein







● Molecule 1: Flagellar M-ring protein





● Molecule 1: Flagellar M-ring protein






• Molecule 1: Flagellar M-ring protein





• Molecule 1: Flagellar M-ring protein






• Molecule 1: Flagellar M-ring protein





ARG
GLU
MET
ALA
THR
SER
ASP
ASN
ASP
PRO
ARG
VAL
VAL
ALA
ALA
VAL
VAL
ILE

• Molecule 1: Flagellar M-ring protein





• Molecule 1: Flagellar M-ring protein








• Molecule 1: Flagellar M-ring protein




• Molecule 1: Flagellar M-ring protein





• Molecule 1: Flagellar M-ring protein



• Molecule 1: Flagellar M-ring protein

Chain BA:




A multi-row visualization of Chain BA residues. Each row contains a list of amino acid abbreviations (e.g., MET, SER, ALA) and corresponding residue IDs (e.g., Q125, F126, S127). The residues are color-coded: green for high quality (34%), yellow for medium (10%), and grey for low or missing (55%). Red diamond markers are placed above certain residues to indicate specific quality issues or flags.

● Molecule 1: Flagellar M-ring protein



Chain CA:




A multi-row visualization of Chain CA residues. Each row contains a list of amino acid abbreviations (e.g., MET, SER, ALA) and corresponding residue IDs (e.g., Q125, F126, S127). The residues are color-coded: green for high quality (30%), yellow for medium (15%), and grey for low or missing (55%). Red diamond markers are placed above certain residues to indicate specific quality issues or flags.



• Molecule 1: Flagellar M-ring protein



• Molecule 1: Flagellar M-ring protein



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C33	Depositor
Number of particles used	16411	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	51.557	Depositor
Minimum defocus (nm)	1000	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	0.981	Depositor
Minimum map value	-0.272	Depositor
Average map value	-0.001	Depositor
Map value standard deviation	0.016	Depositor
Recommended contour level	0.173	Depositor
Map size (\AA)	1046.784, 1046.784, 1046.784	wwPDB
Map dimensions	768, 768, 768	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (\AA)	1.363, 1.363, 1.363	Depositor

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.25	0/1969	0.55	0/2668
1	AA	0.27	0/2017	0.56	3/2748 (0.1%)
1	B	0.28	1/2017 (0.0%)	0.62	2/2748 (0.1%)
1	BA	0.26	0/1969	0.55	1/2668 (0.0%)
1	C	0.26	0/1969	0.56	2/2668 (0.1%)
1	CA	0.26	0/1969	0.58	0/2668
1	D	0.25	0/1969	0.56	1/2668 (0.0%)
1	DA	0.25	0/2017	0.53	0/2748
1	E	0.25	0/2017	0.54	1/2748 (0.0%)
1	EA	0.26	0/1969	0.55	1/2668 (0.0%)
1	F	0.26	0/1969	0.56	0/2668
1	FA	0.25	0/1969	0.55	0/2668
1	G	0.26	0/1969	0.57	0/2668
1	GA	0.25	0/2017	0.53	1/2748 (0.0%)
1	H	0.25	0/2017	0.55	1/2748 (0.0%)
1	HA	0.26	0/1969	0.58	2/2668 (0.1%)
1	I	0.27	0/1969	0.58	4/2668 (0.1%)
1	J	0.26	0/1969	0.57	0/2668
1	K	0.28	1/2017 (0.0%)	0.60	1/2748 (0.0%)
1	L	0.26	0/1969	0.57	2/2668 (0.1%)
1	M	0.26	0/1969	0.58	0/2668
1	N	0.25	0/2017	0.53	0/2748
1	O	0.26	0/1969	0.58	3/2668 (0.1%)
1	P	0.26	0/1969	0.57	0/2668
1	Q	0.25	0/2017	0.53	1/2748 (0.0%)
1	R	0.25	0/1969	0.54	0/2668
1	S	0.26	0/1969	0.58	0/2668
1	T	0.25	0/2017	0.53	0/2748
1	V	0.53	2/1969 (0.1%)	0.74	3/2668 (0.1%)
1	W	0.26	0/1969	0.56	0/2668
1	X	0.28	0/2017	0.59	2/2748 (0.1%)
1	Y	0.25	0/1969	0.54	0/2668
1	Z	0.25	0/1969	0.55	0/2668
All	All	0.27	4/65505 (0.0%)	0.57	31/88924 (0.0%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	V	182	PRO	CG-CD	-18.78	0.88	1.50
1	V	182	PRO	N-CD	7.22	1.57	1.47
1	K	172	PRO	CA-CB	-5.63	1.42	1.53
1	B	172	PRO	CA-CB	-5.62	1.42	1.53

The worst 5 of 31 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	V	182	PRO	N-CD-CG	-19.71	73.63	103.20
1	B	172	PRO	N-CA-CB	-14.47	85.93	103.30
1	K	172	PRO	N-CA-CB	-14.42	85.99	103.30
1	X	172	PRO	N-CA-CB	-13.88	86.64	103.30
1	V	182	PRO	CA-CB-CG	-11.00	83.09	104.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1945	0	1931	78	0
1	AA	2005	0	1599	34	0
1	B	2005	0	1599	36	0
1	BA	1945	0	1931	58	0
1	C	1945	0	1931	58	0
1	CA	1945	0	1931	89	0
1	D	1945	0	1931	77	0
1	DA	2005	0	1599	38	0
1	E	2005	0	1599	36	0
1	EA	1945	0	1931	60	0
1	F	1945	0	1931	74	0
1	FA	1945	0	1931	82	0
1	G	1945	0	1931	73	0
1	GA	2005	0	1599	36	0
1	H	2005	0	1599	36	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	HA	1945	0	1931	64	0
1	I	1945	0	1931	68	0
1	J	1945	0	1931	82	0
1	K	2005	0	1599	40	0
1	L	1945	0	1931	64	0
1	M	1945	0	1931	79	0
1	N	2005	0	1599	32	0
1	O	1945	0	1931	60	0
1	P	1945	0	1931	90	0
1	Q	2005	0	1599	35	0
1	R	1945	0	1931	64	0
1	S	1945	0	1931	76	0
1	T	2005	0	1599	35	0
1	V	1945	0	1931	74	0
1	W	1945	0	1931	79	0
1	X	2005	0	1599	39	0
1	Y	1945	0	1931	69	0
1	Z	1945	0	1931	76	0
All	All	64845	0	60071	1553	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 1553 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:FA:191:ILE:HG13	1:FA:220:LEU:HB3	1.57	0.87
1:R:228:ARG:HD3	1:R:231:ASN:HD21	1.38	0.85
1:GA:409:MET:HE1	1:GA:434:ASN:HB2	1.61	0.83
1:CA:219:LEU:HD21	1:HA:193:ALA:HA	1.59	0.82
1:CA:191:ILE:HG13	1:CA:220:LEU:HB3	1.65	0.79

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	248/560 (44%)	237 (96%)	10 (4%)	1 (0%)	34	67
1	AA	303/560 (54%)	291 (96%)	9 (3%)	3 (1%)	15	46
1	B	303/560 (54%)	291 (96%)	9 (3%)	3 (1%)	15	46
1	BA	248/560 (44%)	242 (98%)	5 (2%)	1 (0%)	34	67
1	C	248/560 (44%)	242 (98%)	5 (2%)	1 (0%)	34	67
1	CA	248/560 (44%)	237 (96%)	10 (4%)	1 (0%)	34	67
1	D	248/560 (44%)	239 (96%)	8 (3%)	1 (0%)	34	67
1	DA	303/560 (54%)	292 (96%)	8 (3%)	3 (1%)	15	46
1	E	303/560 (54%)	290 (96%)	10 (3%)	3 (1%)	15	46
1	EA	248/560 (44%)	242 (98%)	5 (2%)	1 (0%)	34	67
1	F	248/560 (44%)	242 (98%)	5 (2%)	1 (0%)	34	67
1	FA	248/560 (44%)	239 (96%)	8 (3%)	1 (0%)	34	67
1	G	248/560 (44%)	238 (96%)	9 (4%)	1 (0%)	34	67
1	GA	303/560 (54%)	291 (96%)	9 (3%)	3 (1%)	15	46
1	H	303/560 (54%)	290 (96%)	10 (3%)	3 (1%)	15	46
1	HA	248/560 (44%)	242 (98%)	5 (2%)	1 (0%)	34	67
1	I	248/560 (44%)	240 (97%)	7 (3%)	1 (0%)	34	67
1	J	248/560 (44%)	239 (96%)	8 (3%)	1 (0%)	34	67
1	K	303/560 (54%)	292 (96%)	8 (3%)	3 (1%)	15	46
1	L	248/560 (44%)	242 (98%)	5 (2%)	1 (0%)	34	67
1	M	248/560 (44%)	238 (96%)	9 (4%)	1 (0%)	34	67
1	N	303/560 (54%)	293 (97%)	8 (3%)	2 (1%)	22	55
1	O	248/560 (44%)	243 (98%)	4 (2%)	1 (0%)	34	67
1	P	248/560 (44%)	238 (96%)	9 (4%)	1 (0%)	34	67
1	Q	303/560 (54%)	290 (96%)	10 (3%)	3 (1%)	15	46
1	R	248/560 (44%)	243 (98%)	4 (2%)	1 (0%)	34	67
1	S	248/560 (44%)	237 (96%)	10 (4%)	1 (0%)	34	67
1	T	303/560 (54%)	291 (96%)	9 (3%)	3 (1%)	15	46
1	V	248/560 (44%)	242 (98%)	5 (2%)	1 (0%)	34	67
1	W	248/560 (44%)	240 (97%)	7 (3%)	1 (0%)	34	67

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	X	303/560 (54%)	291 (96%)	9 (3%)	3 (1%)	15	46
1	Y	248/560 (44%)	242 (98%)	5 (2%)	1 (0%)	34	67
1	Z	248/560 (44%)	237 (96%)	10 (4%)	1 (0%)	34	67
All	All	8789/18480 (48%)	8483 (96%)	252 (3%)	54 (1%)	29	57

5 of 54 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	74	ASN
1	B	394	LYS
1	B	403	PRO
1	C	394	LYS
1	E	394	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	218/467 (47%)	215 (99%)	3 (1%)	67	83
1	AA	142/467 (30%)	140 (99%)	2 (1%)	67	83
1	B	142/467 (30%)	138 (97%)	4 (3%)	43	70
1	BA	218/467 (47%)	216 (99%)	2 (1%)	78	90
1	C	218/467 (47%)	213 (98%)	5 (2%)	50	74
1	CA	218/467 (47%)	215 (99%)	3 (1%)	67	83
1	D	218/467 (47%)	215 (99%)	3 (1%)	67	83
1	DA	142/467 (30%)	139 (98%)	3 (2%)	53	76
1	E	142/467 (30%)	140 (99%)	2 (1%)	67	83
1	EA	218/467 (47%)	215 (99%)	3 (1%)	67	83
1	F	218/467 (47%)	212 (97%)	6 (3%)	43	70
1	FA	218/467 (47%)	215 (99%)	3 (1%)	67	83
1	G	218/467 (47%)	215 (99%)	3 (1%)	67	83

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	GA	142/467 (30%)	140 (99%)	2 (1%)	67	83
1	H	142/467 (30%)	140 (99%)	2 (1%)	67	83
1	HA	218/467 (47%)	216 (99%)	2 (1%)	78	90
1	I	218/467 (47%)	216 (99%)	2 (1%)	78	90
1	J	218/467 (47%)	216 (99%)	2 (1%)	78	90
1	K	142/467 (30%)	140 (99%)	2 (1%)	67	83
1	L	218/467 (47%)	216 (99%)	2 (1%)	78	90
1	M	218/467 (47%)	213 (98%)	5 (2%)	50	74
1	N	142/467 (30%)	140 (99%)	2 (1%)	67	83
1	O	218/467 (47%)	215 (99%)	3 (1%)	67	83
1	P	218/467 (47%)	216 (99%)	2 (1%)	78	90
1	Q	142/467 (30%)	141 (99%)	1 (1%)	84	92
1	R	218/467 (47%)	214 (98%)	4 (2%)	59	79
1	S	218/467 (47%)	217 (100%)	1 (0%)	88	94
1	T	142/467 (30%)	140 (99%)	2 (1%)	67	83
1	V	218/467 (47%)	217 (100%)	1 (0%)	88	94
1	W	218/467 (47%)	213 (98%)	5 (2%)	50	74
1	X	142/467 (30%)	139 (98%)	3 (2%)	53	76
1	Y	218/467 (47%)	214 (98%)	4 (2%)	59	79
1	Z	218/467 (47%)	216 (99%)	2 (1%)	78	90
All	All	6358/15411 (41%)	6267 (99%)	91 (1%)	68	83

5 of 91 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	W	231	ASN
1	AA	259	ASN
1	W	415	LEU
1	Y	231	ASN
1	CA	219	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 26 such sidechains are listed below:

Mol	Chain	Res	Type
1	S	434	ASN
1	X	434	ASN
1	FA	434	ASN
1	W	131	ASN
1	Y	131	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-41101. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 384



Y Index: 384



Z Index: 384


6.2.2 Raw map



X Index: 384



Y Index: 384




Z Index: 384


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 348



Y Index: 419



Z Index: 295


6.3.2 Raw map



X Index: 348



Y Index: 384




Z Index: 295


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.173. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


6.6.1 emd_41101_msk_1.map [i](#)



X



Y




Z

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [\(i\)](#)



The volume at the recommended contour level is 1012 nm³; this corresponds to an approximate mass of 914 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [i](#)




*Reported resolution corresponds to spatial frequency of 0.294 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.294\AA^{-1}

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.40	-	-
Author-provided FSC curve	3.42	3.99	3.51
Unmasked-calculated*	8.14	12.64	8.28

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 8.14 differs from the reported value 3.4 by more than 10 %

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-41101 and PDB model 8T8P. Per-residue inclusion information can be found in section [3](#) on page [6](#).

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.173 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.173).





































































9.4 Atom inclusion [i](#)



At the recommended contour level, 86% of all backbone atoms, 80% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.173) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.7960	 0.3530
A	 0.8170	 0.3420
AA	 0.7580	 0.3710
B	 0.7590	 0.3650
BA	 0.8150	 0.3460
C	 0.8150	 0.3470
CA	 0.8160	 0.3400
D	 0.8210	 0.3430
DA	 0.7590	 0.3660
E	 0.7550	 0.3680
EA	 0.8100	 0.3460
F	 0.8170	 0.3470
FA	 0.8210	 0.3420
G	 0.8130	 0.3430
GA	 0.7550	 0.3670
H	 0.7540	 0.3700
HA	 0.8110	 0.3440
I	 0.8170	 0.3470
J	 0.8180	 0.3430
K	 0.7590	 0.3700
L	 0.8170	 0.3480
M	 0.8180	 0.3410
N	 0.7590	 0.3690
O	 0.8130	 0.3480
P	 0.8170	 0.3420
Q	 0.7580	 0.3680
R	 0.8100	 0.3460
S	 0.8210	 0.3420
T	 0.7520	 0.3700
V	 0.8130	 0.3490
W	 0.8230	 0.3430
X	 0.7500	 0.3680
Y	 0.8120	 0.3460
Z	 0.8240	 0.3450

