

Full wwPDB X-ray Structure Validation Report (i)

May 14, 2020 – 05:25 pm BST

PDB ID	:	1TA2
Title	:	Crystal structure of thrombin in complex with compound 1
Authors	:	Tucker, T.J.; Brady, S.F.; Lumma, W.C.; Lewis, S.D.; Gardel, S.J.; Naylor-
		Olsen, A.M.; Yan, Y.; Sisko, J.T.; Stauffer, K.J.; Lucas, B.Y.; Lynch, J.J.;
		Cook, J.J.; Stranieri, M.T.; Holahan, M.A.; Lyle, E.A.; Baskin, E.P.; Chen,
		IW.; Dancheck, K.B.; Krueger, J.A.; Cooper, C.M.; Vacca, J.P.
Deposited on	:	2004-05-19
Resolution	:	2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.


The following versions of software and data (see references (1)) were used in the production of this report:

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
Clashscore	141614	5643(2.30-2.30)
Ramachandran outliers	138981	5575(2.30-2.30)
Sidechain outliers	138945	5575(2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain				
1	А	287	75%	18%	•	•	
2	В	11	82%		18%		



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 2510 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called thrombin.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	А	276	Total 2228	C 1419	N 392	O 402	${ m S}$ 15	0	0	0

• Molecule 2 is a protein called Hirudin.


Mol	Chain	Residues		Ate	\mathbf{oms}			ZeroOcc	AltConf	Trace
2	В	11	Total 96	C 60	N 11	0 24	S 1	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
В	362	ALA	GLU	CONFLICT	UNP P28504
В	365	ALA	GLN	CONFLICT	UNP P28504

• Molecule 3 is 1-(2-AMINO-3,3-DIPHENYL-PROPIONYL)-PYRROLIDINE-3-CAR BOXYLIC ACID 2,5-DICHLORO-BENZYLAMIDE (three-letter code: 176) (formula: C₂₇H₂₇Cl₂N₃O₂).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
2	Δ	1	Total	С	Cl	Ν	Ο	0	0
0			34	27	2	3	2	0	U

• Molecule 4 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	150	Total O 150 150	0	0
4	В	2	Total O 2 2	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are colorcoded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.



• Molecule 1: thrombin



4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source	
Space group	C 1 2 1	Depositor	
Cell constants	70.81Å 72.36Å 72.96Å	Depositor	
a, b, c, α , β , γ	90.00° 100.90° 90.00°	Depositor	
Resolution (Å)	8.00 - 2.30	Depositor	
% Data completeness	(Not available) (8.00-2.30)	Depositor	
(in resolution range)	(1000 available) (0.00-2.00)		
R_{merge}	0.06	Depositor	
R_{sym}	0.06	Depositor	
Refinement program	X-PLOR 3.1	Depositor	
R, R_{free}	0.150 , (Not available)	Depositor	
Estimated twinning fraction	No twinning to report.	Xtriage	
Total number of atoms	2510	wwPDB-VP	
Average B, all atoms $(Å^2)$	32.0	wwPDB-VP	



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: 176, TYS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
	Cham	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	А	0.87	0/2282	1.64	48/3079~(1.6%)	
2	В	0.80	0/80	1.21	0/105	
All	All	0.87	0/2362	1.62	48/3184~(1.5%)	

There are no bond length outliers.

All (48) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
1	А	237	TRP	CD1-CG-CD2	9.64	114.01	106.30
1	А	29	TRP	CD1-CG-CD2	9.62	114.00	106.30
1	А	96	TRP	CD1-CG-CD2	9.03	113.52	106.30
1	А	73	ARG	NE-CZ-NH1	8.79	124.69	120.30
1	А	215	TRP	CD1-CG-CD2	8.70	113.26	106.30
1	А	207	TRP	CD1-CG-CD2	8.60	113.18	106.30
1	А	94	TYR	CB-CG-CD2	-8.21	116.07	121.00
1	А	60(D)	TRP	CD1-CG-CD2	8.02	112.71	106.30
1	А	96	TRP	CE2-CD2-CG	-7.88	101.00	107.30
1	А	141	TRP	CD1-CG-CD2	7.75	112.50	106.30
1	А	29	TRP	CE2-CD2-CG	-7.68	101.16	107.30
1	А	97	ARG	NE-CZ-NH2	-7.67	116.47	120.30
1	А	215	TRP	CE2-CD2-CG	-7.58	101.23	107.30
1	А	237	TRP	CE2-CD2-CG	-7.57	101.24	107.30
1	А	93	ARG	NE-CZ-NH1	-7.30	116.65	120.30
1	А	77(A)	ARG	NE-CZ-NH2	-7.15	116.72	120.30
1	А	207	TRP	CE2-CD2-CG	-7.11	101.61	107.30
1	А	141	TRP	CE2-CD2-CG	-6.98	101.72	107.30
1	А	165	ARG	NE-CZ-NH1	6.89	123.75	120.30
1	А	29	TRP	CG-CD1-NE1	-6.78	103.33	110.10
1	А	51	TRP	CE2-CD2-CG	-6.71	101.94	107.30
1	А	51	TRP	CG-CD2-CE3	6.67	139.90 Continued on m	133.90

Continued on next page...



Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
1	А	77(A)	ARG	NE-CZ-NH1	6.66	123.63	120.30
1	А	237	TRP	CG-CD1-NE1	-6.59	103.50	110.10
1	А	165	ARG	NE-CZ-NH2	-6.52	117.04	120.30
1	А	60(D)	TRP	CE2-CD2-CG	-6.48	102.12	107.30
1	А	225	TYR	CB-CG-CD2	-6.27	117.24	121.00
1	А	97	ARG	NE-CZ-NH1	6.27	123.43	120.30
1	А	231	VAL	CG1-CB-CG2	-6.26	100.89	110.90
1	А	14(A)	LYS	CA-CB-CG	6.20	127.03	113.40
1	А	51	TRP	CD1-CG-CD2	5.85	110.98	106.30
1	А	207	TRP	CG-CD1-NE1	-5.77	104.33	110.10
1	А	206	ARG	NE-CZ-NH2	5.76	123.18	120.30
1	А	173	ARG	NE-CZ-NH2	-5.71	117.44	120.30
1	А	96	TRP	CB-CG-CD1	-5.70	119.59	127.00
1	А	207	TRP	CG-CD2-CE3	5.65	138.98	133.90
1	А	41	LEU	CB-CG-CD1	-5.43	101.77	111.00
1	А	51	TRP	CB-CG-CD1	-5.43	119.94	127.00
1	А	106	MET	CG-SD-CE	-5.41	91.55	100.20
1	А	94	TYR	CB-CG-CD1	5.41	124.24	121.00
1	А	180	MET	CA-CB-CG	5.38	122.44	113.30
1	А	215	TRP	CG-CD2-CE3	5.31	138.68	133.90
1	А	57	HIS	CA-CB-CG	5.25	122.53	113.60
1	А	173	ARG	CA-CB-CG	5.25	124.95	113.40
1	А	96	TRP	CG-CD1-NE1	-5.20	104.90	110.10
1	А	173	ARG	NE-CZ-NH1	5.16	122.88	120.30
1	А	9	LYS	CA-CB-CG	-5.14	102.09	113.40
1	А	215	TRP	CB-CG-CD1	-5.05	120.43	127.00

Continued from previous page...

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	2228	0	2193	25	0
2	В	96	0	77	3	0
3	А	34	0	27	4	0

Continued on next page...



Continued from previous page...

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

A tom 1	Atom 9	Interatomic	Clash
Atom-1	Atom-2	${ m distance}~({ m \AA})$	overlap (Å)
1:A:38:GLN:HE22	2:B:364:LEU:HD21	1.62	0.64
1:A:75:ARG:HH11	2:B:357:GLU:HB2	1.63	0.63
1:A:14(G):LEU:HD21	1:A:202:LYS:HD3	1.82	0.61
3:A:401:176:HD1	3:A:401:176:HD2A	1.84	0.60
3:A:401:176:CD1	3:A:401:176:HD2A	2.34	0.58
1:A:18:GLU:HG3	1:A:187:ARG:HG3	1.88	0.56
1:A:33:LEU:HD12	1:A:42:CYS:HB2	1.89	0.54
1:A:16:ILE:N	4:A:796:HOH:O	2.42	0.53
1:A:38:GLN:NE2	2:B:364:LEU:HD21	2.25	0.51
1:A:73:ARG:HD2	4:A:660:HOH:O	2.11	0.50
1:A:217:GLU:HG2	3:A:401:176:CE4	2.41	0.50
1:A:107:LYS:NZ	1:A:246:GLY:HA3	2.28	0.49
1:A:67:ARG:HG2	1:A:82:ILE:HG12	1.95	0.48
1:A:165:ARG:NH2	1:A:177:THR:O	2.46	0.48
1:A:35:ARG:HB2	1:A:41:LEU:HD21	1.96	0.47
1:A:75:ARG:HA	1:A:75:ARG:HD3	1.68	0.47
1:A:110:LYS:HD3	1:A:111:PRO:CD	2.46	0.46
1:A:182:CYS:HA	1:A:226:GLY:O	2.17	0.45
1:A:99:LEU:HD11	3:A:401:176:HG1	2.00	0.44
1:A:85:LEU:HD22	1:A:106:MET:HB3	2.01	0.42
1:A:110:LYS:HD3	1:A:111:PRO:HD3	2.00	0.42
1:A:31:VAL:HB	1:A:44:ALA:HB3	2.02	0.42
1:A:186(D):LYS:HB2	1:A:186(D):LYS:NZ	2.35	0.42
1:A:211:GLY:HA2	1:A:229:THR:O	2.20	0.41
1:A:126:ARG:HB3	1:A:127:GLU:OE1	2.20	0.41
1:A:91:HIS:CE1	1:A:101:ARG:HD3	2.56	0.40
1:A:51:TRP:CZ2	1:A:107:LYS:HD3	2.56	0.40

There are no symmetry-related clashes.



Chain Non-H H(added) Clashes Symm-Clashes Mol H(model) 1504А $\mathbf{2}$ 0 0 0 4 В 2 0 0 0 0 All All 0 25100229727

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	А	270/287~(94%)	258~(96%)	12 (4%)	0	100	100
2	В	8/11~(73%)	8 (100%)	0	0	100	100
All	All	278/298~(93%)	266~(96%)	12 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	А	240/251~(96%)	230~(96%)	10~(4%)	30 42		
2	В	8/8~(100%)	8 (100%)	0	100 100		
All	All	248/259~(96%)	238~(96%)	10 (4%)	31 44		

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	11	SER
1	А	14(A)	LYS
1	А	41	LEU
1	А	112	VAL
1	А	124	PRO
1	А	126	ARG
1	А	151	GLN
1	А	182	CYS

Continued on next page...



Continued from previous page...

Mol	Chain	\mathbf{Res}	Type
1	А	186(D)	LYS
1	А	204(B)	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	А	30	GLN
1	А	156	GLN
1	А	204(B)	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Tune	Chain	Dec	Link	Bond lengths		Bond angles			
	Type	Chain	\mathbf{Res}		Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2
2	TYS	В	363	2	15, 16, 17	<mark>3.28</mark>	2 (13%)	18,22,24	1.12	<mark>1 (5%)</mark>

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	TYS	В	363	2	-	1/10/11/13	0/1/1/1

All (2) bond length outliers are listed below:



Mol	Chain	\mathbf{Res}	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(Å)
2	В	363	TYS	OH-CZ	-10.85	1.24	1.42
2	В	363	TYS	OH-S	-6.26	1.48	1.58

All (1) bond angle outliers are listed below:

Mol	Chain	\mathbf{Res}	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	В	363	TYS	OH-S-O2	3.06	116.62	107.71

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	В	363	TYS	O-C-CA-CB

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	e Chain	Res	Link	Bond lengths			Bond angles		
WIOI			ries		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z >2
3	176	А	401	-	$36,\!37,\!37$	0.70	0	$48,\!51,\!51$	1.27	5 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	176	А	401	-	-	4/29/39/39	0/4/4/4

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
3	А	401	176	C0-CA0-N0	-3.59	104.74	110.22
3	А	401	176	CB'-CG'-CL1	-3.00	114.86	119.73
3	А	401	176	CD'-CG'-CB'	2.86	126.82	122.20
3	А	401	176	CD'-CE'-CDB	-2.48	116.62	119.24
3	А	401	176	CD4-CG2-CD3	2.08	120.89	118.29

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	А	401	176	O-C-CA-N
3	А	401	176	N'-C-CA-N
3	А	401	176	N'-CA'-CB'-CG'
3	А	401	176	N0-CA0-CB0-CG2


There are no ring outliers.

1 monomer is involved in 4 short contacts:

\mathbb{N}	/lol	Chain	\mathbf{Res}	Type	Clashes	Symm-Clashes
	3	А	401	176	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

