



Full wwPDB X-ray Structure Validation Report

May 22, 2020 – 12:46 pm BST

PDB ID : 1TAZ
Title : Catalytic Domain Of Human Phosphodiesterase 1B
Authors : Zhang, K.Y.J.; Card, G.L.; Suzuki, Y.; Artis, D.R.; Fong, D.; Gillette, S.;
Hsieh, D.; Neiman, J.; West, B.L.; Zhang, C.; Milburn, M.V.; Kim, S.-H.;
Schlessinger, J.; Bollag, G.
Deposited on : 2004-05-19
Resolution : 1.77 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.77 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	9185 (1.80-1.76)
Clashscore	141614	10184 (1.80-1.76)
Ramachandran outliers	138981	10051 (1.80-1.76)
Sidechain outliers	138945	10050 (1.80-1.76)
RSRZ outliers	127900	9032 (1.80-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	365	

2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 2900 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Calcium/calmodulin-dependent 3',5'-cyclic nucleotide phosphodiesterase 1B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	322	2691	1720	464	491	16	0	10	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	142	GLY	-	EXPRESSION TAG	UNP Q01064
A	143	SER	-	EXPRESSION TAG	UNP Q01064
A	144	HIS	-	EXPRESSION TAG	UNP Q01064
A	145	MET	-	EXPRESSION TAG	UNP Q01064

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mg	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	207	Total	O	0	0
			207	207		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Calcium/calmodulin-dependent 3',5'-cyclic nucleotide phosphodiesterase 1B



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	87.23Å 87.23Å 134.15Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	72.55 – 1.77 53.17 – 1.70	Depositor EDS
% Data completeness (in resolution range)	99.7 (72.55-1.77) 99.5 (53.17-1.70)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.26 (at 1.70Å)	Xtrriage
Refinement program	REFMAC 5.1.25	Depositor
R, R_{free}	0.182 , 0.193 0.191 , 0.185	Depositor DCC
R_{free} test set	2913 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	28.8	Xtrriage
Anisotropy	0.383	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 44.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2900	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.22% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ZN, CME

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.38	0/2738	0.66	3/3705 (0.1%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	426	ASP	CB-CG-OD2	5.64	123.38	118.30
1	A	438	ASP	CB-CG-OD2	5.50	123.25	118.30
1	A	167	ASP	CB-CG-OD2	5.26	123.04	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2691	0	2667	31	0
2	A	1	0	0	0	0
3	A	1	0	0	0	0
4	A	207	0	0	0	0
All	All	2900	0	2667	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:238[B]:PHE:CE1	1:A:365[B]:LEU:HD11	1.78	1.17
1:A:381:HIS:HE1	1:A:496:TRP:HE1	1.16	0.93
1:A:238[B]:PHE:CE1	1:A:365[B]:LEU:CD1	2.57	0.88
1:A:238[B]:PHE:CD1	1:A:365[B]:LEU:HD11	2.08	0.88
1:A:238[A]:PHE:HD1	1:A:365[A]:LEU:CD1	1.88	0.86
1:A:238[A]:PHE:HD1	1:A:365[A]:LEU:HD11	1.47	0.77
1:A:238[A]:PHE:CD1	1:A:365[A]:LEU:HD11	2.29	0.66
1:A:238[A]:PHE:CD1	1:A:365[A]:LEU:CD1	2.76	0.63
1:A:373:HIS:ND1	1:A:381:HIS:HD2	1.97	0.61
1:A:238[B]:PHE:CD1	1:A:365[B]:LEU:CD1	2.82	0.60
1:A:238[A]:PHE:HD1	1:A:365[A]:LEU:HD12	1.67	0.57
1:A:365[B]:LEU:HD13	1:A:436:LEU:HD11	1.86	0.57
1:A:164[A]:TRP:HA	1:A:237:CME:HZ3	1.87	0.57
1:A:178:HIS:HD2	1:A:212:GLU:OE1	1.88	0.56
1:A:381:HIS:CE1	1:A:496:TRP:HE1	2.08	0.56
1:A:164[A]:TRP:CZ2	1:A:485:TRP:HB3	2.45	0.52
1:A:164[B]:TRP:CD1	1:A:484:THR:HG21	2.49	0.47
1:A:238[B]:PHE:HE1	1:A:365[B]:LEU:HD11	1.65	0.47
1:A:214:GLY:HA3	1:A:265:TYR:CE2	2.50	0.46
1:A:341:GLN:O	1:A:345:THR:HG23	2.16	0.45
1:A:358:LYS:HB3	1:A:359:PRO:HD3	1.98	0.45
1:A:425:ILE:HG23	1:A:489:ILE:HD12	1.99	0.45
1:A:183:ILE:HD12	1:A:229:ALA:HB1	1.99	0.45
1:A:180:LEU:H	1:A:225:GLN:NE2	2.16	0.44
1:A:220:ASN:HD22	1:A:394[B]:ARG:HH22	1.66	0.43
1:A:373:HIS:ND1	1:A:381:HIS:CD2	2.81	0.43
1:A:381:HIS:HE1	1:A:496:TRP:NE1	1.99	0.42
1:A:192:ASN:HA	1:A:192:ASN:HD22	1.67	0.41
1:A:425:ILE:CG2	1:A:489:ILE:HG23	2.50	0.41
1:A:374:PRO:HA	1:A:381:HIS:CD2	2.56	0.41
1:A:164[B]:TRP:HA	1:A:237:CME:HZ3	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	323/365 (88%)	319 (99%)	4 (1%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	299/328 (91%)	288 (96%)	11 (4%)	34 17

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	164[A]	TRP
1	A	164[B]	TRP
1	A	192	ASN
1	A	219	LYS
1	A	225	GLN
1	A	251[A]	ILE
1	A	251[B]	ILE
1	A	310	ASN
1	A	355	ARG
1	A	388	LEU
1	A	502	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	172	ASN
1	A	178	HIS
1	A	192	ASN
1	A	220	ASN
1	A	225	GLN
1	A	310	ASN
1	A	314	ASN
1	A	381	HIS
1	A	419	GLN
1	A	445	GLN
1	A	492	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	CME	A	237	1	8,9,10	1.48	1 (12%)	5,9,11	1.54	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CME	A	237	1	-	3/5/8/10	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	237	CME	OH-CZ	-3.68	1.23	1.42

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	237	CME	OH-CZ-CE	2.06	118.94	110.83

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	237	CME	CZ-CE-SD-SG
1	A	237	CME	N-CA-CB-SG
1	A	237	CME	CE-SD-SG-CB

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	237	CME	2	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	321/365 (87%)	0.42	31 (9%) 7 7	14, 23, 43, 55	0

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	501	ALA	7.6
1	A	473	ASP	7.2
1	A	474	PRO	6.0
1	A	164[A]	TRP	5.7
1	A	500	ALA	5.2
1	A	445	GLN	4.1
1	A	412	ARG	4.0
1	A	404	LEU	3.9
1	A	413	THR	3.7
1	A	502	SER	3.7
1	A	405	PRO	3.6
1	A	402	LEU	3.3
1	A	238[A]	PHE	3.1
1	A	415	THR	3.0
1	A	400	ALA	3.0
1	A	494	GLN	2.9
1	A	414	SER	2.8
1	A	490	GLN	2.7
1	A	397	ASP	2.6
1	A	401	GLU	2.5
1	A	499	ARG	2.5
1	A	307	ASP	2.5
1	A	441	GLU	2.5
1	A	479	VAL	2.2
1	A	475	ASN	2.2
1	A	150	TYR	2.1
1	A	355	ARG	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	398	LYS	2.1
1	A	146	VAL	2.1
1	A	351	GLN	2.0
1	A	393	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CME	A	237	10/11	0.82	0.16	21,22,40,41	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	MG	A	1002	1/1	0.99	0.08	17,17,17,17	0
2	ZN	A	1001	1/1	1.00	0.06	22,22,22,22	0

6.5 Other polymers [i](#)

There are no such residues in this entry.