

# Full wwPDB X-ray Structure Validation Report (i)

#### May 25, 2020 - 05:58 am BST

) :	5TCH
е :	Crystal structure of tryptophan synthase from M. tuberculosis - ligand-free
	form, TrpA-G66V mutant
s :	Michalska, K.; Maltseva, N.; Jedrzejczak, R.; Wellington, S.; Nag, P.P.; Fisher,
	S.L.; Schreiber, S.L.; Hung, D.T.; Joachimiak, A.; Center for Structural Ge-
	nomics of Infectious Diseases (CSGID)
n :	2016-09-15
n :	2.35  Å(reported)
	D : e : s : n : n :

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as 541 be (2020)
Xtriage (Phenix)	:	1.13
$\mathrm{EDS}$	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
$\operatorname{Refmac}$	:	5.8.0158
$\operatorname{CCP4}$	:	7.0.044  (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11

## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R <sub>free</sub>	130704	1164 (2.36-2.36)
Clashscore	141614	1232(2.36-2.36)
Ramachandran outliers	138981	1211(2.36-2.36)
Sidechain outliers	138945	1212(2.36-2.36)
RSRZ outliers	127900	$1150 \ (2.36-2.36)$

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	Δ	276	3%	100/
	A	270	88%	• 10%
1	С	276	89%	• 10%
1	Е	276	87%	• 11%
1	G	276	2% 	• 11%
2	В	410	% 96%	•••
2	D	410	% 95%	•••



Mol	Chain	Length	Quality of chain	
2	F	410	% 95% •	•
2	Н	410	2% 95%	•••



## 5TCH

## 2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 20207 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	Δ	240	Total	С	Ν	Ο	S	0	2	0
		249	1826	1145	329	347	5	0	Z	0
1	C	247	Total	С	Ν	Ο	S	0	2	0
	I G	247	1815	1139	329	342	5	0	2	0
1	F	9.46	Total	С	Ν	Ο	S	0	0	0
		240	1785	1121	320	339	5			0
1	1 C	248	Total	С	Ν	Ο	S	0	0	0
	248	1802	1132	322	343	5	0	0	0	

• Molecule 1 is a protein called Tryptophan synthase alpha chain.

There are 28 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	66	VAL	GLY	engineered mutation	UNP P9WFY1
A	271	HIS	-	expression tag	UNP P9WFY1
A	272	HIS	-	expression tag	UNP P9WFY1
A	273	HIS	-	expression tag	UNP P9WFY1
A	274	HIS	-	expression tag	UNP P9WFY1
А	275	HIS	-	expression tag	UNP P9WFY1
А	276	HIS	-	expression tag	UNP P9WFY1
G	66	VAL	GLY	engineered mutation	UNP P9WFY1
G	271	HIS	-	expression tag	UNP P9WFY1
G	272	HIS	-	expression tag	UNP P9WFY1
G	273	HIS	-	expression tag	UNP P9WFY1
G	274	HIS	-	expression tag	UNP P9WFY1
G	275	HIS	-	expression tag	UNP P9WFY1
G	276	HIS	-	expression tag	UNP P9WFY1
Е	66	VAL	GLY	engineered mutation	UNP P9WFY1
E	271	HIS	-	expression tag	UNP P9WFY1
E	272	HIS	-	expression tag	UNP P9WFY1
Е	273	HIS	-	expression tag	UNP P9WFY1
Е	274	HIS	-	expression tag	UNP P9WFY1
Е	275	HIS	-	expression tag	UNP P9WFY1
Е	276	HIS	-	expression tag	UNP P9WFY1



Chain	Residue	Modelled	Actual	Comment	Reference
С	66	VAL	GLY	engineered mutation	UNP P9WFY1
С	271	HIS	-	expression tag	UNP P9WFY1
С	272	HIS	-	expression tag	UNP P9WFY1
С	273	HIS	-	expression tag	UNP P9WFY1
С	274	HIS	-	expression tag	UNP P9WFY1
С	275	HIS	-	expression tag	UNP P9WFY1
С	276	HIS	-	expression tag	UNP P9WFY1

• Molecule 2 is a protein called Tryptophan synthase beta chain.

Mol	Chain	Residues		Atoms						AltConf	Trace
9	В	405	Total	С	Ν	Ο	Р	S	0	2	0
	D	400	3047	1901	551	580	1	14	0		0
9	Ц	406	Total	С	Ν	Ο	Р	S	0	1	0
		400	3044	1899	551	580	1	13	0		0
0	Б	200	Total	С	Ν	Ο	Р	S	0	1	0
		599	2995	1869	540	572	1	13	0		
2 D	398	Total	С	Ν	Ο	Р	S	0	0	0	
		2980	1861	538	567	1	13		0		

• Molecule 3 is MALONATE ION (three-letter code: MLI) (formula:  $C_3H_2O_4$ ).



Mo	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	1	$\begin{array}{ccc} \text{Total}  \text{C}  \text{O} \\ 7  3  4 \end{array}$	0	0
3	В	1	$\begin{array}{ccc} \text{Total}  \text{C}  \text{O} \\ 7  3  4 \end{array}$	0	0



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	G	1	$\begin{array}{ccc} \text{Total}  \text{C}  \text{O} \\ 7  3  4 \end{array}$	0	0
3	Н	1	$\begin{array}{ccc} \text{Total}  \text{C}  \text{O} \\ 7  3  4 \end{array}$	0	0
3	Ε	1	$\begin{array}{ccc} \text{Total}  \text{C}  \text{O} \\ 7  3  4 \end{array}$	0	0
3	F	1	$\begin{array}{ccc} \text{Total}  \text{C}  \text{O} \\ 7  3  4 \end{array}$	0	0
3	С	1	$\begin{array}{ccc} \text{Total}  \text{C}  \text{O} \\ 7  3  4 \end{array}$	0	0
3	D	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 7 & 3 & 4 \end{array}$	0	0

• Molecule 4 is FORMIC ACID (three-letter code: FMT) (formula:  $CH_2O_2$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	1	$\begin{array}{ccc} \text{Total}  \text{C}  \text{O} \\ 3  1  2 \end{array}$	0	0
4	В	1	$\begin{array}{ccc} \text{Total}  \text{C}  \text{O} \\ 3  1  2 \end{array}$	0	0
4	В	1	$\begin{array}{ccc} \text{Total}  \text{C}  \text{O} \\ 3  1  2 \end{array}$	0	0
4	В	1	$\begin{array}{ccc} \text{Total}  \text{C}  \text{O} \\ 3  1  2 \end{array}$	0	0
4	G	1	$\begin{array}{ccc} \text{Total}  \text{C}  \text{O} \\ 3  1  2 \end{array}$	0	0



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	G	1	$\begin{array}{ccc} \text{Total}  \text{C}  \text{O} \\ 3  1  2 \end{array}$	0	0
4	Н	1	$\begin{array}{ccc} \text{Total}  \text{C}  \text{O} \\ 3  1  2 \end{array}$	0	0
4	Н	1	$\begin{array}{ccc} \text{Total}  \text{C}  \text{O} \\ 3  1  2 \end{array}$	0	0
4	Н	1	$\begin{array}{ccc} \text{Total}  \text{C}  \text{O} \\ 3  1  2 \end{array}$	0	0
4	Н	1	$\begin{array}{ccc} \text{Total}  \text{C}  \text{O} \\ 3  1  2 \end{array}$	0	0
4	Ε	1	$\begin{array}{ccc} \text{Total}  \text{C}  \text{O} \\ 3  1  2 \end{array}$	0	0
4	Ε	1	$\begin{array}{ccc} \text{Total}  \text{C}  \text{O} \\ 3  1  2 \end{array}$	0	0
4	F	1	$\begin{array}{ccc} \text{Total}  \text{C}  \text{O} \\ 3  1  2 \end{array}$	0	0
4	F	1	$\begin{array}{ccc} \text{Total}  \text{C}  \text{O} \\ 3  1  2 \end{array}$	0	0
4	С	1	$\begin{array}{ccc} \text{Total}  \text{C}  \text{O} \\ 3  1  2 \end{array}$	0	0
4	D	1	$\begin{array}{ccc} \text{Total}  \text{C}  \text{O} \\ 3  1  2 \end{array}$	0	0
4	D	1	$\begin{array}{ccc} \text{Total}  \text{C}  \text{O} \\ 3  1  2 \end{array}$	0	0
4	D	1	$\begin{array}{ccc} \overline{\text{Total}} & \text{C} & \text{O} \\ 3 & 1 & 2 \end{array}$	0	0
4	D	1	$\begin{array}{c cc} \hline \text{Total} & \text{C} & \text{O} \\ \hline 3 & 1 & 2 \end{array}$	0	0
4	D	1	$\begin{array}{ccc} \text{Total}  \text{C}  \text{O} \\ 3  1  2 \end{array}$	0	0

• Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	А	44	$\begin{array}{cc} \text{Total} & \text{O} \\ 44 & 44 \end{array}$	0	1
5	В	177	Total O 177 177	0	3
5	G	73	Total O 73 73	0	0
5	Н	171	Total O 172 172	0	2



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	Е	9	Total O 9 9	0	0
5	F	122	Total         O           123         123	0	1
5	С	50	$\begin{array}{cc} {\rm Total} & {\rm O} \\ 50 & 50 \end{array}$	0	0
5	D	149	Total O 149 149	0	0



## 3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Chain A: 88% 10% MET VAL VAL GLU GLU GLY GLY GLY GLY ALA ASF <u>ار ق</u> • Molecule 1: Tryptophan synthase alpha chain Chain G: 88% 11% MET VAL GLU GLU GLU GLU THR MET GLY GLY GLY GLY ALA ALA ASP ASP SER HIS HIS HIS HIS HIS HIS • Molecule 1: Tryptophan synthase alpha chain Chain E: 87% 11% THR AET ALA ALA ASP MET VAL VAL GLU GLU SER LEU GLY MET MET ALA HIS HIS HIS HIS HIS • Molecule 1: Tryptophan synthase alpha chain Chain C: 89% . 10% MET VAL ALA ALA GLU GLU SER SER MET GLY GLY GLY GLY ALA ASF ALA ASF • Molecule 2: Tryptophan synthase beta chain Chain B: 96% . .
- Molecule 1: Tryptophan synthase alpha chain







## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	134.32Å $158.81$ Å $166.22$ Å	Deperitor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.00^{\circ}$ $90.00^{\circ}$	Depositor
$\mathbf{P}_{\text{acclution}}(\hat{\mathbf{A}})$	30.00 - 2.35	Depositor
Resolution (A)	29.91 - 2.35	EDS
% Data completeness	99.7 (30.00-2.35)	Depositor
(in resolution range)	$99.8 \ (29.91-2.35)$	EDS
R <sub>merge</sub>	0.12	Depositor
R <sub>sym</sub>	(Not available)	Depositor
$< I/\sigma(I) > 1$	$2.71 (at 2.34 \text{\AA})$	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
D D .	0.175 , $0.206$	Depositor
$\Pi, \Pi_{free}$	0.180 , $0.207$	DCC
$R_{free}$ test set	2883 reflections $(1.95%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	34.4	Xtriage
Anisotropy	0.368	Xtriage
Bulk solvent $k_{sol}(e/A^3), B_{sol}(A^2)$	0.33 , $39.0$	EDS
L-test for twinning <sup>2</sup>	$<  L  > = 0.50, < L^2 > = 0.34$	Xtriage
Estimated twinning fraction	0.006 for -h,l,k	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	20207	wwPDB-VP
Average B, all atoms $(Å^2)$	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 2.91% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



 $<sup>^1 {\</sup>rm Intensities}$  estimated from amplitudes.

## 5 Model quality (i)

## 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: FMT, MLI, LLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bo	nd lengths	Bo	ond angles
	Chain	RMSZ	# Z  > 5	RMSZ	# Z  > 5
1	А	0.52	0/1855	0.70	0/2531
1	С	0.53	0/1831	0.70	0/2500
1	Е	0.54	1/1814~(0.1%)	0.67	0/2477
1	G	0.53	0/1844	0.70	0/2516
2	В	0.56	0/3083	0.73	0/4177
2	D	0.59	0/3015	0.74	0/4085
2	F	0.55	0/3030	0.71	0/4106
2	H	0.59	0/3080	0.77	1/4174~(0.0%)
All	All	0.56	1/19552~(0.0%)	0.72	1/26566~(0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	$\mathbf{Res}$	Type	Atoms	Z	Observed(A)	$\operatorname{Ideal}(\operatorname{\AA})$
1	Ε	50	GLU	CG-CD	5.10	1.59	1.51

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	Н	220	ARG	NE-CZ-NH2	5.49	123.04	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	1826	0	1847	1	0
1	С	1802	0	1825	0	0
1	Е	1785	0	1808	1	0
1	G	1815	0	1843	0	0
2	В	3047	0	2959	7	0
2	D	2980	0	2894	2	0
2	F	2995	0	2904	3	0
2	Η	3044	0	2956	6	0
3	А	7	0	2	0	0
3	В	7	0	2	0	0
3	С	7	0	2	0	0
3	D	7	0	2	0	0
3	Ε	7	0	2	0	0
3	F	7	0	2	0	0
3	G	7	0	2	0	0
3	Н	7	0	2	0	0
4	А	3	0	1	0	0
4	В	9	0	3	0	0
4	С	3	0	1	0	0
4	D	15	0	5	0	0
4	Е	6	0	2	0	0
4	F	6	0	2	0	0
4	G	6	0	2	0	0
4	Η	12	0	4	0	0
5	А	44	0	0	0	0
5	В	177	0	0	1	0
5	С	50	0	0	0	0
5	D	149	0	0	0	0
5	Е	9	0	0	0	0
5	F	123	0	0	0	0
5	G	73	0	0	0	0
5	Н	172	0	0	0	0
All	All	20207	0	19072	19	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All (19) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:F:230:ILE:HG21	2:F:238:PRO:HD3	1.87	0.56
2:H:34:LEU:HD11	2:H:188:PHE:HD2	1.76	0.51



Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	${ m distance}~({ m \AA})$	overlap (Å)
2:B:242:VAL:HG12	2:B:267:VAL:HB	1.92	0.50
2:D:129:HIS:CE1	2:D:203:GLY:HA2	2.47	0.49
2:B:113[B]:ARG:HG2	5:B:603:HOH:O	2.11	0.49
2:H:207:GLY:HA2	2:H:294:HIS:O	2.12	0.49
2:H:101:LLP:NZ	2:H:101:LLP:O3	2.38	0.48
2:D:123:GLU:HG3	2:D:184:ILE:CG1	2.45	0.47
2:B:123:GLU:OE1	2:B:200:TYR:OH	2.30	0.47
2:F:123:GLU:HG3	2:F:184:ILE:HG12	1.97	0.45
2:B:207:GLY:HA2	2:B:294:HIS:O	2.18	0.44
1:E:182:ALA:HB1	1:E:202:LEU:HD23	2.01	0.43
2:F:207:GLY:HA2	2:F:294:HIS:O	2.19	0.43
1:A:140:ASP:N	1:A:140:ASP:OD1	2.50	0.42
2:B:129:HIS:CE1	2:B:203:GLY:HA2	2.55	0.42
2:B:5:ILE:HD11	2:H:225:GLU:HG2	2.01	0.42
2:H:230:ILE:HG21	2:H:238:PRO:HD3	2.03	0.41
2:H:129:HIS:CE1	2:H:203:GLY:HA2	2.56	0.40
2:B:402:LYS:HG3	2:B:407:LEU:HD12	2.03	0.40

There are no symmetry-related clashes.

#### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	$\mathbf{ntiles}$
1	А	247/276~(90%)	244~(99%)	2 (1%)	1 (0%)	34	38
1	С	244/276~(88%)	239~(98%)	5 (2%)	0	100	100
1	Е	242/276~(88%)	236~(98%)	5 (2%)	1 (0%)	34	38
1	G	245/276~(89%)	240~(98%)	4 (2%)	1 (0%)	34	38
2	В	404/410~(98%)	399~(99%)	5(1%)	0	100	100
2	D	395/410~(96%)	387~(98%)	8 (2%)	0	100	100
2	F	397/410~(97%)	$388 \ (98\%)$	9 $(2\%)$	0	100	100



0 0 1000	contracta from proceeder pagette										
Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles				
2	Η	404/410~(98%)	397~(98%)	7(2%)	0	100	100				
All	All	2578/2744~(94%)	2530~(98%)	45~(2%)	3~(0%)	51	63				

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	А	239	GLY
1	Е	239	GLY
1	G	239	GLY

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	182/201~(90%)	178~(98%)	4 (2%)	52 63
1	С	180/201~(90%)	178~(99%)	2(1%)	73 84
1	Ε	178/201~(89%)	175~(98%)	3~(2%)	60 72
1	G	181/201~(90%)	179~(99%)	2(1%)	73 84
2	В	300/301~(100%)	298~(99%)	2(1%)	84 91
2	D	293/301~(97%)	290~(99%)	3~(1%)	76 85
2	F	295/301~(98%)	293~(99%)	2(1%)	84 91
2	Η	299/301~(99%)	295 (99%)	4 (1%)	69 80
All	All	1908/2008~(95%)	1886 (99%)	22 (1%)	71 82

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	53	CYS
1	А	140	ASP
1	А	169	LEU
1	А	181	TYR
2	В	123	GLU



Mol	Chain	Res	Type
2	В	256	HIS
1	G	140	ASP
1	G	181	TYR
2	Н	128	GLN
2	Н	256	HIS
2	Н	300	LEU
2	Н	315	SER
1	Е	140	ASP
1	Е	149	SER
1	Е	181	TYR
2	F	23	SER
2	F	256	HIS
1	С	140	ASP
1	С	181	TYR
2	D	128	GLN
2	D	182	ASP
2	D	256	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Tune	Chain	Res	Link	Link Bond lengths			Bond angles		
	туре				Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	LLP	F	101	2	23,24,25	2.49	6 (26%)	25,32,34	1.46	3 (12%)
2	LLP	D	101	2	23,24,25	2.62	7 (30%)	25,32,34	1.66	6 (24%)
2	LLP	В	101	2	23,24,25	2.69	5 (21%)	25,32,34	1.61	4 (16%)



Mol	Type	Chain	$\mathbf{Res}$	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	LLP	Н	101	2	23,24,25	2.64	7 (30%)	25,32,34	1.69	6 (24%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	LLP	F	101	2	-	4/16/17/19	0/1/1/1
2	LLP	D	101	2	-	3/16/17/19	0/1/1/1
2	LLP	В	101	2	-	2/16/17/19	0/1/1/1
2	LLP	Н	101	2	-	5/16/17/19	0/1/1/1

All (25) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(Å)
2	В	101	LLP	C4-C4'	8.86	1.63	1.46
2	D	101	LLP	C4-C4'	8.55	1.62	1.46
2	Н	101	LLP	C4-C4'	8.06	1.61	1.46
2	F	101	LLP	C4-C4'	7.97	1.61	1.46
2	В	101	LLP	C4'-NZ	5.32	1.45	1.27
2	Н	101	LLP	C4-C5	-5.24	1.35	1.42
2	F	101	LLP	C4'-NZ	5.16	1.44	1.27
2	D	101	LLP	C4'-NZ	5.09	1.44	1.27
2	Н	101	LLP	C4'-NZ	4.87	1.43	1.27
2	В	101	LLP	C4-C5	-3.61	1.37	1.42
2	F	101	LLP	C4-C5	25 -3.49 1.37		1.42
2	D	101	LLP	C4-C5	-3.38	1.37	1.42
2	D	101	LLP	C6-N1	3.18	1.41	1.34
2	В	101	LLP	C2'-C2	2.99	1.55	1.50
2	Н	101	LLP	C2'-C2	2.86	1.55	1.50
2	F	101	LLP	C6-N1	2.85	1.40	1.34
2	В	101	LLP	C6-N1	2.81	1.40	1.34
2	D	101	LLP	C2'-C2	2.78	1.55	1.50
2	F	101	LLP	C2'-C2	2.62	1.54	1.50
2	Н	101	LLP	C6-N1	2.58	1.39	1.34
2	Н	101	LLP	CB-CA	-2.31	1.50	1.53
2	D	101	LLP	P-OP1	2.29	1.57	1.50
2	Н	101	LLP	$\overline{\text{C5'-C5}}$	2.20	1.56	1.50
2	D	101	LLP	C5'-C5	2.17	1.56	1.50
2	F	101	LLP	P-OP4	-2.12	1.53	1.60



Mol

2

2

2 2

2

2

2

2 2

2

2

2 2

2

2

2

2

2

2

F

Η

D

D

В

) bond ar	) bond angle outliers are listed below:											
Chain	Res	Type	Atoms	Z	$Observed(^{o})$	Ideal( <sup>o</sup>						
D	101	LLP	C4-C4'-NZ	-4.98	101.44	124.31						
В	101	LLP	C4-C4'-NZ	-4.90	101.81	124.31						
F	101	LLP	C4-C4'-NZ	-4.44	103.93	124.31						
Н	101	LLP	C4-C4'-NZ	-4.28	104.64	124.31						
Н	101	LLP	C3-C4-C5	3.50	120.95	118.26						
В	101	LLP	OP4-C5'-C5	3.27	115.59	109.35						
D	101	LLP	C3-C4-C4'	-2.93	114.95	120.41						
D	101	LLP	OP4-C5'-C5	2.37	113.87	109.35						
F	101	LLP	C2'-C2-N1	2.28	122.12	117.67						
D	101	LLP	C3-C4-C5	2.27	120.00	118.26						
В	101	LLP	OP4-P-OP1	2.25	112.80	106.47						
Н	101	LLP	C2'-C2-N1	2.23	122.02	117.67						
H	101	LLP	OP4-C5'-C5	2.18	113.51	109.35						
Н	101	LLP	$C\overline{E}-NZ-C4'$	-2.18	112.20	118.90						

C2'-C2-C3

C2'-C2-C3

C5-C4-C4'

C2'-C2-C3

C5-C6-N1

-2.16

-2.10

2.08

-2.07

-2.06

118.22

118.29

124.97

118.33

120.39

All (19)

There are no chirality outliers.

All (14) torsion outliers are listed below:

101

101

101

101

101

LLP

LLP

LLP

LLP

LLP

Mol	Chain	Res	Type	Atoms
2	Н	101	LLP	C4-C4'-NZ-CE
2	Н	101	LLP	C5'-OP4-P-OP1
2	F	101	LLP	C4-C4'-NZ-CE
2	Н	101	LLP	CG-CD-CE-NZ
2	F	101	LLP	CG-CD-CE-NZ
2	D	101	LLP	C3-C4-C4'-NZ
2	F	101	LLP	CD-CE-NZ-C4'
2	В	101	LLP	CD-CE-NZ-C4'
2	Н	101	LLP	CD-CE-NZ-C4'
2	F	101	LLP	C3-C4-C4'-NZ
2	D	101	LLP	C5-C4-C4'-NZ
2	D	101	LLP	CD-CE-NZ-C4'
2	В	101	LLP	C3-C4-C4'-NZ
2	Н	101	LLP	C3-C4-C4'-NZ

There are no ring outliers.

1 monomer is involved in 1 short contact:

 $(^{o})$ 

120.89

120.89

121.56

120.89

123.82



Mol	Chain	Res	Type	Clashes	Symm-Clashes	
2	Н	101	LLP	1	0	

### 5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry (i)

28 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Type		Chain	Dog	Tink	B	Bond lengths			Bond angles		
	Type	Ullalli	nes		Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2	
4	FMT	D	503	-	0,2,2	0.00	-	$_{0,1,1}$	0.00	-	
4	FMT	D	506	-	0,2,2	0.00	-	$_{0,1,1}$	0.00	-	
3	MLI	G	301	-	$0,\!6,\!6$	0.00	-	$0,\!7,\!7$	0.00	-	
4	FMT	G	302	-	0,2,2	0.00	-	$_{0,1,1}$	0.00	-	
4	FMT	Н	504	-	0,2,2	0.00	-	$_{0,1,1}$	0.00	-	
3	MLI	H	501	-	$0,\!6,\!6$	0.00	-	$_{0,7,7}$	0.00	-	
3	MLI	D	501	-	0,6,6	0.00	-	0,7,7	0.00	-	
4	FMT	В	503	-	0,2,2	0.00	-	$_{0,1,1}$	0.00	-	
4	FMT	Н	502	-	0,2,2	0.00	-	$_{0,1,1}$	0.00	-	
4	FMT	E	303	-	0,2,2	0.00	-	$_{0,1,1}$	0.00	-	
3	MLI	E	301	-	$0,\!6,\!6$	0.00	-	$_{0,7,7}$	0.00	-	
3	MLI	В	501	-	0,6,6	0.00	-	0,7,7	0.00	-	
4	FMT	A	302	-	0,2,2	0.00	-	$_{0,1,1}$	0.00	-	
3	MLI	А	301	-	$0,\!6,\!6$	0.00	-	$0,\!7,\!7$	0.00	-	
4	FMT	В	502	-	0,2,2	0.00	-	$_{0,1,1}$	0.00	-	
4	FMT	Н	503	-	0,2,2	0.00	-	$_{0,1,1}$	0.00	-	
4	FMT	D	505	-	0,2,2	0.00	-	$_{0,1,1}$	0.00	-	
4	FMT	Е	302	-	0,2,2	0.00	-	$_{0,1,1}$	0.00	-	
4	FMT	С	302	-	0,2,2	0.00	-	$_{0,1,1}$	0.00	-	
3	MLI	F	501	-	$0,\!6,\!6$	0.00	-	$0,\!7,\!7$	0.00	-	
4	FMT	F	503	-	0,2,2	0.00	-	$0,\!1,\!1$	0.00	-	
4	FMT	G	303	-	0,2,2	0.00	-	$\overline{0,\!1,\!1}$	0.00	-	
4	FMT	Н	505	-	0,2,2	0.00	-	$0,\!1,\!1$	0.00	-	



Mol	True	Chain	Dec	Tinle	B	ond leng	gths	E	Bond ang	gles
	туре	Chain	nes		Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	FMT	D	502	-	0,2,2	0.00	-	$0,\!1,\!1$	0.00	-
3	MLI	С	301	-	0,6,6	0.00	-	0,7,7	0.00	-
4	FMT	F	502	-	0,2,2	0.00	-	$0,\!1,\!1$	0.00	-
4	FMT	В	504	-	0,2,2	0.00	-	$0,\!1,\!1$	0.00	-
4	FMT	D	504	-	0,2,2	0.00	-	0,1,1	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MLI	А	301	-	-	0/0/4/4	-
3	MLI	С	301	-	-	0/0/4/4	-
3	MLI	Е	301	-	-	0/0/4/4	-
3	MLI	G	301	-	-	0/0/4/4	-
3	MLI	В	501	-	-	0/0/4/4	-
3	MLI	Н	501	-	-	0/0/4/4	-
3	MLI	D	501	-	-	0/0/4/4	-
3	MLI	F	501	-	-	0/0/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

### 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Fit of model and data (i)

## 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	#RSRZ $>$ 2	$\mathbf{OWAB}(\mathbf{A}^2)$	Q<0.9
1	А	249/276~(90%)	-0.17	7 (2%) 53 64	34, 45, 64, 88	0
1	С	248/276~(89%)	0.03	12 (4%) 30 43	30, 50, 81, 101	0
1	Ε	246/276~(89%)	0.78	37 (15%) 2 3	48, 80, 118, 136	0
1	G	247/276~(89%)	-0.31	5 (2%) 65 75	28, 43, 60, 77	0
2	В	404/410~(98%)	-0.44	3 (0%) 87 92	23, 32, 52, 80	0
2	D	397/410~(96%)	-0.34	3 (0%) 86 91	23, 32, 52, 66	0
2	F	398/410~(97%)	-0.34	6 (1%) 73 81	23, 35, 57, 81	0
2	Н	405/410~(98%)	-0.39	7 (1%) 70 78	22, 31, 53, 84	0
All	All	2594/2744~(94%)	-0.20	80 (3%) 49 61	22, 38, 82, 136	0

All (80) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	С	8	GLU	7.1
2	D	22	PRO	4.9
1	Е	197	GLN	4.9
1	А	267	GLY	4.8
1	Е	224	ALA	4.7
2	F	22	PRO	4.5
1	Е	9	ALA	4.4
1	Е	262	ALA	4.3
1	С	9	ALA	4.2
2	Н	408	GLY	4.2
1	Е	21	ALA	3.9
1	Е	23	ASN	3.8
1	С	133	ILE	3.7
1	Е	24	ARG	3.7
2	F	9	THR	3.7
1	Ε	196	SER	3.6



Mol	Chain	Res	Type	RSRZ	
1	Е	222	SER	3.6	
2	Н	22	PRO	3.6	
1	Е	20	ARG	3.5	
1	Е	153	ARG	3.5	
1	Е	208	ALA	3.5	
2	Н	23	SER	3.3	
1	Е	221	ARG	3.3	
1	С	196	SER	3.3	
1	Е	133	ILE	3.1	
1	Е	265	ARG	3.1	
1	А	133	ILE	3.1	
1	Е	218	LEU	3.0	
1	G	133	ILE	3.0	
1	С	197	GLN	3.0	
2	F	21	GLY	2.8	
1	С	23	ASN	2.8	
2	Н	6	ALA	2.8	
1	А	8	GLU	2.7	
1	Е	13	GLY	2.7	
2	В	408	GLY	2.6	
1	А	67	MET	2.6	
1	А	196	SER	2.6	
1	Е	52	GLY	2.6	
1	Е	225	GLN	2.6	
2	Н	8	PRO	2.6	
1	Е	201	GLU	2.6	
1	G	196	SER	2.6	
1	Е	14	PRO	2.6	
1	Е	248	GLU	2.6	
2	В	8	PRO	2.6	
1	G	21	ALA	2.6	
1	Е	211	ASP	2.5	
1	E	231	GLN	2.5	
2	F	407	LEU	2.5	
1	E	198	ALA	2.5	
1	Е	98	ILE	2.5	
1	C	21	ALA	2.5	
1	C	211	ASP	2.5	
1	C	105	VAL	2.4	
2	F	176	GLY	2.4	
1	E	223	ARG	2.4	
1	A	20	ARG	2.4	



Mol	Chain	Res	Type	RSRZ
1	Е	209	VAL	2.4
2	Н	409	ASN	2.4
2	Н	4	ALA	2.3
1	Е	263	GLY	2.3
2	D	277	GLU	2.3
1	С	266	LEU	2.3
1	А	106	MET	2.3
1	Е	132	LEU	2.2
1	Е	251	PRO	2.2
1	Е	228	GLN	2.2
1	G	23	ASN	2.2
1	С	201	GLU	2.2
1	Е	105	VAL	2.2
2	D	10	SER	2.2
1	С	104	VAL	2.2
2	F	177	SER	2.2
1	Е	102	ARG	2.1
1	Е	232	TYR	2.1
1	G	197	GLN	2.1
1	Е	36	PRO	2.1
1	Е	28	ILE	2.1
2	В	5	ILE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	LLP	F	101	24/25	0.98	0.16	24,27,29,29	0
2	LLP	D	101	24/25	0.98	0.14	$23,\!24,\!26,\!27$	0
2	LLP	Н	101	24/25	0.98	0.14	22,24,26,27	0
2	LLP	В	101	24/25	0.99	0.14	25,27,28,29	0

## 6.3 Carbohydrates (i)

There are no carbohydrates in this entry.



## 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathbf{A}^2)$	Q<0.9
4	FMT	Н	503	3/3	0.75	0.19	$56,\!56,\!58,\!58$	0
4	FMT	Н	505	3/3	0.80	0.29	$54,\!54,\!55,\!60$	0
4	FMT	С	302	3/3	0.82	0.22	$52,\!52,\!56,\!57$	0
4	FMT	F	503	3/3	0.83	0.18	70,70,74,74	0
4	FMT	Н	504	3/3	0.84	0.19	$52,\!52,\!54,\!56$	0
4	FMT	Е	302	3/3	0.86	0.15	$59,\!59,\!60,\!61$	0
3	MLI	D	501	7/7	0.86	0.23	57,63,69,69	0
4	FMT	В	503	3/3	0.87	0.13	64,64,67,68	0
4	FMT	G	302	3/3	0.87	0.23	45,45,52,53	0
4	FMT	D	504	3/3	0.87	0.24	62,62,62,65	0
3	MLI	В	501	7/7	0.88	0.18	$53,\!59,\!67,\!68$	0
3	MLI	Е	301	7/7	0.88	0.14	75,79,81,87	0
3	MLI	F	501	7/7	0.88	0.23	58,64,71,74	0
4	FMT	Е	303	3/3	0.89	0.28	80,80,81,82	0
4	FMT	D	502	3/3	0.90	0.36	$65,\!65,\!68,\!68$	0
4	FMT	D	506	3/3	0.91	0.17	$57,\!57,\!58,\!60$	0
4	FMT	Н	502	3/3	0.92	0.16	$53,\!53,\!54,\!54$	0
3	MLI	Н	501	7/7	0.92	0.14	$51,\!54,\!56,\!58$	0
4	FMT	А	302	3/3	0.92	0.22	43,43,46,47	0
3	MLI	А	301	7/7	0.92	0.13	44,47,49,51	0
3	MLI	С	301	7/7	0.93	0.13	$59,\!63,\!65,\!66$	0
4	FMT	F	502	3/3	0.94	0.23	$60,\!60,\!61,\!62$	0
4	FMT	G	303	3/3	0.94	0.15	$53,\!53,\!54,\!54$	0
4	FMT	В	502	3/3	0.95	0.11	47,47,48,49	0
3	MLI	G	301	7/7	0.95	0.10	$51,\!54,\!56,\!56$	0
4	FMT	В	504	3/3	0.95	0.20	61,61,62,62	0
4	FMT	D	503	3/3	0.95	0.18	$50,\!50,\!50,\!51$	0
4	FMT	D	505	3/3	0.96	0.12	$53,\!53,\!53,\!54$	0

## 6.5 Other polymers (i)

There are no such residues in this entry.

