



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 29, 2024 – 01:38 PM EST

PDB ID : 8TC7
Title : Human asparaginyl-tRNA synthetase, apo form
Authors : Dogovski, C.; Metcalfe, R.D.; Xie, S.C.; Morton, C.J.; Tilley, L.; Griffin, M.D.W.
Deposited on : 2023-06-30
Resolution : 1.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	453	
1	B	453	
1	C	453	
1	D	453	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 15123 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Asparagine-tRNA ligase, cytoplasmic.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	426	3471	2215	593	637	26	0	2	0
1	B	429	3488	2224	597	641	26	0	2	0
1	C	428	3464	2210	590	638	26	0	0	0
1	D	428	3461	2207	591	637	26	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	96	GLY	-	expression tag	UNP O43776
A	97	MET	-	expression tag	UNP O43776
B	96	GLY	-	expression tag	UNP O43776
B	97	MET	-	expression tag	UNP O43776
C	96	GLY	-	expression tag	UNP O43776
C	97	MET	-	expression tag	UNP O43776
D	96	GLY	-	expression tag	UNP O43776
D	97	MET	-	expression tag	UNP O43776

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 6 3 3	0	0
2	A	1	Total C O 6 3 3	0	0
2	A	1	Total C O 6 3 3	0	0
2	A	1	Total C O 6 3 3	0	0
2	A	1	Total C O 6 3 3	0	0
2	A	1	Total C O 6 3 3	0	0
2	A	1	Total C O 6 3 3	0	0
2	B	1	Total C O 6 3 3	0	0
2	B	1	Total C O 6 3 3	0	0
2	B	1	Total C O 6 3 3	0	0
2	B	1	Total C O 6 3 3	0	0
2	B	1	Total C O 6 3 3	0	0
2	B	1	Total C O 6 3 3	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total 6	C 3	O 3	0	0
2	B	1	Total 6	C 3	O 3	0	0
2	C	1	Total 6	C 3	O 3	0	0
2	C	1	Total 6	C 3	O 3	0	0
2	C	1	Total 6	C 3	O 3	0	0
2	D	1	Total 6	C 3	O 3	0	0
2	D	1	Total 6	C 3	O 3	0	0
2	D	1	Total 6	C 3	O 3	0	0

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total 1	Cl 1	0	0
3	C	1	Total 1	Cl 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	288	Total 288	O 288	0	0
4	B	329	Total 329	O 329	0	0
4	C	229	Total 229	O 229	0	0
4	D	259	Total 259	O 259	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Asparagine-tRNA ligase, cytoplasmic




- Molecule 1: Asparagine-tRNA ligase, cytoplasmic



- Molecule 1: Asparagine-tRNA ligase, cytoplasmic



- Molecule 1: Asparagine-tRNA ligase, cytoplasmic



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	113.49Å 127.19Å 163.16Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.58 – 1.90 46.58 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.7 (46.58-1.90) 99.8 (46.58-1.90)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.34 (at 1.90Å)	Xtrriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.180 , 0.217 0.187 , 0.221	Depositor DCC
R_{free} test set	9081 reflections (4.90%)	wwPDB-VP
Wilson B-factor (Å ²)	28.9	Xtrriage
Anisotropy	0.660	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 49.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	15123	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 40.40 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.7501e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.64	4/3555 (0.1%)	0.74	0/4810
1	B	0.63	0/3572	0.76	0/4832
1	C	0.57	1/3547 (0.0%)	0.71	0/4800
1	D	0.61	0/3544	0.72	0/4795
All	All	0.61	5/14218 (0.0%)	0.73	0/19237

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	452	CYS	CB-SG	-5.71	1.72	1.81
1	A	483	GLU	CD-OE2	5.29	1.31	1.25
1	C	165	CYS	CB-SG	-5.28	1.73	1.81
1	A	483	GLU	CG-CD	5.22	1.59	1.51
1	A	165	CYS	CB-SG	-5.09	1.73	1.81

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3471	0	3399	21	0
1	B	3488	0	3413	22	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	3464	0	3394	19	0
1	D	3461	0	3390	18	1
2	A	42	0	54	5	1
2	B	54	0	70	2	0
2	C	18	0	23	1	0
2	D	18	0	24	1	0
3	B	1	0	0	1	0
3	C	1	0	0	0	0
4	A	288	0	0	6	0
4	B	329	0	0	10	0
4	C	229	0	0	7	0
4	D	259	0	0	1	0
All	All	15123	0	13767	78	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 78 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:610:CL:CL	4:B:968:HOH:O	2.32	0.84
1:C:483:GLU:OE2	4:C:701:HOH:O	2.00	0.77
1:B:375:HIS:ND1	4:B:705:HOH:O	2.18	0.75
1:B:231:ASN:HB3	2:B:603:GOL:H32	1.70	0.72
1:D:219:LEU:HD13	1:D:221:ASN:HD22	1.58	0.67

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:142:LYS:NZ	2:A:607:GOL:O1[2_455]	2.13	0.07

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	420/453 (93%)	410 (98%)	9 (2%)	1 (0%)	47	38
1	B	423/453 (93%)	414 (98%)	7 (2%)	2 (0%)	29	18
1	C	420/453 (93%)	414 (99%)	5 (1%)	1 (0%)	47	38
1	D	420/453 (93%)	410 (98%)	9 (2%)	1 (0%)	47	38
All	All	1683/1812 (93%)	1648 (98%)	30 (2%)	5 (0%)	41	31

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	324	GLU
1	A	469	VAL
1	C	469	VAL
1	D	469	VAL
1	B	469	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	380/398 (96%)	378 (100%)	2 (0%)	88	89
1	B	381/398 (96%)	375 (98%)	6 (2%)	62	60
1	C	380/398 (96%)	377 (99%)	3 (1%)	81	82
1	D	379/398 (95%)	373 (98%)	6 (2%)	62	60
All	All	1520/1592 (96%)	1503 (99%)	17 (1%)	73	73

5 of 17 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	287	LEU
1	D	404	ASP
1	B	424	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	198	HIS
1	C	287	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	D	143	ASN
1	D	166	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 24 ligands modelled in this entry, 2 are monoatomic - leaving 22 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	B	604	-	5,5,5	1.29	0	5,5,5	1.01	0
2	GOL	C	601	-	5,5,5	0.98	0	5,5,5	0.77	0
2	GOL	B	605	-	5,5,5	0.97	0	5,5,5	0.97	0
2	GOL	A	604	-	5,5,5	0.73	0	5,5,5	1.11	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GOL	B	601	-	5,5,5	1.05	0	5,5,5	1.56	1 (20%)
2	GOL	C	602	-	5,5,5	0.81	0	5,5,5	1.01	0
2	GOL	A	601	-	5,5,5	0.65	0	5,5,5	1.40	1 (20%)
2	GOL	D	601	-	5,5,5	0.85	0	5,5,5	1.56	1 (20%)
2	GOL	A	603	-	5,5,5	1.40	1 (20%)	5,5,5	0.82	0
2	GOL	D	602	-	5,5,5	1.29	1 (20%)	5,5,5	1.32	1 (20%)
2	GOL	D	603	-	5,5,5	1.20	0	5,5,5	0.78	0
2	GOL	B	608	-	5,5,5	1.15	0	5,5,5	0.75	0
2	GOL	B	609	-	5,5,5	1.25	1 (20%)	5,5,5	1.05	0
2	GOL	B	606	-	5,5,5	1.58	1 (20%)	5,5,5	0.94	0
2	GOL	B	607	-	5,5,5	0.64	0	5,5,5	1.09	1 (20%)
2	GOL	A	606	-	5,5,5	2.53	2 (40%)	5,5,5	1.14	0
2	GOL	A	607	-	5,5,5	1.04	0	5,5,5	1.24	0
2	GOL	A	602	-	5,5,5	1.42	1 (20%)	5,5,5	0.83	0
2	GOL	A	605	-	5,5,5	0.99	0	5,5,5	1.01	0
2	GOL	C	603	-	5,5,5	1.79	2 (40%)	5,5,5	2.02	2 (40%)
2	GOL	B	602	-	5,5,5	1.17	1 (20%)	5,5,5	1.46	1 (20%)
2	GOL	B	603	-	5,5,5	1.13	0	5,5,5	1.13	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	B	604	-	-	1/4/4/4	-
2	GOL	C	601	-	-	0/4/4/4	-
2	GOL	B	605	-	-	2/4/4/4	-
2	GOL	A	604	-	-	0/4/4/4	-
2	GOL	B	601	-	-	2/4/4/4	-
2	GOL	C	602	-	-	0/4/4/4	-
2	GOL	A	601	-	-	3/4/4/4	-
2	GOL	D	601	-	-	0/4/4/4	-
2	GOL	A	603	-	-	2/4/4/4	-
2	GOL	D	602	-	-	2/4/4/4	-
2	GOL	D	603	-	-	0/4/4/4	-
2	GOL	B	608	-	-	2/4/4/4	-
2	GOL	B	609	-	-	4/4/4/4	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	B	606	-	-	2/4/4/4	-
2	GOL	B	607	-	-	2/4/4/4	-
2	GOL	A	606	-	-	2/4/4/4	-
2	GOL	A	607	-	-	0/4/4/4	-
2	GOL	A	602	-	-	2/4/4/4	-
2	GOL	A	605	-	-	0/4/4/4	-
2	GOL	C	603	-	-	0/4/4/4	-
2	GOL	B	602	-	-	0/4/4/4	-
2	GOL	B	603	-	-	2/4/4/4	-

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	606	GOL	O3-C3	4.43	1.61	1.42
2	A	606	GOL	C3-C2	3.19	1.64	1.51
2	B	606	GOL	C1-C2	3.15	1.64	1.51
2	C	603	GOL	O3-C3	-2.80	1.30	1.42
2	A	602	GOL	O2-C2	-2.70	1.35	1.43

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	603	GOL	C3-C2-C1	-3.74	97.14	111.70
2	B	602	GOL	C3-C2-C1	-2.51	101.93	111.70
2	D	602	GOL	C3-C2-C1	-2.41	102.33	111.70
2	A	601	GOL	O2-C2-C3	2.27	119.14	109.12
2	B	601	GOL	O1-C1-C2	-2.24	99.47	110.20

There are no chirality outliers.

5 of 28 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	601	GOL	O1-C1-C2-O2
2	A	603	GOL	C1-C2-C3-O3
2	B	601	GOL	O1-C1-C2-O2
2	B	601	GOL	O1-C1-C2-C3
2	B	605	GOL	C1-C2-C3-O3

There are no ring outliers.

8 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	605	GOL	1	0
2	A	601	GOL	1	0
2	D	603	GOL	1	0
2	A	607	GOL	1	1
2	A	602	GOL	2	0
2	A	605	GOL	1	0
2	C	603	GOL	1	0
2	B	603	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	426/453 (94%)	0.14	23 (5%) 25 29	22, 38, 72, 105	0
1	B	429/453 (94%)	-0.04	12 (2%) 53 56	23, 37, 77, 118	0
1	C	428/453 (94%)	0.26	34 (7%) 12 14	24, 43, 89, 117	0
1	D	428/453 (94%)	0.23	35 (8%) 11 13	25, 41, 89, 128	0
All	All	1711/1812 (94%)	0.15	104 (6%) 21 24	22, 39, 81, 128	0

The worst 5 of 104 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	109	LEU	9.5
1	A	109	LEU	8.6
1	A	110	PRO	7.3
1	D	193	GLN	6.5
1	D	191	GLY	6.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	GOL	B	606	6/6	0.55	0.28	60,67,73,74	0
3	CL	C	604	1/1	0.81	0.15	83,83,83,83	0
2	GOL	A	607	6/6	0.82	0.36	65,67,71,73	0
2	GOL	A	606	6/6	0.86	0.18	36,37,46,52	0
2	GOL	A	605	6/6	0.87	0.30	47,65,67,81	0
2	GOL	B	603	6/6	0.87	0.25	63,68,74,78	0
2	GOL	D	601	6/6	0.88	0.24	43,54,57,62	0
2	GOL	C	601	6/6	0.88	0.17	48,53,54,74	0
2	GOL	B	605	6/6	0.90	0.18	59,65,66,74	0
2	GOL	D	602	6/6	0.90	0.12	41,51,61,67	0
2	GOL	C	603	6/6	0.90	0.19	38,41,45,54	0
2	GOL	B	609	6/6	0.91	0.16	35,44,50,60	0
2	GOL	A	601	6/6	0.91	0.12	35,42,44,48	0
2	GOL	B	607	6/6	0.91	0.16	47,54,58,73	0
2	GOL	A	604	6/6	0.93	0.10	47,49,61,64	0
2	GOL	D	603	6/6	0.93	0.13	35,44,51,52	0
2	GOL	B	601	6/6	0.93	0.19	44,52,58,60	0
2	GOL	A	602	6/6	0.94	0.11	45,48,61,62	0
2	GOL	B	608	6/6	0.94	0.22	40,61,65,72	0
2	GOL	B	604	6/6	0.94	0.12	50,58,62,73	0
2	GOL	C	602	6/6	0.95	0.13	47,51,55,57	0
2	GOL	B	602	6/6	0.95	0.10	36,40,45,45	0
2	GOL	A	603	6/6	0.96	0.14	43,49,52,53	0
3	CL	B	610	1/1	0.97	0.21	70,70,70,70	0

6.5 Other polymers [i](#)

There are no such residues in this entry.