

wwPDB X-ray Structure Validation Summary Report (i)

May 25, 2020 – 04:59 pm BST

PDB ID	:	5THQ
Title	:	Comprehensive Analysis of a Novel Ketoreductase for Pentangular Polyphenol
		Biosynthesis
Authors	:	Valentic, T.R.; Tsai, S.C.; Brady, S.F.
Deposited on	:	2016-09-30
$\operatorname{Resolution}$:	2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
buster-report	:	1.1.7(2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
$\operatorname{CCP4}$:	$7.0.044 (\mathrm{Gargrove})$
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries},{ m resolution\ range}({ m \AA}))$
R_{free}	130704	5042(2.30-2.30)
Clashscore	141614	5643(2.30-2.30)
Ramachandran outliers	138981	5575(2.30-2.30)
Sidechain outliers	138945	5575(2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	Δ	272			22/
	A	212	85%	6%	9%
1	В	272	79%	13%	8%
1	С	272	3% 77%	14%	• 8%
1	D	272	% 	10%	• 8%
1	Е	272	78%	14%	8%
1	F	272	76%	15%	• 8%



Continued from previous page...

Mol	Chain	Length	Quality of chain			
1	G	272	% 	11%	•	9%
1	Н	272	% 	13%	•	8%



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 15849 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues		Ate	oms			ZeroOcc	AltConf	Trace
1	G	248	Total	С	Ν	Ο	S	0	0	0
1	G	240	1858	1166	341	349	2	0	0	0
1	В	250	Total	С	Ν	Ο	S	0	0	0
	D	230	1875	1177	343	353	2	0	0	0
1	С	250	Total	С	Ν	Ο	S	0	0	0
	U	230	1875	1177	343	353	2	0	0	0
1	F	250	Total	С	Ν	Ο	S	0	0	0
	Г	230	1875	1177	343	353	2	0	0	0
1	А	248	Total	С	Ν	Ο	S	0	0	0
	А	240	1858	1166	341	349	2	0	0	0
1	Е	250	Total	С	Ν	Ο	S	0	0	0
		230	1875	1177	343	353	2	0	0	0
1	Н	250	Total	С	Ν	Ο	S	0	0	0
	11	230	1875	1177	343	353	2		0	0
1	D	250	Total	С	Ν	Ο	S	0	0	0
		200	1875	1177	343	353	2		U	U

• Molecule 1 is a protein called 3-oxoacyl-ACP reductase.

There are 160 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
G	-19	MET	-	initiating methionine	UNP A0A023PKG5
G	-18	GLY	-	expression tag	UNP A0A023PKG5
G	-17	SER	-	expression tag	UNP A0A023PKG5
G	-16	SER	-	expression tag	UNP A0A023PKG5
G	-15	HIS	-	expression tag	UNP A0A023PKG5
G	-14	HIS	-	expression tag	UNP A0A023PKG5
G	-13	HIS	-	expression tag	UNP A0A023PKG5
G	-12	HIS	-	expression tag	UNP A0A023PKG5
G	-11	HIS	-	expression tag	UNP A0A023PKG5
G	-10	HIS	-	expression tag	UNP A0A023PKG5
G	-9	SER	-	expression tag	UNP A0A023PKG5
G	-8	SER	-	expression tag	UNP A0A023PKG5
G	-7	GLY	_	expression tag	UNP A0A023PKG5



		vious page	A at 1	Comerce eret	Defer
Chain	Residue	Modelled	Actual	Comment	Reference
G	-6	LEU	-	expression tag	UNP A0A023PKG5
G	-5	VAL	-	expression tag	UNP A0A023PKG5
G	-4	PRO	-	expression tag	UNP A0A023PKG5
G	-3	ARG	-	expression tag	UNP A0A023PKG5
G	-2	GLY	-	expression tag	UNP A0A023PKG5
G	-1	SER	-	expression tag	UNP A0A023PKG5
G	0	HIS	-	expression tag	UNP A0A023PKG5
B	-19	MET	-	initiating methionine	UNP A0A023PKG5
B	-18	GLY	_	expression tag	UNP A0A023PKG5
В	-17	SER	_	expression tag	UNP A0A023PKG5
В	-16	SER	-	expression tag	UNP A0A023PKG5
В	-15	HIS	-	expression tag	UNP A0A023PKG5
В	-14	HIS	-	expression tag	UNP A0A023PKG5
В	-13	HIS	-	expression tag	UNP A0A023PKG5
В	-12	HIS	-	expression tag	UNP A0A023PKG5
В	-11	HIS	-	expression tag	UNP A0A023PKG5
В	-10	HIS	-	expression tag	UNP A0A023PKG5
В	-9	SER	-	expression tag	UNP A0A023PKG5
В	-8	SER	-	expression tag	UNP A0A023PKG5
В	-7	GLY	-	expression tag	UNP A0A023PKG5
В	-6	LEU	_	expression tag	UNP A0A023PKG5
В	-5	VAL	-	expression tag	UNP A0A023PKG5
В	-4	PRO	_	expression tag	UNP A0A023PKG5
В	-3	ARG	_	expression tag	UNP A0A023PKG5
В	-2	GLY	_	expression tag	UNP A0A023PKG5
В	-1	SER	-	expression tag	UNP A0A023PKG5
В	0	HIS	-	expression tag	UNP A0A023PKG5
С	-19	MET	-	initiating methionine	UNP A0A023PKG5
С	-18	GLY	-	expression tag	UNP A0A023PKG5
С	-17	SER	-	expression tag	UNP A0A023PKG5
С	-16	SER	-	expression tag	UNP A0A023PKG5
С	-15	HIS	-	expression tag	UNP A0A023PKG5
С	-14	HIS	-	expression tag	UNP A0A023PKG5
С	-13	HIS	-	expression tag	UNP A0A023PKG5
С	-12	HIS	-	expression tag	UNP A0A023PKG5
С	-11	HIS	-	expression tag	UNP A0A023PKG5
С	-10	HIS	-	expression tag	UNP A0A023PKG5
С	-9	SER	-	expression tag	UNP A0A023PKG5
С	-8	SER	-	expression tag	UNP A0A023PKG5
С	-7	GLY	-	expression tag	UNP A0A023PKG5
С	-6	LEU	_	expression tag	UNP A0A023PKG5
С	-5	VAL	_	expression tag	UNP A0A023PKG5
	1	1	1		tinued on nert nage



Continu	Residue	vious page Modelled	Actual	Comment	Reference
С	-4	PRO		expression tag	UNP A0A023PKG5
C		ARG	_	expression tag	UNP A0A023PKG5
<u> </u>	-3	GLY	_	expression tag	UNP A0A023PKG5
<u> </u>	-2	SER	_	expression tag	UNP A0A023PKG5
<u> </u>	0	HIS	_	expression tag	UNP A0A023PKG5
F	-19	MET	_	initiating methionine	UNP A0A023PKG5
F	-13	GLY	_	expression tag	UNP A0A023PKG5
F	-17	SER	_	expression tag	UNP A0A023PKG5
F	-16	SER	_	expression tag	UNP A0A023PKG5
F	-10	HIS	_	expression tag	UNP A0A023PKG5
F	-14	HIS	_	expression tag	UNP A0A023PKG5
F	-14	HIS	_	expression tag	UNP A0A023PKG5
F	-13	HIS	_	expression tag	UNP A0A023PKG5
F	-12	HIS	_	expression tag	UNP A0A023PKG5
F	-11	HIS	_	expression tag	UNP A0A023PKG5
F	-9	SER	_	expression tag	UNP A0A023PKG5
F	-3	SER	_	expression tag	UNP A0A023PKG5
 F	-7	GLY	_	expression tag	UNP A0A023PKG5
F	-6	LEU	_	expression tag	UNP A0A023PKG5
F	-0	VAL	_	expression tag	UNP A0A023PKG5
F		PRO	_	expression tag	UNP A0A023PKG5
F		ARG	_	expression tag	UNP A0A023PKG5
F	-3	GLY	_	expression tag	UNP A0A023PKG5
F	-2	SER	_	expression tag	UNP A0A023PKG5
F	0	HIS	_	expression tag	UNP A0A023PKG5
A	-19	MET	_	initiating methionine	UNP A0A023PKG5
A	-18	GLY	_	expression tag	UNP A0A023PKG5
A	-17	SER	-	expression tag	UNP A0A023PKG5
A	-16	SER		expression tag	UNP A0A023PKG5
A	-15	HIS	_	expression tag	UNP A0A023PKG5
A	-14	HIS		expression tag	UNP A0A023PKG5
A	-13	HIS	_	expression tag	UNP A0A023PKG5
A	-12	HIS	_	expression tag	UNP A0A023PKG5
A	-11	HIS	_	expression tag	UNP A0A023PKG5
A	-10	HIS	_	expression tag	UNP A0A023PKG5
A	-9	SER	_	expression tag	UNP A0A023PKG5
A	-8	SER	_	expression tag	UNP A0A023PKG5
A	-7	GLY	-	expression tag	UNP A0A023PKG5
A	-6	LEU	_	expression tag	UNP A0A023PKG5
A	-5	VAL	_	expression tag	UNP A0A023PKG5
A	-4	PRO	_	expression tag	UNP A0A023PKG5
A	-3	ARG	_	expression tag	UNP A0A023PKG5
11		11100			tinued on nert nage




Continu Chain	Residue	vious page Modelled	Actual	Commont	Reference
				Comment	
<u>A</u>	-2	GLY	-	expression tag	UNP A0A023PKG5
A	-1	SER	-	expression tag	UNP A0A023PKG5
A	0	HIS	-	expression tag	UNP A0A023PKG5
E	-19	MET	-	initiating methionine	UNP A0A023PKG5
Е	-18	GLY	-	expression tag	UNP A0A023PKG5
Е	-17	SER	_	expression tag	UNP A0A023PKG5
Е	-16	SER	-	expression tag	UNP A0A023PKG5
Е	-15	HIS	-	expression tag	UNP A0A023PKG5
E	-14	HIS	-	expression tag	UNP A0A023PKG5
Ε	-13	HIS	-	expression tag	UNP A0A023PKG5
Ε	-12	HIS	-	expression tag	UNP A0A023PKG5
Ε	-11	HIS	-	expression tag	UNP A0A023PKG5
Е	-10	HIS	-	expression tag	UNP A0A023PKG5
Ε	-9	SER	-	expression tag	UNP A0A023PKG5
Е	-8	SER	_	expression tag	UNP A0A023PKG5
Е	-7	GLY	_	expression tag	UNP A0A023PKG5
Е	-6	LEU	_	expression tag	UNP A0A023PKG5
Е	-5	VAL	-	expression tag	UNP A0A023PKG5
Е	-4	PRO	-	expression tag	UNP A0A023PKG5
Е	-3	ARG	-	expression tag	UNP A0A023PKG5
Е	-2	GLY	-	expression tag	UNP A0A023PKG5
Е	-1	SER	-	expression tag	UNP A0A023PKG5
Е	0	HIS	-	expression tag	UNP A0A023PKG5
Н	-19	MET	-	initiating methionine	UNP A0A023PKG5
Н	-18	GLY	-	expression tag	UNP A0A023PKG5
Н	-17	SER	-	expression tag	UNP A0A023PKG5
Н	-16	SER	-	expression tag	UNP A0A023PKG5
Н	-15	HIS	-	expression tag	UNP A0A023PKG5
Н	-14	HIS	-	expression tag	UNP A0A023PKG5
Н	-13	HIS	-	expression tag	UNP A0A023PKG5
Н	-12	HIS	-	expression tag	UNP A0A023PKG5
Н	-11	HIS	-	expression tag	UNP A0A023PKG5
Н	-10	HIS	-	expression tag	UNP A0A023PKG5
Н	-9	SER	-	expression tag	UNP A0A023PKG5
Н	-8	SER	-	expression tag	UNP A0A023PKG5
Η	-7	GLY	-	expression tag	UNP A0A023PKG5
Н	-6	LEU	_	expression tag	UNP A0A023PKG5
Н	-5	VAL	_	expression tag	UNP A0A023PKG5
Н	-4	PRO	_	expression tag	UNP A0A023PKG5
Н	-3	ARG	-	expression tag	UNP A0A023PKG5
H	-2	GLY	-	expression tag	UNP A0A023PKG5
H	-1	SER	_	expression tag	UNP A0A023PKG5
11				1 0	tinued on next nage



5THQ	
------	--

Chain	Residue	Modelled	Actual	Comment	Reference
Н	0	HIS	-	expression tag	UNP A0A023PKG5
D	-19	MET	-	initiating methionine	UNP A0A023PKG5
D	-18	GLY	-	expression tag	UNP A0A023PKG5
D	-17	SER	-	expression tag	UNP A0A023PKG5
D	-16	SER	-	expression tag	UNP A0A023PKG5
D	-15	HIS	-	expression tag	UNP A0A023PKG5
D	-14	HIS	-	expression tag	UNP A0A023PKG5
D	-13	HIS	-	expression tag	UNP A0A023PKG5
D	-12	HIS	-	expression tag	UNP A0A023PKG5
D	-11	HIS	-	expression tag	UNP A0A023PKG5
D	-10	HIS	-	expression tag	UNP A0A023PKG5
D	-9	SER	_	expression tag	UNP A0A023PKG5
D	-8	SER	-	expression tag	UNP A0A023PKG5
D	-7	GLY	-	expression tag	UNP A0A023PKG5
D	-6	LEU	-	expression tag	UNP A0A023PKG5
D	-5	VAL	-	expression tag	UNP A0A023PKG5
D	-4	PRO	-	expression tag	UNP A0A023PKG5
D	-3	ARG	-	expression tag	UNP A0A023PKG5
D	-2	GLY	-	expression tag	UNP A0A023PKG5
D	-1	SER	-	expression tag	UNP A0A023PKG5
D	0	HIS	-	expression tag	UNP A0A023PKG5

• Molecule 2 is NADPH DIHYDRO-NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NDP) (formula: C₂₁H₃₀N₇O₁₇P₃).





rп		`
5	ΓH(.)
<u> </u>		с.

Mol	Chain	Residues		Ate	oms			ZeroOcc	AltConf	
2	G	1	Total	С	Ν	Ο	Р	0	0	
	G	L	48	21	7	17	3	0	0	
2	В	1	Total	С	Ν	Ο	Р	0	0	
	D	L	48	21	7	17	3	0	0	
2	С	1	Total	С	Ν	Ο	Р	0	0	
	U	L	48	21	7	17	3	0	0	
2	F	1	Total	С	Ν	Ο	Р	0	0	
	Ľ	I	48	21	7	17	3	0		
2	А	1	Total	С	Ν	Ο	Р	0	0	
	Л	I	48	21	7	17	3		0	
2	Е	1	Total	С	Ν	Ο	Р	0	0	
		L	48	21	7	17	3	0	0	
2	Н	1	Total	С	Ν	Ο	Р	0	0	
	11	L	48	21	7	17	3	0	U	
2	D	1	Total	С	Ν	Ο	Р	0	0	
	D		48	21	7	17	3	0		


• Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	G	54	$\begin{array}{cc} \text{Total} & \text{O} \\ 54 & 54 \end{array}$	0	0
3	В	60	Total O 60 60	0	0
3	С	36	Total O 36 36	0	0
3	F	49	Total O 49 49	0	0
3	А	80	Total O 80 80	0	0
3	Ε	65	$\begin{array}{cc} {\rm Total} & {\rm O} \\ 65 & 65 \end{array}$	0	0
3	Н	77	Total O 77 77	0	0
3	D	78	Total O 78 78	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: 3-oxoacyl-ACP reductase



MET MET SER SER SER RIS RIS RIS RIS RIS RIS RIS RIS RIS RI	115 616 737 737 737 737 737 734 746 746 746 746 753 746 753 753 753 753 754 755 754 755 755 754 755 754 755 755	168 168 197 197 198 198 198 198 100 1112 1114 1114
1126 1130 1130 1130 1156 1156 1156 1156 1160 1160 1171 1160 1185 1185 1185	H196 11199 11199 1200 1200 1200 1200 1200 12	
• Molecule 1: 3-oxoacyl-ACP redu	ctase	
Chain A:	85%	6% 9%
MET OLY SER SER SER HIS HIS HIS HIS SER HIS SER VIL VIL VIL VIL VIL VIL VIL VIL VIL VIL	198 0112 0112 0113 0134 1151 1151 1151 1165 1165 1165 1165 116	1198 1198 1200 1200 1214 1214 1245 1245 1245
• Molecule 1: 3-oxoacyl-ACP redu	ctase	
Chain E:	78%	14% 8%
MET SER SER SER HIS SER HIS SER RIS SER SER SER SER SER SER SER SER SER SE	115 115 115 115 126 128 128 128 128 128 128 128 128	K72 E76 094 1126 1126 1126 1128 1128 1128
P131 K145 K145 1146 P149 P160 P161 P163 P163 P163 P163 P163 P195 P196 P196 P196 P196 P196 P196 P196 P196	M206 1214 2234 1245 1245 8252	
• Molecule 1: 3-oxoacyl-ACP redu	ctase	
Chain H:	78%	13% • 8%
MET MET SER SER SER HIS HIS HIS SER RIS SER RIS SER RIS CLY CLU CLU CLU CLU CLU CLU CLU	115 115 122 122 123 123 124 126 145 145 145 145 145 145 145 145 145 145	E76 1883 1994 11133 11133 11151
R154 K161 H162 H163 A163 A163 A163 A163 A163 A163 A163 A	S234 1245 S252	
• Molecule 1: 3-oxoacyl-ACP redu	ctase	
Chain D:	81%	10% • 8%
MET SER SER SER HIS HIS HIS SER PRO CLV CLV CLV SER SER SER SER SER SER SER SER SER SER	K11 66 966 785 785 7112 7145 7145 7154 7154 7154 7154	4198 8191 1198 1198 1198 1198 1198 1200 1200 1200 1200
A204 K208 T212 K213 T214 S234 L236 L236 L236 L236 L245 L245 S262		



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
$\begin{array}{c} \text{Cell constants} \\ \text{a, b, c, } \alpha, \beta, \gamma \end{array}$	$\frac{105.34 \text{\AA}}{90.00^{\circ}} \frac{86.54 \text{\AA}}{106.97 \text{\AA}} \frac{106.97 \text{\AA}}{90.00^{\circ}}$	Depositor
Resolution (Å)	$\begin{array}{rrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrr$	Depositor EDS
% Data completeness	$98.3 \ (86.31 - 2.30)$	Depositor
(in resolution range)	94.2 (86.30-2.30)	EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$3.65 \;({ m at}\; 2.29{ m \AA})$	Xtriage
Refinement program	PHENIX (1.10_2155: ???)	Depositor
R, R_{free}	$\begin{array}{rrrr} 0.169 & , & 0.217 \\ 0.169 & , & 0.217 \end{array}$	Depositor DCC
R_{free} test set	1993 reflections (2.50%)	wwPDB-VP
Wilson B-factor $(Å^2)$	26.6	Xtriage
Anisotropy	0.418	Xtriage
Bulk solvent $k_{sol}(e/A^3)$, $B_{sol}(A^2)$	0.33 , 40.2	EDS
L-test for twinning ²	$< L > = 0.49, < L^2 > = 0.32$	Xtriage
Estimated twinning fraction	0.019 for l,-k,h	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	15849	wwPDB-VP
Average B, all atoms $(Å^2)$	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 24.56 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.7459e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NDP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Chain		Bo	nd lengths	Bond angles		
			# Z > 5	RMSZ	# Z > 5	
1	А	0.48	0/1888	0.74	2/2558~(0.1%)	
1	В	0.48	0/1905	0.76	1/2581~(0.0%)	
1	С	0.44	0/1905	0.69	1/2581~(0.0%)	
1	D	0.51	0/1905	0.70	0/2581	
1	Ε	0.53	1/1905~(0.1%)	0.72	0/2581	
1	F	0.50	0/1905	0.71	0/2581	
1	G	0.53	0/1888	0.77	4/2558~(0.2%)	
1	Н	0.66	4/1905~(0.2%)	0.72	0/2581	
All	All	0.52	5/15206~(0.0%)	0.73	8/20602~(0.0%)	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	С	0	2
1	D	0	1
1	F	0	2
All	All	0	5

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(Å)
1	Н	203	ARG	CZ-NH2	-10.45	1.19	1.33
1	Н	203	ARG	NE-CZ	-9.78	1.20	1.33
1	Н	203	ARG	CD-NE	-7.60	1.33	1.46
1	Н	203	ARG	CZ-NH1	-7.55	1.23	1.33
1	Е	194	GLU	CD-OE2	5.30	1.31	1.25

The worst 5 of 8 bond angle outliers are listed below:



Mol	Chain	\mathbf{Res}	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
1	В	8	ARG	NE-CZ-NH2	-8.84	115.88	120.30
1	G	84	ARG	NE-CZ-NH1	8.78	124.69	120.30
1	А	169	ARG	NE-CZ-NH2	-7.80	116.40	120.30
1	G	65	ARG	CG-CD-NE	-7.26	96.55	111.80
1	G	84	ARG	NE-CZ-NH2	-6.98	116.81	120.30

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	С	194	GLU	Sidechain
1	С	201	ASP	Peptide
1	D	199	PRO	Peptide
1	F	202	GLU	Peptide
1	F	204	ALA	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	1858	0	1886	14	0
1	В	1875	0	1903	31	0
1	С	1875	0	1903	41	0
1	D	1875	0	1903	27	0
1	Е	1875	0	1903	31	1
1	F	1875	0	1903	51	0
1	G	1858	0	1886	33	0
1	Н	1875	0	1903	24	1
2	А	48	0	25	1	0
2	В	48	0	24	2	0
2	С	48	0	25	2	0
2	D	48	0	25	1	0
2	Е	48	0	25	6	0
2	F	48	0	25	9	0
2	G	48	0	25	2	0
2	Н	48	0	26	3	0
3	А	80	0	0	2	0
3	В	60	0	0	12	0



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes				
3	С	36	0	0	5	0				
3	D	78	0	0	6	0				
3	Ε	65	0	0	3	0				
3	F	49	0	0	4	0				
3	G	54	0	0	3	0				
3	Н	77	0	0	6	0				
All	All	15849	0	15390	242	1				

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 242 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:49:SER:HB3	3:F:402:HOH:O	1.25	1.30
1:B:53:GLU:HB2	3:B:401:HOH:O	1.29	1.30
1:D:3:LEU:N	3:D:401:HOH:O	1.85	1.09
1:G:151:ILE:HB	1:G:154:ARG:NH1	1.68	1.08
1:C:58:GLY:C	3:C:401:HOH:O	1.93	1.05

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:45:ASP:OD2	1:H:83:ARG:NH1[2_554]	2.14	0.06

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured Allowed		Outliers	Percentiles	
1	А	246/272~(90%)	239~(97%)	6(2%)	1 (0%)	34 42	



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	В	248/272~(91%)	241~(97%)	7 (3%)	0	100	100
1	С	248/272~(91%)	244 (98%)	4 (2%)	0	100	100
1	D	248/272~(91%)	240~(97%)	8 (3%)	0	100	100
1	Ε	248/272~(91%)	243~(98%)	4 (2%)	1 (0%)	34	42
1	F	248/272~(91%)	241 (97%)	7 (3%)	0	100	100
1	G	246/272~(90%)	242~(98%)	3 (1%)	1 (0%)	34	42
1	Н	248/272~(91%)	242 (98%)	6 (2%)	0	100	100
All	All	1980/2176~(91%)	1932~(98%)	45 (2%)	3 (0%)	47	58

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	Е	200	ASP
1	А	200	ASP
1	G	200	ASP

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	191/212~(90%)	191~(100%)	0	100 100
1	В	193/212~(91%)	192~(100%)	1 (0%)	88 95
1	С	193/212~(91%)	191~(99%)	2(1%)	76 87
1	D	193/212~(91%)	191~(99%)	2(1%)	76 87
1	Ε	193/212~(91%)	191~(99%)	2(1%)	76 87
1	F	193/212~(91%)	191~(99%)	2(1%)	76 87
1	G	191/212~(90%)	188~(98%)	3~(2%)	62 78
1	Н	193/212~(91%)	188~(97%)	5(3%)	46 63
All	All	1540/1696~(91%)	1523~(99%)	17 (1%)	73 86

5 of 17 residues with a non-rotameric side chain are listed below:



Mol	Chain	Res	Type
1	F	196	HIS
1	Е	201	ASP
1	Н	180	LEU
1	F	3	LEU
1	Н	203	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	В	100	HIS
1	С	100	HIS
1	А	79	GLN
1	D	100	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	B	ond leng	gths	B	ond ang	les
	туре	Cham	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NDP	F	301	-	45,52,52	4.06	16 (35%)	53,80,80	2.42	7 (13%)



Mol	Tune	Chain	Res	Link	B	ond leng	gths	B	ond ang	les
	Type	Unam	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NDP	D	301	-	45,52,52	4.09	18 (40%)	53,80,80	2.12	5 (9%)
2	NDP	В	301	-	45,52,52	4.09	20 (44%)	53,80,80	2.00	5 (9%)
2	NDP	G	301	-	45,52,52	4.02	19 (42%)	53,80,80	2.13	<mark>5 (9%)</mark>
2	NDP	Н	301	-	45,52,52	<u>3.91</u>	19 (42%)	53,80,80	2.10	<mark>6 (11%)</mark>
2	NDP	Е	301	-	45,52,52	<mark>3.99</mark>	17 (37%)	53,80,80	2.49	<mark>6 (11%)</mark>
2	NDP	С	301	-	45,52,52	<mark>3.95</mark>	20 (44%)	53,80,80	2.20	<mark>6 (11%)</mark>
2	NDP	А	301	-	45,52,52	3.94	20 (44%)	53,80,80	1.97	<mark>6 (11%)</mark>

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NDP	F	301	-	-	7/30/77/77	0/5/5/5
2	NDP	D	301	-	-	4/30/77/77	0/5/5/5
2	NDP	В	301	-	-	6/30/77/77	0/5/5/5
2	NDP	G	301	-	-	4/30/77/77	0/5/5/5
2	NDP	Н	301	-	-	10/30/77/77	0/5/5/5
2	NDP	Е	301	-	-	9/30/77/77	0/5/5/5
2	NDP	С	301	-	-	9/30/77/77	0/5/5/5
2	NDP	А	301	-	-	7/30/77/77	0/5/5/5

The worst 5 of 149 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(Å)
2	F	301	NDP	O4B-C1B	15.54	1.62	1.41
2	В	301	NDP	O4B-C1B	15.32	1.62	1.41
2	D	301	NDP	O4B-C1B	15.19	1.62	1.41
2	Е	301	NDP	O4B-C1B	15.16	1.62	1.41
2	G	301	NDP	O4B-C1B	15.08	1.62	1.41

The worst 5 of 46 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
2	F	301	NDP	C5A-C6A-N6A	11.18	137.35	120.35
2	Е	301	NDP	C5A-C6A-N6A	10.97	137.02	120.35
2	С	301	NDP	C5A-C6A-N6A	9.37	134.59	120.35
2	G	301	NDP	C5A-C6A-N6A	9.27	134.44	120.35



Continued from previous page...

Mol	Chain	\mathbf{Res}	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	D	301	NDP	C5A-C6A-N6A	8.93	133.92	120.35

There are no chirality outliers.

5 of 56 torsion outliers are listed below:

Mol	Chain	\mathbf{Res}	Type	Atoms
2	F	301	NDP	C5D-O5D-PN-O3
2	F	301	NDP	C5D-O5D-PN-O2N
2	В	301	NDP	C2B-O2B-P2B-O1X
2	G	301	NDP	C2B-O2B-P2B-O1X
2	Е	301	NDP	C2B-O2B-P2B-O1X


There are no ring outliers.

8	monomers	are	invol	ved	in	26	short	contacts:
---	----------	----------------------	-------	-----	----	----	------------------------	-----------


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	F	301	NDP	9	0
2	D	301	NDP	1	0
2	В	301	NDP	2	0
2	G	301	NDP	2	0
2	Н	301	NDP	3	0
2	Е	301	NDP	6	0
2	С	301	NDP	2	0
2	A	301	NDP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







































5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$<$ RSRZ $>$	$\# RSRZ {>}2$	$\mathbf{OWAB}(\mathbf{\AA}^2)$	Q<0.9
1	А	248/272~(91%)	-0.23	2 (0%) 86 89	20, 29, 51, 78	0
1	В	250/272~(91%)	-0.15	4 (1%) 72 77	21, 32, 61, 94	0
1	С	250/272~(91%)	0.07	8 (3%) 47 54	24, 38, 69, 92	0
1	D	250/272~(91%)	-0.21	2 (0%) 86 89	19, 29, 55, 86	0
1	Ε	250/272~(91%)	-0.13	7 (2%) 53 60	19, 29, 59, 94	0
1	F	250/272~(91%)	0.02	9 (3%) 42 49	22, 36, 70, 101	0
1	G	248/272~(91%)	-0.17	4 (1%) 72 77	21,33,56,89	0
1	Η	250/272~(91%)	-0.18	3 (1%) 79 83	19, 28, 55, 89	0
All	All	1996/2176~(91%)	-0.12	39 (1%) 65 71	19,31,63,101	0

The worst 5 of 39 RSRZ outliers are listed below:

Mol	Chain	\mathbf{Res}	Type	RSRZ
1	С	252	SER	6.0
1	F	198	LEU	4.6
1	F	201	ASP	4.6
1	Н	252	SER	4.0
1	D	199	PRO	3.9

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.




6.4 Ligands (i)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathbf{A}^2)$	$Q{<}0.9$
2	NDP	F	301	48/48	0.90	0.21	$39,\!48,\!57,\!58$	0
2	NDP	Е	301	48/48	0.94	0.13	$24,\!36,\!43,\!47$	0
2	NDP	В	301	48/48	0.95	0.13	$31,\!38,\!48,\!49$	0
2	NDP	D	301	48/48	0.95	0.14	$29,\!38,\!43,\!45$	0
2	NDP	С	301	48/48	0.95	0.16	$33,\!48,\!57,\!63$	0
2	NDP	G	301	48/48	0.96	0.13	$26,\!38,\!44,\!49$	0
2	NDP	Н	301	48/48	0.97	0.13	$27,\!33,\!39,\!41$	0
2	NDP	А	301	48/48	0.97	0.12	$26,\!31,\!37,\!39$	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





















6.5 Other polymers (i)

There are no such residues in this entry.

