

Full wwPDB X-ray Structure Validation Report (i)

Jul 31, 2023 – 12:46 AM EDT

PDB ID : 1TJS

Title : E. COLI THYMIDYLATE SYNTHASE Authors : Stout, T.J.; Sage, C.R.; Stroud, R.M.

Deposited on : 1997-06-27

Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul: 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : NOT EXECUTED EDS : NOT EXECUTED

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.34

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# \mathrm{Entries})$	$(\# ext{Entries}, ext{ resolution range}(ext{Å}))$
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain		
1	A	265	75%	19%	5%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	PO4	A	265	-	X	-	-



2 Entry composition (i)


There are 3 unique types of molecules in this entry. The entry contains 2257 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called THYMIDYLATE SYNTHASE.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	A	265	Total 2153	C 1375	N 371	O 395	S 12	0	0	0

• Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	
2	A	1	Total 5	O 4	P 1	0	0

• Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	99	Total O 99 99	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: THYMIDYLATE SYNTHASE





4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source	
Space group	I 21 3	Depositor	
Cell constants	132.94Å 132.94Å 132.94Å	Depositor	
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor	
Resolution (Å)	8.00 - 2.20	Depositor	
% Data completeness	91.4 (8.00-2.20)	Depositor	
(in resolution range)	31.4 (0.00 2.20)	Берозног	
R_{merge}	0.06	Depositor	
R_{sym}	0.06	Depositor	
Refinement program	X-PLOR 3.843	Depositor	
R, R_{free}	0.191 , 0.253	Depositor	
Estimated twinning fraction	No twinning to report.	Xtriage	
Total number of atoms	2257	wwPDB-VP	
Average B, all atoms (Å ²)	24.0	wwPDB-VP	



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, FMT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Chain		Bond	$\mathbf{lengths}$	Bond angles		
MIOI	Mol Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.38	0/2210	0.71	0/3000	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2153	0	2080	45	0
2	A	5	0	0	1	0
3	A	99	0	0	3	0
All	All	2257	0	2080	45	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	1100111 1		$egin{aligned} \operatorname{Clash} \ \operatorname{overlap}\ (\begin{subarray}{c} \begin{subarray}{c} \begi$
1:A:215:GLN:HE21	1:A:215:GLN:H	1.14	0.91

Continued on next page...



 $Continued\ from\ previous\ page...$

Continued from prev		Interatomic	Clash
Atom-1	Atom-2	${\rm distance}\ (\mathring{\rm A})$	overlap (Å)
1:A:61:TRP:CD1	1:A:66:ASP:HB3	2.33	0.64
1:A:145:PRO:O	1:A:166:ARG:HD3	1.99	0.63
1:A:215:GLN:H	1:A:215:GLN:NE2	1.93	0.62
1:A:68:ASN:HD22	1:A:70:ALA:H	1.47	0.62
1:A:112:ILE:HD11	1:A:240:PHE:CE1	2.35	0.61
1:A:212:HIS:HE1	1:A:261:PRO:O	1.84	0.59
1:A:147:HIS:HB2	1:A:163:LEU:HD21	1.84	0.59
1:A:45:VAL:CG1	1:A:50:CYS:SG	2.92	0.57
1:A:231:ILE:HG23	1:A:248:GLU:HB2	1.87	0.57
1:A:69:ILE:HD11	1:A:90:LEU:HD11	1.86	0.56
1:A:60:LEU:O	1:A:64:GLN:HG2	2.06	0.55
1:A:61:TRP:NE1	1:A:66:ASP:HB3	2.23	0.53
1:A:215:GLN:HE21	1:A:215:GLN:N	1.96	0.53
1:A:245:GLU:H	1:A:245:GLU:CD	2.13	0.52
1:A:190:GLN:NE2	1:A:242:TYR:OH	2.42	0.52
1:A:130:VAL:HG22	1:A:150:PHE:CE1	2.45	0.52
1:A:68:ASN:ND2	1:A:70:ALA:H	2.09	0.49
1:A:37:ASN:ND2	1:A:39:GLN:H	2.10	0.49
1:A:115:VAL:HG21	1:A:130:VAL:HG12	1.96	0.48
1:A:23:GLY:HA3	3:A:364:HOH:O	2.14	0.47
1:A:134:ASN:C	1:A:134:ASN:HD22	2.18	0.46
1:A:171:PHE:CZ	1:A:215:GLN:HB3	2.51	0.46
1:A:212:HIS:HD2	3:A:330:HOH:O	1.99	0.46
1:A:45:VAL:HG13	1:A:50:CYS:SG	2.55	0.46
1:A:166:ARG:NH2	2:A:265:PO4:O3	2.48	0.46
1:A:2:LYS:O	1:A:6:GLU:HG3	2.16	0.45
1:A:0:FMT:C	1:A:46:THR:H	2.30	0.45
1:A:101:TRP:HB3	1:A:109:ILE:HB	1.98	0.45
1:A:212:HIS:O	1:A:216:THR:HG23	2.17	0.45
1:A:212:HIS:CE1	1:A:260:ALA:HB1	2.52	0.44
1:A:140:LYS:HB2	1:A:140:LYS:HE3	1.78	0.44
1:A:47:THR:HA	1:A:255:HIS:CD2	2.54	0.43
1:A:255:HIS:HB3	1:A:256:PRO:HD2	2.01	0.43
1:A:27:LEU:HD23	1:A:27:LEU:HA	1.84	0.42
1:A:181:TYR:O	1:A:185:VAL:HG23	2.20	0.42
1:A:7:LEU:HG	1:A:32:HIS:CE1	2.55	0.41
1:A:264:ILE:HD12	1:A:264:ILE:HA	1.89	0.41
1:A:231:ILE:CG2	1:A:248:GLU:HB2	2.51	0.41
1:A:233:LYS:HG3	1:A:246:ASP:O	2.20	0.41
1:A:239:ILE:O	1:A:239:ILE:HG13	2.20	0.41
1:A:37:ASN:C	1:A:37:ASN:HD22	2.24	0.41

Continued on next page...



Continued from previous page...

Atom-1	Atom-2	$egin{aligned} & ext{Interatomic} \ & ext{distance} \ & ext{(Å)} \end{aligned}$	$egin{aligned} ext{Clash} \ ext{overlap } (ext{Å}) \end{aligned}$
1:A:21:ARG:HD3	3:A:414:HOH:O	2.20	0.41
1:A:208:LEU:HD22	1:A:216:THR:HG21	2.03	0.41
1:A:67:THR:HG23	1:A:67:THR:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	263/265 (99%)	252 (96%)	9 (3%)	2 (1%)	19 19

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	21	ARG
1	A	94	TYR

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mo	Chain	Analysed	Analysed Rotameric		Percentiles	
1	A	233/233 (100%)	210 (90%)	23 (10%)	8 7	

All (23) residues with a non-rotameric sidechain are listed below:



Mol	Chain	Res	Type
1	A	5	LEU
	A		
1	A	7	LEU
1	A	21	ARG
1	A	37	ASN
1	A	39	GLN
1	A	60	LEU
1	A	66	ASP
1	A	67	THR
1	A	76	ASN
1	A A	112	ILE
1	A	130	VAL
1	A A	134	ASN
1	A	143	LEU
1	A	163	LEU
1	A	166	ARG
1	A A	191	GLN
1	A	215	GLN
1	A	220	LEU
1	A	222	ARG
1	A A A	225	ARG
1	A	233	LYS
1	A	245	GLU
1	A	249	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	32	HIS
1	A	37	ASN
1	A	39	GLN
1	A	68	ASN
1	A	76	ASN
1	A	117	ASN
1	A	134	ASN
1	A	147	HIS
1	A	190	GLN
1	A	212	HIS
1	A	215	GLN
1	A	219	GLN



5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains i

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res Link		В	ond len	gths	В	ond ang	gles
WIOI	Туре	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z >2
2	PO4	A	265	-	4,4,4	3.12	4 (100%)	6,6,6	1.83	2 (33%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	Observed(A)	$\operatorname{Ideal}(\text{\AA})$
2	A	265	PO4	P-O1	4.75	1.62	1.50
2	A	265	PO4	P-O3	-2.54	1.47	1.54
2	A	265	PO4	P-O4	-2.29	1.47	1.54
2	A	265	PO4	P-O2	-2.17	1.48	1.54

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^o)$
2	A	265	PO4	O4-P-O3	2.35	115.51	107.97
2	A	265	PO4	O4-P-O1	-2.16	103.01	110.89



There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	265	PO4	1	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

