Full wwPDB X-ray Structure Validation Report ⓘ

Sep 23, 2023 – 02:56 PM EDT

PDB ID	:	5TJ8
Title	:	Structure of WWP2 WW2-2,3-linker-HECT (no WW2 observed)
Authors	:	Chen, Z.; Gabelli, S.B.
Deposited on	:	2016-10-03
Resolution	:	2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references ⓘ) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.35.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	447	4% 67% 19% 5% 9%

2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 3510 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NEDD4-like E3 ubiquitin-protein ligase WWP2,NEDD4-like E3 ubiquitin-protein ligase WWP2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	408	3457	2225	588	622	22	0	1	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	333	GLY	-	expression tag	UNP O00308

- Molecule 2 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Na		
2	A	1	1	1	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	52	52	52	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: NEDD4-like E3 ubiquitin-protein ligase WWP2,NEDD4-like E3 ubiquitin-protein ligase WWP2

Chain A:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α, β, γ	44.23Å 89.27Å 111.47Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.30 47.28 – 2.20	Depositor EDS
% Data completeness (in resolution range)	95.5 (50.00-2.30) 92.5 (47.28-2.20)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	3.02 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
R, R_{free}	0.215 , 0.316 0.238 , 0.303	Depositor DCC
R_{free} test set	1072 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å^2)	42.3	Xtriage
Anisotropy	1.084	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.30 , 46.8	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.94	EDS
Total number of atoms	3510	wwPDB-VP
Average B, all atoms (Å^2)	67.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.58% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| > 5$	RMSZ	$\#\|Z\| > 5$
1	A	0.67	0/3550	0.84	4/4785 (0.1%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	381	GLY	C-N-CA	7.14	139.56	121.70
1	A	575	MET	N-CA-C	-6.70	92.91	111.00
1	A	700	ASN	N-CA-C	-5.12	97.19	111.00
1	A	360	ARG	C-N-CD	5.10	139.11	128.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3457	0	3343	101	0
2	A	1	0	0	0	0
3	A	52	0	0	0	0
All	All	3510	0	3343	101	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

All (101) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:655:ILE:CD1	1:A:708:LEU:HD13	1.77	1.15
1:A:531:GLN:O	1:A:535:MET:HG3	1.54	1.08
1:A:655:ILE:HD11	1:A:708:LEU:HD13	1.35	1.03
1:A:828:LYS:O	1:A:830:THR:N	1.91	1.03
1:A:657:TRP:HA	1:A:657:TRP:CE3	1.98	0.98
1:A:828:LYS:HB2	1:A:831:TRP:HD1	1.27	0.97
1:A:379:LEU:HD12	1:A:380:GLN:N	1.81	0.96
1:A:677:GLU:N	1:A:677:GLU:OE2	2.00	0.94
1:A:573:ASN:O	1:A:575:MET:O	1.86	0.93
1:A:655:ILE:CD1	1:A:708:LEU:CD1	2.46	0.92
1:A:657:TRP:HA	1:A:657:TRP:HE3	1.37	0.90
1:A:825:LYS:O	1:A:826:VAL:HG23	1.70	0.90
1:A:376:ARG:HA	1:A:379:LEU:HG	1.54	0.90
1:A:674:GLN:NE2	1:A:674:GLN:HA	1.87	0.88
1:A:828:LYS:HB2	1:A:831:TRP:CD1	2.09	0.87
1:A:671:TYR:O	1:A:673:ILE:N	2.08	0.86
1:A:697:THR:HB	1:A:700:ASN:HB3	1.57	0.86
1:A:779:PHE:O	1:A:782:VAL:HG12	1.76	0.84
1:A:655:ILE:HD12	1:A:708:LEU:HD13	1.61	0.81
1:A:767:ARG:HB2	1:A:824:ASP:HB3	1.65	0.79
1:A:794:LEU:HD13	1:A:861:ILE:HD11	1.65	0.79
1:A:697:THR:O	1:A:699:GLU:N	2.15	0.78
1:A:672:PHE:HE1	1:A:696:VAL:HA	1.48	0.78
1:A:825:LYS:O	1:A:826:VAL:CG2	2.33	0.76
1:A:380:GLN:OE1	1:A:381:GLY:N	2.18	0.76
1:A:665:GLU:O	1:A:666:CYS:HB3	1.86	0.74
1:A:858:LEU:O	1:A:862:GLU:HG2	1.87	0.74
1:A:655:ILE:HD11	1:A:708:LEU:CD1	2.16	0.72
1:A:699:GLU:OE2	1:A:699:GLU:HA	1.89	0.71
1:A:779:PHE:O	1:A:782:VAL:CG1	2.37	0.70
1:A:655:ILE:HD12	1:A:708:LEU:CD1	2.19	0.70
1:A:665:GLU:HB3	1:A:698:GLU:OE1	1.91	0.70
1:A:535:MET:O	1:A:536:LYS:HB2	1.92	0.68
1:A:376:ARG:CA	1:A:379:LEU:HG	2.24	0.67
1:A:530:GLN:O	1:A:534:ASN:OD1	2.13	0.67
1:A:704:TYR:O	1:A:708:LEU:HB2	1.95	0.67
1:A:573:ASN:HB3	1:A:576:TYR:CE2	2.31	0.66
1:A:804:LEU:HD12	1:A:805:PRO:HD2	1.78	0.66
1:A:672:PHE:CE1	1:A:696:VAL:HB	2.32	0.64
1:A:790:LYS:NZ	1:A:862:GLU:OE1	2.27	0.63

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:376:ARG:HA	1:A:379:LEU:CG	2.27	0.63
1:A:674:GLN:HA	1:A:674:GLN:HE21	1.61	0.63
1:A:825:LYS:C	1:A:826:VAL:HG23	2.19	0.62
1:A:654:SER:O	1:A:658:ILE:HG12	1.99	0.62
1:A:827:GLY:O	1:A:828:LYS:HE2	2.00	0.61
1:A:376:ARG:O	1:A:379:LEU:HG	1.99	0.61
1:A:573:ASN:HB3	1:A:576:TYR:CD2	2.36	0.61
1:A:703:GLU:O	1:A:707:LEU:HB2	2.04	0.58
1:A:697:THR:C	1:A:699:GLU:H	2.07	0.58
1:A:672:PHE:CE1	1:A:696:VAL:HA	2.35	0.57
1:A:804:LEU:HD21	1:A:809:PHE:CE2	2.41	0.56
1:A:379:LEU:HD12	1:A:379:LEU:C	2.27	0.55
1:A:747:LEU:HD11	1:A:792:ILE:HG21	1.89	0.55
1:A:671:TYR:C	1:A:673:ILE:N	2.61	0.54
1:A:670:LEU:C	1:A:696:VAL:HG13	2.27	0.54
1:A:515:VAL:HG13	1:A:542:ARG:HD2	1.90	0.54
1:A:676:MET:HE3	1:A:715:ARG:HG3	1.91	0.52
1:A:766:TYR:CD1	1:A:776:ILE:HD13	2.44	0.52
1:A:697:THR:O	1:A:700:ASN:N	2.34	0.52
1:A:665:GLU:O	1:A:665:GLU:HG2	2.10	0.52
1:A:537:PRO:HG2	1:A:730:GLU:HB3	1.92	0.52
1:A:832:LEU:N	1:A:832:LEU:HD12	2.25	0.51
1:A:652:TYR:O	1:A:656:VAL:HG23	2.11	0.51
1:A:548:MET:CE	1:A:559:ILE:HD11	2.40	0.50
1:A:548:MET:HE3	1:A:559:ILE:HD11	1.94	0.50
1:A:676:MET:CE	1:A:715:ARG:HG3	2.42	0.49
1:A:671:TYR:HB2	1:A:673:ILE:HG12	1.95	0.49
1:A:665:GLU:CB	1:A:698:GLU:OE1	2.61	0.48
1:A:376:ARG:HA	1:A:379:LEU:CD2	2.43	0.48
1:A:376:ARG:C	1:A:379:LEU:HG	2.34	0.48
1:A:657:TRP:CE3	1:A:657:TRP:CA	2.85	0.48
1:A:779:PHE:C	1:A:782:VAL:HG12	2.35	0.47
1:A:520:SER:HB3	1:A:523:THR:OG1	2.15	0.47
1:A:665:GLU:N	1:A:698:GLU:OE1	2.49	0.46
1:A:572:LEU:HD21	1:A:624:THR:HG23	1.97	0.46
1:A:667:GLY:N	1:A:698:GLU:OE2	2.41	0.46
1:A:766:TYR:CE1	1:A:776:ILE:HD13	2.50	0.46
1:A:512:PRO:O	1:A:542:ARG:NH1	2.45	0.46
1:A:561:ARG:HG2	1:A:620:LYS:HD2	1.98	0.46
1:A:393:GLN:HG3	1:A:394:SER:N	2.30	0.46
1:A:665:GLU:CA	1:A:698:GLU:OE1	2.64	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:616:LEU:HD11	1:A:748:MET:HB3	1.98	0.45
1:A:556:TYR:HA	1:A:559:ILE:HG22	1.99	0.44
1:A:828:LYS:HD3	1:A:828:LYS:HA	1.74	0.44
1:A:672:PHE:HE2	1:A:703:GLU:HB3	1.83	0.44
1:A:525:PHE:HE1	1:A:571:VAL:HG12	1.83	0.43
1:A:670:LEU:C	1:A:696:VAL:CG1	2.87	0.43
1:A:766:TYR:CD1	1:A:776:ILE:CD1	3.01	0.43
1:A:537:PRO:CG	1:A:730:GLU:HB3	2.49	0.43
1:A:628:LEU:HB2	1:A:629:PRO:HD3	2.00	0.43
1:A:827:GLY:HA3	1:A:845:PRO:HG3	1.99	0.43
1:A:632:LYS:HG2	1:A:637:LYS:HB2	2.00	0.43
1:A:671:TYR:CB	1:A:673:ILE:HG12	2.49	0.42
1:A:767:ARG:HB2	1:A:824:ASP:CB	2.44	0.42
1:A:505:LEU:HD23	1:A:505:LEU:HA	1.89	0.42
1:A:670:LEU:HD23	1:A:670:LEU:HA	1.84	0.42
1:A:375:GLN:NE2	1:A:586:ASN:ND2	2.68	0.42
1:A:670:LEU:O	1:A:696:VAL:HG13	2.20	0.42
1:A:672:PHE:O	1:A:687:LEU:HB3	2.19	0.42
1:A:671:TYR:CA	1:A:696:VAL:HG12	2.50	0.41
1:A:674:GLN:HG3	1:A:711:TRP:CZ3	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	399/447 (89%)	365 (92%)	25 (6%)	9 (2%)	6 5

All (9) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	668	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	672	PHE
1	A	698	GLU
1	A	826	VAL
1	A	829	GLU
1	A	688	LYS
1	A	536	LYS
1	A	576	TYR
1	A	818	PRO

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	374/406 (92%)	345 (92%)	29 (8%)	12 16

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	360	ARG
1	A	376	ARG
1	A	379	LEU
1	A	380	GLN
1	A	384	GLN
1	A	393	GLN
1	A	394	SER
1	A	493	ARG
1	A	503	ARG
1	A	566	LEU
1	A	601	HIS
1	A	642	LYS
1	A	657	TRP
1	A	666	CYS
1	A	674	GLN
1	A	681	LYS
1	A	692	GLU
1	A	696	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	699	GLU
1	A	700	ASN
1	A	703	GLU
1	A	708	LEU
1	A	712	ARG
1	A	763	SER
1	A	769	TYR
1	A	838[A]	CYS
1	A	838[B]	CYS
1	A	858	LEU
1	A	862	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	375	GLN
1	A	636	ASN
1	A	685	HIS

5.3.3 RNA

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no monosaccharides in this entry.

5.6 Ligand geometry

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å^2)	Q<0.9
1	A	408/447 (91%)	-0.00	18 (4%) 34 41	30, 65, 107, 154	0

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	360	ARG	6.8
1	A	554	LEU	5.2
1	A	657	TRP	4.8
1	A	379	LEU	4.6
1	A	380	GLN	3.7
1	A	358	TRP	3.5
1	A	668	LEU	3.0
1	A	359	GLN	2.8
1	A	378	GLN	2.4
1	A	671	TYR	2.3
1	A	673	ILE	2.3
1	A	375	GLN	2.2
1	A	695	ARG	2.2
1	A	363	ALA	2.2
1	A	369	TYR	2.1
1	A	681	LYS	2.1
1	A	828	LYS	2.0
1	A	364	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	NA	A	901	1/1	0.93	0.19	59,59,59,59	0

6.5 Other polymers (i)

There are no such residues in this entry.