



# wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 7, 2020 – 02:16 PM BST

PDB ID : 1TKR  
Title : Human Dipeptidyl Peptidase IV/CD26 inhibited with Diisopropyl FluoroPhosphate  
Authors : Bjelke, J.R.; Christensen, J.; Branner, S.; Wagtmann, N.; Olsen, C.; Kanstrup, A.B.; Rasmussen, H.B.  
Deposited on : 2004-06-09  
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.13.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13.1


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.









Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	728	<div style="display: flex; align-items: center;"> <div style="width: 2%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 63%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 32%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 1%; height: 10px; background-color: orange; margin-right: 5px;"></div> <div style="width: 2%; height: 10px; background-color: red;"></div> </div>
1	B	728	<div style="display: flex; align-items: center;"> <div style="width: 2%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 58%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 37%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 5%; height: 10px; background-color: orange;"></div> </div>
2	C	3	<div style="display: flex; align-items: center;"> <div style="width: 33%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 67%; height: 10px; background-color: orange;"></div> </div>
3	D	3	<div style="display: flex; align-items: center;"> <div style="width: 33%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 67%; height: 10px; background-color: yellow;"></div> </div>
4	E	2	<div style="display: flex; align-items: center;"> <div style="width: 50%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 50%; height: 10px; background-color: orange;"></div> </div>
4	G	2	<div style="display: flex; align-items: center;"> <div style="width: 100%; height: 10px; background-color: green;"></div> </div>
4	H	2	<div style="display: flex; align-items: center;"> <div style="width: 50%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 50%; height: 10px; background-color: yellow;"></div> </div>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
4	J	2	 50% 50%
5	F	3	 33% 67%
6	I	2	 50% 50%
7	K	3	 33% 67%
7	L	3	 67% 33%
8	M	2	 100%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NAG	D	1	X	-	-	-
3	FUC	D	3	-	-	-	X
4	NAG	J	1	X	-	-	-
5	NDG	F	2	-	-	-	X
8	NAG	M	1	X	-	X	-
8	NDG	M	2	-	-	X	X

## 2 Entry composition [i](#)


There are 11 unique types of molecules in this entry. The entry contains 12996 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Dipeptidyl peptidase IV.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	726	Total	C	N	O	S	0	0	0
			5948	3816	980	1126	26			
1	B	728	Total	C	N	O	S	0	0	0
			5964	3827	982	1129	26			

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-alpha-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	C	3	Total	C	N	O	0	0	0
			38	22	2	14			

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	D	3	Total	C	N	O	0	0	0
			38	22	2	14			

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
4	E	2	28	16	2	10	0	0	0
4	G	2	28	16	2	10	0	0	0
4	H	2	28	16	2	10	0	0	0
4	J	2	28	16	2	10	0	0	0

- Molecule 5 is an oligosaccharide called alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-alpha-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
5	F	3	39	22	2	15	0	0	0

- Molecule 6 is an oligosaccharide called alpha-L-fucopyranose-(1-6)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
6	I	2	24	14	1	9	0	0	0

- Molecule 7 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
7	K	3	Total	C	N	O	0	0	0
			39	22	2	15			
7	L	3	Total	C	N	O	0	0	0
			39	22	2	15			

- Molecule 8 is an oligosaccharide called 2-acetamido-2-deoxy-alpha-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
8	M	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 9 is DIISOPROPYL PHOSPHONATE (three-letter code: DFP) (formula: C<sub>6</sub>H<sub>15</sub>O<sub>3</sub>P).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
9	A	1	Total	C	O	P	0	0
			10	6	3	1		
9	B	1	Total	C	O	P	0	0
			10	6	3	1		

- Molecule 10 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
10	B	1	14	8	1	5	0	0

- Molecule 11 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
11	A	378	378	378	0	0
11	B	315	315	315	0	0







- Molecule 2: 2-acetamido-2-deoxy-alpha-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain C: 33% 67%

MAG1  
MAG2  
FUC3

- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D: 33% 67%

MAG1  
MAG2  
FUC3

- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E: 50% 50%

MAG1  
MAG2

- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G:  100%

  
MAG1  
MAG2

- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain H:  50% 50%

  
MAG1  
MAG2

- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain J:  50% 50%

  
MAG1  
MAG2

- Molecule 5: alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-alpha-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:  33% 67%

  
MAG1  
MAG2  
MAG3

- Molecule 6: alpha-L-fucopyranose-(1-6)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain I:  50% 50%

  
MAG1  
FUG2

- Molecule 7: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain K:  33% 67%

  
MAG1  
MAG2  
BGL3

- Molecule 7: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain L:  67% 33%

  
MAG1  
MAG2  
BGL3

- Molecule 8: 2-acetamido-2-deoxy-alpha-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain M:

100%

MAG1  
MAG2

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	119.22Å 123.46Å 131.00Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.60 – 2.70 30.59 – 2.70	Depositor EDS
% Data completeness (in resolution range)	92.7 (30.60-2.70) 92.4 (30.59-2.70)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.23 (at 2.68Å)	Xtrriage
Refinement program	CNX 2000.1	Depositor
R, $R_{free}$	0.203 , 0.268 0.198 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	25.3	Xtrriage
Anisotropy	0.429	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 43.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.023 for k,h,-l	Xtrriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	12996	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	27.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.21% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, NAG, DFP, NDG, FUC, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.36	0/6119	0.62	0/8321
1	B	0.37	0/6136	0.61	0/8344
All	All	0.37	0/12255	0.61	0/16665

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5948	0	5659	227	0
1	B	5964	0	5676	267	0
2	C	38	0	33	1	0
3	D	38	0	34	6	0
4	E	28	0	25	1	0
4	G	28	0	25	0	0
4	H	28	0	25	0	0
4	J	28	0	25	2	0
5	F	39	0	33	4	0
6	I	24	0	22	2	0
7	K	39	0	34	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	L	39	0	34	0	0
8	M	28	0	24	9	0
9	A	10	0	14	2	0
9	B	10	0	14	2	0
10	B	14	0	13	0	0
11	A	378	0	0	21	0
11	B	315	0	0	21	0
All	All	12996	0	11690	505	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 21.

The worst 5 of 505 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
9:A:1:DFP:H21	11:A:1159:HOH:O	1.30	1.28
1:A:289:ALA:HB1	1:A:290:PRO:HA	1.34	1.08
1:B:289:ALA:HB1	1:B:290:PRO:HA	1.38	1.03
1:B:458:SER:HB3	1:B:471:ARG:HB2	1.44	0.99
1:B:358:ARG:HH11	1:B:358:ARG:HB3	1.22	0.98

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	724/728 (100%)	656 (91%)	60 (8%)	8 (1%)	14 34
1	B	726/728 (100%)	641 (88%)	74 (10%)	11 (2%)	10 26
All	All	1450/1456 (100%)	1297 (89%)	134 (9%)	19 (1%)	12 30

5 of 19 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	320	GLN
1	B	289	ALA
1	B	320	GLN
1	B	399	LYS
1	B	401	THR

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	651/653 (100%)	606 (93%)	45 (7%)	15	35
1	B	653/653 (100%)	610 (93%)	43 (7%)	16	38
All	All	1304/1306 (100%)	1216 (93%)	88 (7%)	16	37

5 of 88 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	658	ARG
1	B	74	ASN
1	B	689	MET
1	A	689	MET
1	A	761	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 50 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	112	GLN
1	B	170	ASN
1	B	712	HIS
1	B	119	ASN
1	B	138	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

27 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	NAG	C	1	1,2	14,14,15	0.70	0	17,19,21	1.28	2 (11%)
2	NDG	C	2	2	14,14,15	0.68	0	17,19,21	0.81	1 (5%)
2	FUC	C	3	2	10,10,11	0.49	0	14,14,16	0.46	0
3	NAG	D	1	1,3	14,14,15	0.77	0	17,19,21	0.99	0
3	NAG	D	2	3	14,14,15	0.68	0	17,19,21	0.67	0
3	FUC	D	3	3	10,10,11	0.51	0	14,14,16	0.37	0
4	NAG	E	1	1,4	14,14,15	0.53	0	17,19,21	0.91	1 (5%)
4	NAG	E	2	4	14,14,15	0.47	0	17,19,21	0.68	0
5	NAG	F	1	1,5	14,14,15	0.59	0	17,19,21	1.03	2 (11%)
5	NDG	F	2	5	14,14,15	0.95	1 (7%)	17,19,21	0.89	0
5	MAN	F	3	5	11,11,12	0.66	0	15,15,17	0.50	0
4	NAG	G	1	1,4	14,14,15	0.48	0	17,19,21	0.60	0
4	NAG	G	2	4	14,14,15	0.51	0	17,19,21	0.60	0
4	NAG	H	1	1,4	14,14,15	0.49	0	17,19,21	0.84	1 (5%)
4	NAG	H	2	4	14,14,15	0.69	0	17,19,21	0.57	0
6	NAG	I	1	1,6	14,14,15	0.54	0	17,19,21	0.64	0
6	FUC	I	2	6	10,10,11	0.53	0	14,14,16	0.44	0
4	NAG	J	1	1,4	14,14,15	0.68	0	17,19,21	0.81	1 (5%)
4	NAG	J	2	4	14,14,15	0.49	0	17,19,21	0.74	0
7	NAG	K	1	1,7	14,14,15	0.46	0	17,19,21	0.87	1 (5%)
7	NAG	K	2	7	14,14,15	0.46	0	17,19,21	0.82	1 (5%)
7	BMA	K	3	7	11,11,12	0.59	0	15,15,17	0.22	0
7	NAG	L	1	1,7	14,14,15	0.51	0	17,19,21	0.61	0
7	NAG	L	2	7	14,14,15	0.63	0	17,19,21	1.11	2 (11%)
7	BMA	L	3	7	11,11,12	0.43	0	15,15,17	0.37	0



Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
8	NAG	M	1	1,8	14,14,15	0.61	0	17,19,21	0.95	1 (5%)
8	NDG	M	2	8	14,14,15	0.74	1 (7%)	17,19,21	0.77	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	1,2	-	0/6/23/26	0/1/1/1
2	NDG	C	2	2	-	1/6/23/26	0/1/1/1
2	FUC	C	3	2	-	-	0/1/1/1
3	NAG	D	1	1,3	1/1/5/7	0/6/23/26	0/1/1/1
3	NAG	D	2	3	-	3/6/23/26	0/1/1/1
3	FUC	D	3	3	-	-	0/1/1/1
4	NAG	E	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	E	2	4	-	2/6/23/26	0/1/1/1
5	NAG	F	1	1,5	-	1/6/23/26	0/1/1/1
5	NDG	F	2	5	-	1/6/23/26	0/1/1/1
5	MAN	F	3	5	-	2/2/19/22	0/1/1/1
4	NAG	G	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	G	2	4	-	1/6/23/26	0/1/1/1
4	NAG	H	1	1,4	-	2/6/23/26	0/1/1/1
4	NAG	H	2	4	-	2/6/23/26	0/1/1/1
6	NAG	I	1	1,6	-	2/6/23/26	0/1/1/1
6	FUC	I	2	6	-	-	0/1/1/1
4	NAG	J	1	1,4	1/1/5/7	1/6/23/26	0/1/1/1
4	NAG	J	2	4	-	2/6/23/26	0/1/1/1
7	NAG	K	1	1,7	-	0/6/23/26	0/1/1/1
7	NAG	K	2	7	-	0/6/23/26	0/1/1/1
7	BMA	K	3	7	-	2/2/19/22	0/1/1/1
7	NAG	L	1	1,7	-	0/6/23/26	0/1/1/1
7	NAG	L	2	7	-	0/6/23/26	0/1/1/1
7	BMA	L	3	7	-	0/2/19/22	0/1/1/1
8	NAG	M	1	1,8	1/1/5/7	1/6/23/26	0/1/1/1
8	NDG	M	2	8	-	0/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	F	2	NDG	C1-C2	2.85	1.56	1.52
8	M	2	NDG	C1-C2	2.11	1.55	1.52

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	1	NAG	C4-C3-C2	3.36	115.95	111.02
5	F	1	NAG	C2-N2-C7	-2.85	118.84	122.90
8	M	1	NAG	C4-C3-C2	2.57	114.79	111.02
7	L	2	NAG	C4-C3-C2	-2.49	107.36	111.02
7	K	1	NAG	C2-N2-C7	-2.43	119.44	122.90

All (3) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	J	1	NAG	C1
8	M	1	NAG	C1
3	D	1	NAG	C1

5 of 23 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	K	3	BMA	O5-C5-C6-O6
4	J	2	NAG	O5-C5-C6-O6
5	F	3	MAN	C4-C5-C6-O6
5	F	3	MAN	O5-C5-C6-O6
7	K	3	BMA	C4-C5-C6-O6

There are no ring outliers.

12 monomers are involved in 25 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	1	NAG	1	0
3	D	2	NAG	5	0
5	F	1	NAG	1	0
4	J	1	NAG	2	0
8	M	1	NAG	9	0
2	C	2	NDG	1	0
5	F	3	MAN	2	0
8	M	2	NDG	8	0
6	I	1	NAG	2	0
5	F	2	NDG	2	0


*Continued on next page...*


Continued from previous page...


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	1	NAG	5	0
4	E	1	NAG	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



























## 5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
9	DFP	B	1	1	6,9,9	1.52	2 (33%)	6,11,11	0.43	0
9	DFP	A	1	1	6,9,9	1.51	2 (33%)	6,11,11	0.41	0
10	NAG	B	769	1	14,14,15	0.54	0	17,19,21	0.67	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
9	DFP	B	1	1	-	2/4/8/8	-
9	DFP	A	1	1	-	2/4/8/8	-
10	NAG	B	769	1	-	0/6/23/26	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
9	A	1	DFP	O1P-C1	-2.52	1.42	1.45
9	B	1	DFP	O1P-C1	-2.51	1.42	1.45
9	B	1	DFP	O2P-C1'	-2.29	1.42	1.45
9	A	1	DFP	O2P-C1'	-2.17	1.43	1.45

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
9	B	1	DFP	C3'-C1'-O2P-P
9	A	1	DFP	C3'-C1'-O2P-P
9	B	1	DFP	C2'-C1'-O2P-P
9	A	1	DFP	C2'-C1'-O2P-P

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
9	B	1	DFP	2	0
9	A	1	DFP	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å <sup>2</sup> )	Q < 0.9
1	A	726/728 (99%)	-0.52	6 (0%) 86 87	9, 22, 44, 67	0
1	B	728/728 (100%)	-0.36	13 (1%) 68 70	6, 24, 54, 94	0
All	All	1454/1456 (99%)	-0.44	19 (1%) 77 78	6, 23, 48, 94	0

The worst 5 of 19 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	39	SER	4.1
1	B	99	GLY	3.7
1	B	39	SER	3.2
1	B	97	GLU	3.1
1	A	73	GLU	3.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q < 0.9
5	MAN	F	3	11/12	0.47	0.36	86,88,88,89	0
4	NAG	H	2	14/15	0.58	0.40	72,75,76,77	0
5	NDG	F	2	14/15	0.64	0.41	67,74,76,82	0
3	NAG	D	2	14/15	0.68	0.35	77,79,84,84	0
3	NAG	D	1	14/15	0.70	0.30	65,69,77,80	0

*Continued on next page...*




*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
8	NDG	M	2	14/15	0.72	0.41	79,81,84,85	0
3	FUC	D	3	10/11	0.74	0.51	83,86,87,88	0
4	NAG	J	2	14/15	0.78	0.30	65,67,68,69	0
8	NAG	M	1	14/15	0.81	0.32	63,67,72,76	0
2	NDG	C	2	14/15	0.81	0.45	72,74,76,77	0
4	NAG	G	2	14/15	0.82	0.35	65,69,71,72	0
6	NAG	I	1	14/15	0.84	0.23	63,66,71,72	0
4	NAG	J	1	14/15	0.84	0.30	55,58,62,64	0
4	NAG	E	2	14/15	0.85	0.34	54,56,59,59	0
7	BMA	K	3	11/12	0.85	0.29	64,66,67,68	0
6	FUC	I	2	10/11	0.86	0.35	74,75,75,75	0
7	BMA	L	3	11/12	0.86	0.24	53,56,58,59	0
2	FUC	C	3	10/11	0.88	0.24	63,64,65,65	0
2	NAG	C	1	14/15	0.89	0.20	54,58,62,67	0
7	NAG	K	2	14/15	0.89	0.26	50,54,59,62	0
4	NAG	H	1	14/15	0.89	0.28	57,61,65,70	0
4	NAG	G	1	14/15	0.91	0.20	49,52,55,60	0
4	NAG	E	1	14/15	0.91	0.21	39,41,45,50	0
7	NAG	L	2	14/15	0.92	0.19	32,38,44,47	0
5	NAG	F	1	14/15	0.92	0.17	39,42,48,58	0
7	NAG	K	1	14/15	0.95	0.14	28,30,35,43	0
7	NAG	L	1	14/15	0.95	0.14	20,27,32,33	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around Chain C:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around Chain D:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around Chain E:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around Chain G:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around Chain H:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around Chain J:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around Chain F:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)













**Electron density around Chain L:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around Chain M:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
10	NAG	B	769	14/15	0.77	0.39	68,73,74,75	0
9	DFP	A	1	10/10	0.96	0.19	16,23,27,29	0
9	DFP	B	1	10/10	0.97	0.16	25,29,35,35	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.