



wwPDB X-ray Structure Validation Summary Report ⓘ

May 29, 2020 – 10:58 pm BST

PDB ID : 3TKK
Title : Crystal Structure Analysis of a recombinant predicted acetamidase/ formamidase from the thermophile thermoanaerobacter tengcongensis
Authors : Qian, M.; Huang, Q.; Wu, G.; Lai, L.; Tang, Y.; Pei, J.; Kusunoki, M.
Deposited on : 2011-08-26
Resolution : 1.99 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.99 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	301	67% (green), 28% (yellow), 2% (orange), 2% (red), 1% (grey)
1	B	301	73% (green), 23% (yellow), 2% (orange), 2% (red), 1% (grey)
1	C	301	72% (green), 24% (yellow), 5% (orange), 1% (red), 1% (grey)
1	D	301	67% (green), 27% (yellow), 5% (orange), 1% (red), 1% (grey)

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 9872 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Predicted acetamidase/formamidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	301	2257	1436	369	442	10	0	0	0
1	B	301	2257	1436	369	442	10	0	0	0
1	C	301	2257	1436	369	442	10	0	0	0
1	D	301	2257	1436	369	442	10	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	SER	-	EXPRESSION TAG	UNP Q8R8S5
A	-1	HIS	-	EXPRESSION TAG	UNP Q8R8S5
B	-2	SER	-	EXPRESSION TAG	UNP Q8R8S5
B	-1	HIS	-	EXPRESSION TAG	UNP Q8R8S5
C	-2	SER	-	EXPRESSION TAG	UNP Q8R8S5
C	-1	HIS	-	EXPRESSION TAG	UNP Q8R8S5
D	-2	SER	-	EXPRESSION TAG	UNP Q8R8S5
D	-1	HIS	-	EXPRESSION TAG	UNP Q8R8S5

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total 1	Ca 1	0	0
2	A	2	Total 2	Ca 2	0	0
2	D	1	Total 1	Ca 1	0	0
2	C	1	Total 1	Ca 1	0	0

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total 1	Zn 1	0	0
3	A	1	Total 1	Zn 1	0	0
3	D	1	Total 1	Zn 1	0	0
3	C	1	Total 1	Zn 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	292	Total 292	O 292	0	0
4	B	153	Total 153	O 153	0	0
4	C	254	Total 254	O 254	0	0
4	D	136	Total 136	O 136	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Predicted acetamidase/formamidase



- Molecule 1: Predicted acetamidase/formamidase




- Molecule 1: Predicted acetamidase/formamidase



- Molecule 1: Predicted acetamidase/formamidase

Chain D:  67% 27% 5%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	41.23Å 152.88Å 100.25Å 90.00° 99.49° 90.00°	Depositor
Resolution (Å)	50.00 – 1.99 40.67 – 1.99	Depositor EDS
% Data completeness (in resolution range)	94.7 (50.00-1.99) 94.7 (40.67-1.99)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.61 (at 2.00Å)	Xtrriage
Refinement program	REFMAC 5.4.0077	Depositor
R, R_{free}	0.174 , 0.237 0.240 , 0.283	Depositor DCC
R_{free} test set	3985 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	26.1	Xtrriage
Anisotropy	0.692	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 30.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.023 for h,-k,-h-l	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	9872	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.19% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: CSD, ZN, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.52	25/2285 (1.1%)	1.22	6/3088 (0.2%)
1	B	1.03	0/2285	0.98	5/3088 (0.2%)
1	C	1.29	5/2285 (0.2%)	1.11	7/3088 (0.2%)
1	D	1.04	3/2285 (0.1%)	0.95	1/3088 (0.0%)
All	All	1.24	33/9140 (0.4%)	1.07	19/12352 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

The worst 5 of 33 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	192	VAL	CB-CG1	-11.70	1.28	1.52
1	A	137	VAL	CB-CG2	6.98	1.67	1.52
1	A	69	ALA	CA-CB	6.56	1.66	1.52
1	C	198	GLU	CG-CD	6.54	1.61	1.51
1	C	146	CYS	CB-SG	-6.31	1.71	1.82

The worst 5 of 19 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	8	ASP	CB-CG-OD1	7.59	125.14	118.30
1	A	202	LYS	CD-CE-NZ	6.92	127.61	111.70
1	B	284	ARG	NE-CZ-NH1	-6.66	116.97	120.30
1	C	235	LEU	CB-CG-CD2	-6.41	100.11	111.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	157	ASP	CB-CG-OD1	6.16	123.85	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	160	LEU	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2257	0	2294	74	1
1	B	2257	0	2292	90	0
1	C	2257	0	2293	75	0
1	D	2257	0	2293	107	0
2	A	2	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	292	0	0	14	1
4	B	153	0	0	9	0
4	C	254	0	0	14	0
4	D	136	0	0	11	0
All	All	9872	0	9172	327	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

The worst 5 of 327 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:45:ASN:HB3	1:B:110:VAL:CG2	1.22	1.57
1:A:135:ILE:CG1	1:A:135:ILE:CD1	1.78	1.56
1:A:161:ILE:CD1	1:A:161:ILE:CG1	1.76	1.56
1:B:45:ASN:CB	1:B:110:VAL:HG23	1.54	1.33
1:B:45:ASN:CB	1:B:110:VAL:CG2	2.05	1.33

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:114:LYS:NZ	4:A:455:HOH:O[1_455]	1.55	0.65

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	298/301 (99%)	276 (93%)	19 (6%)	3 (1%)	15	9
1	B	298/301 (99%)	273 (92%)	21 (7%)	4 (1%)	12	6
1	C	298/301 (99%)	277 (93%)	17 (6%)	4 (1%)	12	6
1	D	298/301 (99%)	276 (93%)	16 (5%)	6 (2%)	7	3
All	All	1192/1204 (99%)	1102 (92%)	73 (6%)	17 (1%)	11	5

5 of 17 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	194	VAL
1	B	44	SER
1	B	194	VAL
1	C	52	MET
1	C	194	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	245/245 (100%)	229 (94%)	16 (6%)	17	12
1	B	245/245 (100%)	226 (92%)	19 (8%)	12	8
1	C	245/245 (100%)	224 (91%)	21 (9%)	10	6
1	D	245/245 (100%)	227 (93%)	18 (7%)	14	9
All	All	980/980 (100%)	906 (92%)	74 (8%)	13	8

5 of 74 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	237	LYS
1	C	115	ASP
1	D	180	LEU
1	B	249	LEU
1	C	78	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 20 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	253	HIS
1	C	10	HIS
1	C	244	HIS
1	B	214	ASN
1	B	244	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	CSD	A	37	1,3	3,7,8	2.23	1 (33%)	1,8,10	7.72	1 (100%)
1	CSD	B	37	1,3	3,7,8	0.83	0	1,8,10	5.78	1 (100%)
1	CSD	C	37	1,3	3,7,8	0.60	0	1,8,10	9.74	1 (100%)
1	CSD	D	37	1,3	3,7,8	2.30	1 (33%)	1,8,10	6.47	1 (100%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSD	A	37	1,3	-	2/2/6/8	-
1	CSD	B	37	1,3	-	2/2/6/8	-
1	CSD	C	37	1,3	-	1/2/6/8	-
1	CSD	D	37	1,3	-	0/2/6/8	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	37	CSD	CB-SG	3.72	1.99	1.79
1	A	37	CSD	CB-SG	3.20	1.97	1.79

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	37	CSD	OD1-SG-CB	9.74	124.07	105.54
1	A	37	CSD	OD1-SG-CB	7.72	120.23	105.54
1	D	37	CSD	OD1-SG-CB	6.47	117.85	105.54
1	B	37	CSD	OD1-SG-CB	5.78	116.54	105.54

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	37	CSD	N-CA-CB-SG
1	A	37	CSD	CA-CB-SG-OD1
1	B	37	CSD	N-CA-CB-SG
1	B	37	CSD	CA-CB-SG-OD1
1	C	37	CSD	CA-CB-SG-OD1

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	D	37	CSD	2	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 9 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers

Unable to reproduce the depositors R factor - this section is therefore empty.