



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 11, 2023 – 02:32 PM EDT

PDB ID : 7TLT
Title : SARS-CoV-2 Spike-derived peptide S489-497 (YFPLQSYGF) presented by HLA-A*29:02
Authors : Murdolo, L.D.; Szeto, C.; Gras, S.
Deposited on : 2022-01-18
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	365	
1	C	365	
2	B	100	
2	D	100	
3	E	9	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	9	 89% 11%

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 6403 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HLA class I histocompatibility antigen, A alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	275	Total	C	N	O	S	0	4	0
			2240	1388	407	434	11			
1	C	275	Total	C	N	O	S	0	5	0
			2254	1396	410	437	11			

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	99	Total	C	N	O	S	0	2	0
			841	536	142	159	4			
2	D	100	Total	C	N	O	S	0	2	0
			847	540	142	161	4			


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	MET	-	initiating methionine	UNP P61769
D	0	MET	-	initiating methionine	UNP P61769

- Molecule 3 is a protein called Spike protein S1 peptide.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	E	9	Total	C	N	O	0	0	0
			81	57	10	14			
3	F	9	Total	C	N	O	0	0	0
			81	57	10	14			

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total O S 5 4 1	0	0
4	A	1	Total O S 5 4 1	0	0
4	B	1	Total O S 5 4 1	0	0
4	D	1	Total O S 5 4 1	0	0
4	F	1	Total O S 5 4 1	0	0

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total Mg 1 1	0	0
5	B	1	Total Mg 1 1	0	0

- Molecule 6 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	B	1	Total Na 1 1	0	0
6	C	1	Total Na 1 1	0	0

- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	10	Total O 10 10	0	0
7	B	6	Total O 6 6	0	0
7	C	9	Total O 9 9	0	0
7	D	2	Total O 2 2	0	0
7	E	1	Total O 1 1	0	0
7	F	2	Total O 2 2	0	0



- Molecule 3: Spike protein S1 peptide

Chain E: 78% 22%



- Molecule 3: Spike protein S1 peptide

Chain F: 89% 11%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	48.81Å 64.47Å 76.42Å 106.53° 92.43° 97.51°	Depositor
Resolution (Å)	41.29 – 2.30 41.29 – 2.30	Depositor EDS
% Data completeness (in resolution range)	95.4 (41.29-2.30) 95.4 (41.29-2.30)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.62 (at 2.29Å)	Xtrriage
Refinement program	BUSTER 2.10.3	Depositor
R, R_{free}	0.192 , 0.238 0.204 , 0.258	Depositor DCC
R_{free} test set	1826 reflections (4.89%)	wwPDB-VP
Wilson B-factor (Å ²)	39.7	Xtrriage
Anisotropy	0.716	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 43.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.52$, $\langle L^2 \rangle = 0.35$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6403	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.39% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, NA, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.53	0/2308	0.71	0/3131
1	C	0.52	0/2322	0.67	0/3150
2	B	0.50	0/867	0.69	0/1171
2	D	0.50	0/873	0.68	0/1180
3	E	0.58	0/85	0.61	0/113
3	F	0.56	0/85	0.56	0/113
All	All	0.52	0/6540	0.69	0/8858

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2240	0	2108	8	0
1	C	2254	0	2120	10	0
2	B	841	0	810	2	0
2	D	847	0	816	3	0
3	E	81	0	72	1	0
3	F	81	0	72	1	0
4	A	10	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	5	0	0	0	0
4	D	5	0	0	0	0
4	F	5	0	0	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
6	B	1	0	0	0	0
6	C	1	0	0	0	0
7	A	10	0	0	0	0
7	B	6	0	0	0	0
7	C	9	0	0	0	0
7	D	2	0	0	0	0
7	E	1	0	0	0	0
7	F	2	0	0	0	0
All	All	6403	0	5998	21	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:45:MET:CE	1:A:67:VAL:HG21	2.36	0.56
1:C:258:THR:HG23	1:C:260:HIS:NE2	2.23	0.54
1:C:195:SER:HB3	1:C:198[A]:GLU:H	1.76	0.51
1:C:195:SER:HB3	1:C:198[B]:GLU:H	1.76	0.51
1:A:194:VAL:HB	1:A:198:GLU:HG3	1.92	0.51
1:C:103:VAL:HG11	1:C:165:VAL:HG23	1.92	0.50
1:C:73:THR:OG1	3:F:6:SER:HB2	2.13	0.48
1:A:117:ALA:HB2	2:B:60:TRP:CE2	2.48	0.48
1:A:63:GLN:HE22	3:E:2:PHE:H	1.60	0.47
1:A:28:VAL:HG11	1:A:179:LEU:HD13	1.95	0.47
1:C:175:GLY:HA3	1:C:179:LEU:HD22	1.99	0.45
2:D:69:GLU:H	2:D:69:GLU:CD	2.22	0.43
1:C:103:VAL:CG1	1:C:165:VAL:HG23	2.50	0.42
1:A:178:THR:HA	1:A:181:ARG:HD3	2.01	0.42
1:A:190:THR:HG21	2:B:98:ASP:OD1	2.20	0.42
1:C:121:LYS:HZ1	1:C:122:ASP:H	1.67	0.41
1:C:203:CYS:CB	1:C:259[A]:CYS:HG	2.29	0.41
2:D:21:ASN:HB3	2:D:70:PHE:CE1	2.55	0.41
1:A:45:MET:HE2	1:A:67:VAL:HG21	2.01	0.41
2:D:24:ASN:HB3	2:D:65:LEU:HD11	2.03	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:218:GLN:HB2	1:C:258:THR:HG22	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	277/365 (76%)	271 (98%)	6 (2%)	0	100	100
1	C	278/365 (76%)	272 (98%)	6 (2%)	0	100	100
2	B	99/100 (99%)	97 (98%)	2 (2%)	0	100	100
2	D	100/100 (100%)	97 (97%)	3 (3%)	0	100	100
3	E	7/9 (78%)	6 (86%)	1 (14%)	0	100	100
3	F	7/9 (78%)	6 (86%)	1 (14%)	0	100	100
All	All	768/948 (81%)	749 (98%)	19 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	235/298 (79%)	226 (96%)	9 (4%)	33	47
1	C	236/298 (79%)	225 (95%)	11 (5%)	26	37

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	96/95 (101%)	91 (95%)	5 (5%)	23	32
2	D	97/95 (102%)	91 (94%)	6 (6%)	18	25
3	E	8/8 (100%)	7 (88%)	1 (12%)	4	5
3	F	8/8 (100%)	8 (100%)	0	100	100
All	All	680/802 (85%)	648 (95%)	32 (5%)	27	37

All (32) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	35	ARG
1	A	88	SER
1	A	121	LYS
1	A	132[A]	SER
1	A	132[B]	SER
1	A	166	GLU
1	A	229	GLU
1	A	232	GLU
1	A	271	THR
2	B	19	LYS
2	B	20	SER
2	B	40	LEU
2	B	47	GLU
2	B	89	GLN
1	C	35	ARG
1	C	45	MET
1	C	89	GLU
1	C	116	ASP
1	C	179	LEU
1	C	225	THR
1	C	251	SER
1	C	258	THR
1	C	259[A]	CYS
1	C	259[B]	CYS
1	C	270	LEU
2	D	1	ILE
2	D	20	SER
2	D	47	GLU
2	D	68	THR
2	D	70	PHE
2	D	77	GLU
3	E	4	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	63	GLN
1	A	226	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 4 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	SO4	A	402	-	4,4,4	0.16	0	6,6,6	0.11	0
4	SO4	A	401	-	4,4,4	0.32	0	6,6,6	0.45	0
4	SO4	D	101	-	4,4,4	0.22	0	6,6,6	0.33	0
4	SO4	B	101	-	4,4,4	0.16	0	6,6,6	0.08	0
4	SO4	F	101	-	4,4,4	0.17	0	6,6,6	0.12	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	275/365 (75%)	0.04	3 (1%) 80 85	31, 45, 63, 104	0
1	C	275/365 (75%)	0.03	0 100 100	30, 45, 72, 87	0
2	B	99/100 (99%)	0.17	1 (1%) 82 86	31, 45, 76, 87	0
2	D	100/100 (100%)	0.54	11 (11%) 5 7	33, 54, 78, 85	2 (2%)
3	E	9/9 (100%)	0.02	0 100 100	35, 39, 42, 47	0
3	F	9/9 (100%)	0.03	0 100 100	34, 38, 42, 48	0
All	All	767/948 (80%)	0.12	15 (1%) 65 71	30, 46, 73, 104	2 (0%)

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	0	MET	4.4
2	D	79	ALA	4.2
2	D	18	GLY	3.5
2	D	20	SER	3.3
1	A	196	ASP	3.1
2	D	69	GLU	2.7
1	A	192	HIS	2.6
2	D	74	GLU	2.5
2	D	48	LYS	2.3
2	D	21	ASN	2.3
2	B	1	ILE	2.2
2	D	15	ALA	2.2
1	A	274	TRP	2.1
2	D	17	ASN	2.1
2	D	22	PHE	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	NA	B	102	1/1	0.85	0.17	49,49,49,49	0
6	NA	C	401	1/1	0.89	0.25	57,57,57,57	0
4	SO4	A	402	5/5	0.93	0.15	113,113,114,114	0
4	SO4	D	101	5/5	0.93	0.13	72,73,74,75	0
4	SO4	F	101	5/5	0.94	0.20	110,111,112,112	0
5	MG	B	103	1/1	0.95	0.31	71,71,71,71	0
5	MG	A	403	1/1	0.97	0.24	42,42,42,42	0
4	SO4	A	401	5/5	0.98	0.13	60,60,62,62	0
4	SO4	B	101	5/5	0.98	0.10	81,83,84,85	0

6.5 Other polymers [i](#)

There are no such residues in this entry.