



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 5, 2024 – 09:10 AM EST

PDB ID : 1TMO  
Title : TRIMETHYLAMINE N-OXIDE REDUCTASE FROM SHEWANELLA MASSILIA  
Authors : Czjzek, M.; Dos Santos, J.P.; Giordano, G.; Mejean, V.  
Deposited on : 1998-08-03  
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	829	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	2MO	A	801	-	-	X	-

## 2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 6857 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TRIMETHYLAMINE N-OXIDE REDUCTASE.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	794	6255	3972	1080	1167	36	0	0	0

- Molecule 2 is GUANYLATE-O'-PHOSPHORIC ACID MONO-(2-AMINO-5,6-DIMERCAPTO-4-OXO-3,5,6,7,8A,9,10,10A-OCTAHYDRO-4H-8-OXA-1,3,9,10-TETRAAZA-ANTHRACEN-7-YLMETHYL) ESTER (three-letter code: 2MD) (formula: C<sub>20</sub>H<sub>28</sub>N<sub>10</sub>O<sub>13</sub>P<sub>2</sub>S<sub>2</sub>).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
			Total	C	N	O	P	S		
2	A	1	47	20	10	13	2	2	0	0
2	A	1	47	20	10	13	2	2	0	0

- Molecule 3 is MOLYBDENUM (IV)OXIDE (three-letter code: 2MO) (formula: MoO<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	Mo	O		
3	A	1	3	1	2	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	505	505	505	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: TRIMETHYLAMINE N-OXIDE REDUCTASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	123.32Å 140.70Å 59.39Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.50 19.87 – 2.40	Depositor EDS
% Data completeness (in resolution range)	90.5 (20.00-2.50) 82.6 (19.87-2.40)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.93 (at 2.41Å)	Xtrriage
Refinement program	X-PLOR 3.8	Depositor
R, $R_{free}$	0.182 , 0.247 0.209 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	14.5	Xtrriage
Anisotropy	0.533	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 61.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	6857	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	13.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.85% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 2MD, 2MO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.45	0/6422	0.66	0/8702

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6255	0	6069	53	0
2	A	94	0	44	4	0
3	A	3	0	0	5	0
4	A	505	0	0	6	0
All	All	6857	0	6113	57	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (57) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:392:PRO:HG2	1:A:395:ASP:HB2	1.57	0.87
1:A:5:ASN:N	1:A:24:LYS:HG3	1.92	0.84
3:A:801:2MO:OT2	3:A:801:2MO:MO	1.57	0.75
1:A:657:HIS:CE1	2:A:800:2MD:S13	2.81	0.73
3:A:801:2MO:OT2	3:A:801:2MO:OT1	2.07	0.72
1:A:663:SER:HB3	4:A:826:HOH:O	1.91	0.70
3:A:801:2MO:MO	3:A:801:2MO:OT1	1.64	0.69
1:A:422:ILE:HG22	1:A:429:VAL:HB	1.74	0.68
1:A:58:ARG:HG3	1:A:58:ARG:HH11	1.58	0.68
1:A:664:ARG:HH11	1:A:664:ARG:HB3	1.60	0.66
1:A:204:GLU:HG3	1:A:502:ARG:HH12	1.64	0.62
1:A:190:PRO:O	1:A:194:LEU:HB2	2.06	0.56
1:A:550:LEU:O	1:A:553:GLU:HG3	2.05	0.56
1:A:112:HIS:HB3	1:A:437:MET:HG2	1.88	0.55
1:A:15:HIS:HE1	1:A:119:ARG:O	1.89	0.54
1:A:657:HIS:NE2	2:A:800:2MD:S13	2.81	0.54
1:A:633:ARG:HB3	1:A:637:GLY:HA3	1.90	0.54
1:A:187:SER:O	1:A:327:GLY:HA3	2.08	0.54
1:A:678:VAL:HG22	1:A:707:ALA:CB	2.39	0.53
1:A:114:GLY:HA3	1:A:143:LYS:HE3	1.88	0.53
1:A:657:HIS:HE1	2:A:800:2MD:S13	2.32	0.53
1:A:57:VAL:HG12	1:A:483:PRO:HB3	1.89	0.52
1:A:678:VAL:HG22	1:A:707:ALA:HB1	1.94	0.50
1:A:757:LEU:HD12	1:A:757:LEU:O	2.11	0.50
1:A:328:TRP:O	1:A:331:GLN:HG3	2.12	0.49
2:A:800:2MD:S13	3:A:801:2MO:OT2	2.70	0.49
1:A:723:HIS:HD2	4:A:1216:HOH:O	1.95	0.49
1:A:334:GLN:HE22	1:A:631:THR:H	1.61	0.48
1:A:124:LEU:O	1:A:130:HIS:HE1	1.96	0.48
1:A:665:GLU:H	1:A:665:GLU:CD	2.17	0.48
1:A:113:ALA:HB1	1:A:131:MET:HE1	1.95	0.47
1:A:152:ALA:HB1	1:A:337:GLU:HG3	1.97	0.47
1:A:648:GLN:NE2	1:A:721:ARG:HE	2.12	0.47
1:A:503:GLY:HA2	1:A:578:PHE:CD2	2.50	0.46
1:A:257:ALA:O	1:A:261:ILE:HG12	2.16	0.46
1:A:282:VAL:HB	1:A:283:PRO:HD3	1.97	0.46
1:A:759:ILE:HD13	1:A:759:ILE:HA	1.84	0.46
1:A:716:PRO:HG2	1:A:719:ILE:HD12	1.97	0.45
1:A:113:ALA:HB1	1:A:131:MET:CE	2.46	0.45
1:A:133:ARG:O	1:A:137:MET:HG3	2.16	0.45
1:A:149:SER:OG	3:A:801:2MO:OT1	2.36	0.44
1:A:204:GLU:HG3	1:A:502:ARG:HH22	1.82	0.44

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:502:ARG:NE	4:A:955:HOH:O	2.51	0.43
1:A:312:ASP:OD1	1:A:315:LYS:HE3	2.18	0.43
1:A:678:VAL:HG13	1:A:722:ILE:HG12	1.99	0.43
1:A:622:LYS:HG2	4:A:948:HOH:O	2.19	0.42
1:A:258:HIS:CE1	1:A:295:THR:HG22	2.54	0.42
1:A:58:ARG:NH1	4:A:833:HOH:O	2.52	0.42
1:A:202:THR:O	1:A:203:HIS:HB2	2.19	0.42
1:A:680:ILE:O	1:A:711:VAL:HA	2.20	0.42
1:A:143:LYS:HE2	1:A:144:LYS:H	1.84	0.42
1:A:58:ARG:HH11	1:A:58:ARG:CG	2.28	0.42
1:A:269:LYS:HB2	1:A:269:LYS:NZ	2.36	0.41
1:A:675:ARG:N	1:A:675:ARG:HD2	2.35	0.41
1:A:99:LEU:HB3	1:A:138:HIS:CE1	2.56	0.41
1:A:53:ASN:HD21	1:A:470:ASN:HD21	1.70	0.40
1:A:641:ASP:HB2	4:A:1029:HOH:O	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	792/829 (96%)	749 (95%)	40 (5%)	3 (0%)	34 54

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	664	ARG
1	A	660	MET
1	A	388	GLU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	665/692 (96%)	635 (96%)	30 (4%)	27 51

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	23	ARG
1	A	51	VAL
1	A	59	TYR
1	A	64	LEU
1	A	68	LEU
1	A	81	PHE
1	A	115	GLN
1	A	119	ARG
1	A	126	SER
1	A	131	MET
1	A	143	LYS
1	A	144	LYS
1	A	194	LEU
1	A	204	GLU
1	A	217	LYS
1	A	252	LEU
1	A	411	TRP
1	A	416	LEU
1	A	425	ASN
1	A	443	ASN
1	A	496	TYR
1	A	553	GLU
1	A	591	ARG
1	A	595	GLU
1	A	631	THR
1	A	664	ARG
1	A	665	GLU
1	A	675	ARG
1	A	678	VAL
1	A	761	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	15	HIS
1	A	71	HIS
1	A	115	GLN
1	A	130	HIS
1	A	245	ASN
1	A	334	GLN
1	A	425	ASN
1	A	443	ASN
1	A	470	ASN
1	A	573	GLN
1	A	635	HIS
1	A	643	HIS
1	A	648	GLN
1	A	651	HIS
1	A	723	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	2MO	A	801	2,1	0,2,2	-	-	-		
2	2MD	A	799	3	42,52,52	2.23	8 (19%)	47,81,81	2.11	14 (29%)
2	2MD	A	800	3	42,52,52	2.21	9 (21%)	47,81,81	2.68	16 (34%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	2MD	A	799	3	-	2/18/66/66	0/6/6/6
2	2MD	A	800	3	-	3/18/66/66	0/6/6/6

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	800	2MD	O14-C15	8.22	1.39	1.23
2	A	799	2MD	O14-C15	8.06	1.38	1.23
2	A	799	2MD	C7-C14	5.92	1.58	1.53
2	A	800	2MD	C7-C14	5.17	1.57	1.53
2	A	799	2MD	C16-C20	4.93	1.47	1.38
2	A	800	2MD	C11-C12	4.64	1.57	1.53
2	A	800	2MD	C16-C20	4.56	1.46	1.38
2	A	800	2MD	O4'-C1'	3.99	1.46	1.41
2	A	799	2MD	C11-C12	3.67	1.56	1.53
2	A	799	2MD	C6-N1	3.26	1.42	1.37
2	A	799	2MD	C12-C13	-3.23	1.34	1.51
2	A	800	2MD	C6-N1	2.83	1.42	1.37
2	A	799	2MD	O11-C7	-2.58	1.40	1.43
2	A	800	2MD	O11-C7	-2.21	1.40	1.43
2	A	799	2MD	C2'-C1'	-2.10	1.50	1.53
2	A	800	2MD	C12-C13	-2.08	1.40	1.51
2	A	800	2MD	O11-C11	-2.03	1.39	1.44

All (30) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	800	2MD	C11-O11-C7	10.24	126.41	112.52
2	A	800	2MD	O11-C7-N8	7.23	115.99	108.57
2	A	799	2MD	C10-C11-C12	5.31	123.23	112.14
2	A	800	2MD	C17-N18-C20	5.14	122.71	113.43

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	799	2MD	C17-N18-C20	5.02	122.49	113.43
2	A	799	2MD	C11-C12-S12	4.94	122.17	110.16
2	A	800	2MD	C10-C11-C12	4.51	121.56	112.14
2	A	800	2MD	O11-C11-C10	-4.19	98.20	106.67
2	A	799	2MD	O3A-C10-C11	3.95	122.57	108.99
2	A	800	2MD	O14-C15-C16	-3.90	118.30	127.24
2	A	800	2MD	O4'-C1'-C2'	-3.62	101.64	106.93
2	A	800	2MD	O11-C7-C14	-3.52	106.56	108.97
2	A	799	2MD	O11-C7-C14	3.47	111.33	108.97
2	A	799	2MD	O4'-C1'-C2'	-3.35	102.03	106.93
2	A	799	2MD	O14-C15-C16	-3.33	119.61	127.24
2	A	799	2MD	C7-C14-N15	3.15	115.17	108.76
2	A	799	2MD	C17-N17-C15	-3.03	119.57	125.10
2	A	800	2MD	C16-C15-N17	2.95	121.08	112.31
2	A	799	2MD	C16-C15-N17	2.90	120.95	112.31
2	A	800	2MD	O3A-C10-C11	2.87	118.88	108.99
2	A	799	2MD	PA-O3B-PB	-2.86	123.01	132.83
2	A	799	2MD	O11-C11-C10	-2.83	100.96	106.67
2	A	800	2MD	PA-O3B-PB	-2.76	123.34	132.83
2	A	799	2MD	O6-C6-C5	2.76	129.76	124.37
2	A	800	2MD	O6-C6-C5	2.62	129.50	124.37
2	A	800	2MD	N16-C17-N17	2.49	122.02	116.71
2	A	800	2MD	C15-C16-N15	2.28	122.89	116.76
2	A	800	2MD	C17-N17-C15	-2.15	121.17	125.10
2	A	799	2MD	O6-C6-N1	-2.03	118.25	120.65
2	A	800	2MD	N17-C17-N18	-2.00	119.59	123.32

There are no chirality outliers.

All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	799	2MD	O3A-C10-C11-C12
2	A	800	2MD	O3A-C10-C11-O11
2	A	800	2MD	O3A-C10-C11-C12
2	A	799	2MD	O4'-C4'-C5'-O5'
2	A	800	2MD	C10-O3A-PB-O1B


There are no ring outliers.

2 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	801	2MO	5	0
2	A	800	2MD	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	794/829 (95%)	-0.34	5 (0%) 89 90	2, 11, 29, 64	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	5	ASN	4.9
1	A	7	ASP	2.8
1	A	6	GLU	2.6
1	A	395	ASP	2.4
1	A	559	ALA	2.3

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.


### 6.4 Ligands [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	2MD	A	799	47/47	0.94	0.11	1,9,13,23	0
2	2MD	A	800	47/47	0.94	0.12	3,11,16,24	0
3	2MO	A	801	3/3	0.98	0.22	16,16,18,20	0



The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.5 Other polymers [i](#)

There are no such residues in this entry.