

Full wwPDB EM Validation Report (i)

Dec 12, 2022 – 12:53 am GMT

PDB ID : 6TM2

EMDB ID : EMD-10517

Title: Human MUC2 AAs 21-1397

Authors: Javitt, G.; Khmelnitsky, L.; Albert, L.; Elad, N.; Ilani, T.; Diskin, R.; Fass,

D.

Deposited on : 2019-12-03

Resolution : 2.95 Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/EMValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev43

Mogul : 1.8.4, CSD as541be (2020)

MolProbity: 4.02b-467

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

MapQ : 1.9.9

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.31.3

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $ELECTRON\ MICROSCOPY$


The reported resolution of this entry is 2.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# ext{Entries})$	${ m EM~structures} \ (\#{ m Entries})$
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion <40%). The numeric value is given above the bar.





2 Entry composition (i)

There are 7 unique types of molecules in this entry. The entry contains 18418 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Mucin-2.

\mathbf{Mol}	Chain	Residues	Atoms				AltConf	Trace		
1	A	708	Total 5276	C 3278		O 1013	S 67	3	0	
1	В	708	Total 5276	C 3278		O 1013	S 67	3	0	

• Molecule 2 is a protein called Mucin-2.

Mol	Chain	Residues	Atoms			AltConf	Trace		
2	С	407	Total 3153	C 1961	N 541	O 609	S 42	1	0
2	D	407	Total 3153	C 1961	N 541	O 609	S 42	1	0

• Molecule 3 is a protein called Mucin-2.

Mol	Chain	Residues	Atoms			AltConf	Trace		
2	E	90	Total	С	N	О	S	0	0
3	E	90	696	429	120	138	9	0	
2	Г	00	Total	С	N	О	S	0	0
3	3 F	90	696	429	120	138	9	0	

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
Е	1325	THR	PRO	conflict	UNP Q02817
Е	1398	HIS	-	expression tag	UNP Q02817
Е	1399	HIS	-	expression tag	UNP Q02817
Е	1400	HIS	-	expression tag	UNP Q02817
E	1401	HIS	-	expression tag	UNP Q02817
Е	1402	HIS	-	expression tag	UNP Q02817
E	1403	HIS	-	expression tag	UNP Q02817
F	1325	THR	PRO	conflict	UNP Q02817
F	1398	HIS	-	expression tag	UNP Q02817



Chain	Residue	Modelled	Actual	Comment	Reference
F	1399	HIS	-	expression tag	UNP Q02817
F	1400	HIS	-	expression tag	UNP Q02817
F	1401	HIS	-	expression tag	UNP Q02817
F	1402	HIS	-	expression tag	UNP Q02817
F	1403	HIS	-	expression tag	UNP Q02817

 \bullet Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-a cetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms	AltConf	Trace	
4	G	2	Total C N		0	0
			28 16 2			
1	Н	9	Total C N	O	0	0
4	11	<u> </u>	28 16 2	10	U	U

• Molecule 5 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	AltConf
5	A	2	Total Ca 2 2	0
5	В	2	Total Ca 2 2	0
5	С	1	Total Ca 1 1	0
5	D	1	Total Ca 1 1	0
5	E	2	Total Ca 2 2	0
5	F	2	Total Ca 2 2	0

 \bullet Molecule 6 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $\rm C_8H_{15}NO_6).$





Mol	Chain	Residues	Atoms	AltConf
6	Λ	1	Total C N O	0
0	A	1	28 16 2 10	0
6	Λ	1	Total C N O	0
0	A	A 1	28 16 2 10	
6	В	1	Total C N O	0
0	Б	1	28 16 2 10	0
6	В	1	Total C N O	0
0	Б	1	28 16 2 10	U


• Molecule 7 is water.

Mol	Chain	Residues	Atoms	AltConf
7	A	10	Total O 10 10	0
7	В	10	Total O 10 10	0
7	С	12	Total O 12 12	0
7	D	12	Total O 12 12	0
7	E	1	Total O 1 1	0
7	F	1	Total O 1 1	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 2: Mucin-2







 \bullet Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G: 50% 50%



 \bullet Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain H: 50% 50%





4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	178136	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE	Depositor
	CORRECTION	
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose $(e^-/\text{Å}^2)$	30	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K3 BIOQUANTUM (6k x 4k)	Depositor
Maximum map value	2.714	Depositor
Minimum map value	-1.867	Depositor
Average map value	0.001	Depositor
Map value standard deviation	0.037	Depositor
Recommended contour level	0.17	Depositor
Map size (Å)	340.0, 340.0, 340.0	wwPDB
Map dimensions	400, 400, 400	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.85, 0.85, 0.85	Depositor



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.35	0/5406	0.61	0/7355	
1	В	0.35	0/5406	0.61	0/7355	
2	С	0.47	0/3237	0.63	0/4393	
2	D	0.47	0/3237	0.63	0/4393	
3	Е	0.31	0/715	0.56	0/970	
3	F	0.31	0/715	0.56	0/970	
All	All	0.39	0/18716	0.61	0/25436	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a maintain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	В	0	1
2	С	0	1
2	D	0	1
All	All	0	4

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	424	ASP	Peptide
1	В	424	ASP	Peptide
2	С	1126	PRO	Peptide



Mol	Chain	Res	Type	Group
2	D	1126	PRO	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5276	0	4839	33	0
1	В	5276	0	4839	35	0
2	С	3153	0	2916	18	0
2	D	3153	0	2916	17	0
3	Е	696	0	611	3	0
3	F	696	0	611	3	0
4	G	28	0	25	0	0
4	Н	28	0	25	0	0
5	A	2	0	0	0	0
5	В	2	0	0	0	0
5	С	1	0	0	0	0
5	D	1	0	0	0	0
5	Е	2	0	0	0	0
5	F	2	0	0	0	0
6	A	28	0	26	0	0
6	В	28	0	26	0	0
7	A	10	0	0	0	0
7	В	10	0	0	0	0
7	С	12	0	0	0	0
7	D	12	0	0	0	0
7	Ε	1	0	0	0	0
7	F	1	0	0	0	0
All	All	18418	0	16834	101	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (101) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



A + 1	A 4 0	Interatomic	Clash
Atom-1	Atom-2	${\rm distance} \ (\rm \mathring{A})$	overlap (Å)
1:A:35:ASN:N	1:A:172:TYR:HH	1.71	0.89
1:B:35:ASN:N	1:B:172:TYR:HH	1.73	0.87
1:B:60:ASP:OD2	1:B:201:GLN:NE2	2.22	0.72
1:A:60:ASP:OD2	1:A:201:GLN:NE2	2.21	0.72
1:B:618:THR:O	1:B:624:ASN:ND2	2.33	0.62
1:A:618:THR:O	1:A:624:ASN:ND2	2.33	0.61
1:B:640:THR:HG22	1:B:645:MET:HA	1.85	0.59
1:A:640:THR:HG22	1:A:645:MET:HA	1.85	0.59
1:A:623:ASN:ND2	1:A:626:ASP:OD2	2.38	0.57
1:B:623:ASN:ND2	1:B:626:ASP:OD2	2.38	0.56
2:D:929:LEU:HD22	2:D:936:ARG:HD3	1.88	0.56
2:C:929:LEU:HD22	2:C:936:ARG:HD3	1.88	0.55
1:B:218:CYS:O	1:B:251:GLN:NE2	2.40	0.55
1:A:558:LYS:HD2	1:A:563:CYS:HB2	1.89	0.55
1:A:218:CYS:O	1:A:251:GLN:NE2	2.39	0.55
1:B:254:ARG:NH1	1:B:261:ASP:OD1	2.40	0.55
3:E:1353:GLN:NE2	3:E:1378:LEU:O	2.41	0.54
1:A:254:ARG:NH1	1:A:261:ASP:OD1	2.40	0.53
1:B:558:LYS:HD2	1:B:563:CYS:HB2	1.89	0.53
3:F:1353:GLN:NE2	3:F:1378:LEU:O	2.41	0.53
1:A:551:ALA:O	1:A:555:ASN:ND2	2.41	0.53
1:B:704:THR:HA	1:B:714:PRO:HA	1.90	0.53
1:B:551:ALA:O	1:B:555:ASN:ND2	2.41	0.53
1:B:103:ILE:HD11	1:B:126:LEU:HD21	1.91	0.52
1:A:704:THR:HA	1:A:714:PRO:HA	1.91	0.52
1:A:140:ARG:NH1	2:C:933:ASP:OD2	2.43	0.52
1:A:721:TYR:HA	1:A:726:TYR:HA	1.91	0.52
1:B:107[A]:ARG:HD2	1:B:149:ARG:HB3	1.92	0.52
1:A:107[A]:ARG:HD2	1:A:149:ARG:HB3	1.92	0.51
1:A:103:ILE:HD11	1:A:126:LEU:HD21	1.91	0.51
1:B:721:TYR:HA	1:B:726:TYR:HA	1.91	0.51
3:E:1366:LYS:O	3:E:1370:GLN:NE2	2.43	0.51
2:D:811:ARG:HH21	2:D:828:LEU:HG	1.75	0.51
1:B:140:ARG:NH1	2:D:933:ASP:OD2	2.43	0.51
2:D:1015:LEU:O	2:D:1019:ASN:ND2	2.44	0.51
2:C:811:ARG:HH21	2:C:828:LEU:HG	1.75	0.50
2:C:1015:LEU:O	2:C:1019:ASN:ND2	2.44	0.50
2:D:859:THR:OG1	2:D:979:LYS:NZ	2.44	0.50
1:A:379:ARG:HE	2:C:1008:HIS:CD2	2.31	0.49
2:D:942:ASP:OD2	2:D:945:HIS:HB3	2.13	0.49
2:D:1036:PRO:HB3	2:D:1081:SER:HB3	1.95	0.48
1:B:379:ARG:HE	2:D:1008:HIS:CD2	2.31	0.48



Atom-1	Atom-2	Interatomic	Clash
		distance (Å)	overlap (Å)
1:B:486:ARG:HE	1:B:489:SER:HA	1.78	0.48
3:F:1366:LYS:O	3:F:1370:GLN:NE2	2.46	0.48
2:C:1036:PRO:HB3	2:C:1081:SER:HB3	1.95	0.48
2:C:942:ASP:OD2	2:C:945:HIS:HB3	2.13	0.48
1:A:486:ARG:HE	1:A:489:SER:HA	1.79	0.47
1:A:284:ASN:HD21	1:A:302:LEU:HA	1.79	0.47
2:D:1150:GLY:O	2:D:1152:HIS:ND1	2.47	0.47
1:B:284:ASN:HD21	1:B:302:LEU:HA	1.79	0.47
1:B:454:LYS:HZ1	2:D:792:TYR:HE1	1.62	0.47
1:B:85:ALA:HB3	1:B:88:PRO:HD2	1.96	0.47
2:C:961:GLU:HG3	2:C:967:ILE:HG12	1.97	0.47
2:D:806:LEU:HB3	2:D:814:CYS:HB3	1.97	0.46
2:D:961:GLU:HG3	2:D:967:ILE:HG12	1.97	0.46
1:B:503:GLN:HE21	1:B:692:PHE:HD1	1.63	0.46
1:A:85:ALA:HB3	1:A:88:PRO:HD2	1.96	0.46
2:C:806:LEU:HB3	2:C:814:CYS:HB3	1.97	0.46
1:A:503:GLN:HE21	1:A:692:PHE:HD1	1.64	0.46
1:B:127:LEU:HD23	1:B:138:TYR:HD2	1.80	0.46
2:C:1150:GLY:O	2:C:1152:HIS:ND1	2.48	0.45
1:A:282:PRO:HD2	1:A:304:SER:HB2	1.99	0.45
1:A:454:LYS:HZ1	2:C:792:TYR:HE1	1.65	0.45
2:C:859:THR:OG1	2:C:979:LYS:NZ	2.44	0.45
1:A:127:LEU:HD23	1:A:138:TYR:HD2	1.80	0.45
1:B:488:SER:HB2	1:B:671:LEU:HD22	1.99	0.45
2:C:920:ILE:HD11	2:C:976:VAL:HG21	2.00	0.43
1:B:282:PRO:HD2	1:B:304:SER:HB2	1.99	0.43
1:A:488:SER:HB2	1:A:671:LEU:HD22	1.99	0.43
1:B:131:SER:OG	1:B:132:ASP:N	2.52	0.43
1:A:131:SER:OG	1:A:132:ASP:N	2.52	0.43
2:D:920:ILE:HD11	2:D:976:VAL:HG21	2.00	0.43
1:A:503:GLN:HE22	1:A:693:ALA:H	1.67	0.43
1:B:549:THR:HB	1:B:568:ASP:OD2	2.18	0.43
1:A:494:VAL:HB	1:A:502:LEU:HB2	2.01	0.42
1:B:316:GLU:OE2	2:D:890:TYR:HB3	2.19	0.42
1:A:429:LEU:HB2	1:A:448:VAL:HB	2.01	0.42
1:A:549:THR:HB	1:A:568:ASP:OD2	2.18	0.42
1:B:35:ASN:N	1:B:172:TYR:OH	2.46	0.42
1:A:316:GLU:OE2	2:C:890:TYR:HB3	2.19	0.42
1:A:534:LEU:HG	1:A:536:GLY:H	1.85	0.42
3:E:1306:SER:OG	3:E:1307:ASP:N	2.52	0.42
1:A:66:ASP:HB3	1:A:72:LYS:HG3	2.02	0.42



Atom-1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	${\rm distance}({\rm \AA})$	overlap (Å)
1:B:494:VAL:HB	1:B:502:LEU:HB2	2.01	0.42
1:B:503:GLN:HE22	1:B:693:ALA:H	1.67	0.42
1:A:171:ASP:OD2	1:A:179:SER:OG	2.38	0.42
1:B:429:LEU:HB2	1:B:448:VAL:HB	2.01	0.42
1:B:534:LEU:HG	1:B:536:GLY:H	1.85	0.41
2:C:1100:GLN:HE21	2:C:1159:TYR:HE2	1.68	0.41
3:F:1306:SER:OG	3:F:1307:ASP:N	2.52	0.41
1:A:540:LYS:HA	1:A:546:VAL:HA	2.02	0.41
1:B:66:ASP:HB3	1:B:72:LYS:HG3	2.02	0.41
1:B:171:ASP:OD2	1:B:179:SER:OG	2.38	0.41
2:D:1100:GLN:HE21	2:D:1159:TYR:HE2	1.68	0.41
2:C:969:ILE:HB	2:C:977:PHE:HB2	2.03	0.41
2:D:969:ILE:HB	2:D:977:PHE:HB2	2.03	0.41
2:D:1187:CYS:SG	2:D:1188:GLY:N	2.94	0.40
2:C:1187:CYS:SG	2:C:1188:GLY:N	2.94	0.40
1:B:198:LYS:NZ	1:B:206:CYS:O	2.52	0.40
2:C:936:ARG:HH21	2:C:951:THR:HB	1.86	0.40
1:B:188:PHE:HA	1:B:192:GLU:OE2	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	705/729 (97%)	612 (87%)	93 (13%)	0	100	100
1	В	705/729 (97%)	611 (87%)	94 (13%)	0	100	100
2	С	402/448 (90%)	362 (90%)	40 (10%)	0	100	100
2	D	402/448 (90%)	363 (90%)	39 (10%)	0	100	100
3	Е	88/206 (43%)	87 (99%)	1 (1%)	0	100	100
3	F	88/206 (43%)	87 (99%)	1 (1%)	0	100	100



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percen	tiles
All	All	2390/2766 (86%)	2122 (89%)	268 (11%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	562/620~(91%)	561 (100%)	1 (0%)	93 98
1	В	562/620 (91%)	561 (100%)	1 (0%)	93 98
2	С	356/395 (90%)	355 (100%)	1 (0%)	92 97
2	D	356/395 (90%)	355 (100%)	1 (0%)	92 97
3	E	78/189 (41%)	78 (100%)	0	100 100
3	F	78/189 (41%)	78 (100%)	0	100 100
All	All	1992/2408 (83%)	1988 (100%)	4 (0%)	93 98

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	419	LYS
1	В	419	LYS
2	С	998	ARG
2	D	998	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	221	HIS
1	A	284	ASN
1	A	352	HIS
1	A	503	GLN
1	A	652	HIS



Continued from previous page...

Mol	Chain	Res	Type
1	A	665	GLN
1	В	221	HIS
1	В	284	ASN
1	В	352	HIS
1	В	503	GLN
1	В	652	HIS
1	В	665	GLN
2	С	1008	HIS
2	D	1008	HIS
3	Е	1370	GLN
3	F	1370	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bo	Bond lengths			ond ang	les
IVIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NAG	G	1	4,2	14,14,15	0.31	0	17,19,21	0.46	0
4	NAG	G	2	4	14,14,15	0.27	0	17,19,21	0.58	1 (5%)
4	NAG	Н	1	4,2	14,14,15	0.32	0	17,19,21	0.48	0
4	NAG	Н	2	4	14,14,15	0.29	0	17,19,21	0.59	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the



Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	G	1	4,2	-	1/6/23/26	0/1/1/1
4	NAG	G	2	4	-	2/6/23/26	0/1/1/1
4	NAG	Н	1	4,2	-	1/6/23/26	0/1/1/1
4	NAG	Н	2	4	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^{o})$	$\mathbf{Ideal}(^{o})$
4	G	2	NAG	C1-O5-C5	2.03	114.94	112.19
4	Н	2	NAG	C1-O5-C5	2.02	114.92	112.19

There are no chirality outliers.

All (6) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	G	2	NAG	O5-C5-C6-O6
4	Н	2	NAG	O5-C5-C6-O6
4	G	2	NAG	C4-C5-C6-O6
4	Н	2	NAG	C4-C5-C6-O6
4	G	1	NAG	C4-C5-C6-O6
4	Н	1	NAG	C4-C5-C6-O6

There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.









5.6 Ligand geometry (i)

Of 14 ligands modelled in this entry, 10 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Во	Bond lengths			Bond angles		
WIOI	Туре	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2	
6	NAG	В	2004	1	14,14,15	0.33	0	17,19,21	0.49	0	
6	NAG	В	2003	1	14,14,15	1.33	2 (14%)	17,19,21	1.26	1 (5%)	
6	NAG	A	2004	1	14,14,15	0.34	0	17,19,21	0.49	0	



Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
IVIOI	туре	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	NAG	A	2003	1	14,14,15	1.34	2 (14%)	17,19,21	1.27	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	NAG	В	2004	1	-	4/6/23/26	0/1/1/1
6	NAG	В	2003	1	-	2/6/23/26	0/1/1/1
6	NAG	A	2004	1	-	4/6/23/26	0/1/1/1
6	NAG	A	2003	1	-	2/6/23/26	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(A)	Ideal(Å)
6	A	2003	NAG	O5-C1	4.43	1.50	1.43
6	В	2003	NAG	O5-C1	4.42	1.50	1.43
6	A	2003	NAG	C1-C2	2.24	1.55	1.52
6	В	2003	NAG	C1-C2	2.18	1.55	1.52

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$\operatorname{Ideal}({}^{o})$
6	A	2003	NAG	C1-O5-C5	5.03	119.00	112.19
6	В	2003	NAG	C1-O5-C5	5.00	118.97	112.19

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	2004	NAG	O5-C5-C6-O6
6	В	2004	NAG	O5-C5-C6-O6
6	В	2004	NAG	C4-C5-C6-O6
6	A	2003	NAG	O5-C5-C6-O6
6	В	2003	NAG	O5-C5-C6-O6
6	A	2004	NAG	C4-C5-C6-O6
6	A	2004	NAG	C8-C7-N2-C2
6	A	2004	NAG	O7-C7-N2-C2
6	В	2004	NAG	C8-C7-N2-C2



Mol	Chain	Res	Type	Atoms
6	В	2004	NAG	O7-C7-N2-C2
6	A	2003	NAG	C4-C5-C6-O6
6	В	2003	NAG	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.




6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-10517. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections (i)


6.1.1 Primary map



The images above show the map projected in three orthogonal directions.

6.2 Central slices (i)

6.2.1 Primary map





The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices (i)


6.3.1 Primary map



The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal surface views (i)

6.4.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.17. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.



6.5 Mask visualisation (i)


This section was not generated. No masks/segmentation were deposited.



7 Map analysis (i)

This section contains the results of statistical analysis of the map.


7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.



7.2 Volume estimate (i)




The volume at the recommended contour level is $133~\mathrm{nm}^3$; this corresponds to an approximate mass of $120~\mathrm{kDa}$.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



7.3 Rotationally averaged power spectrum (i)




^{*}Reported resolution corresponds to spatial frequency of 0.339 $\rm \AA^{-1}$



8 Fourier-Shell correlation (i)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC (i)



*Reported resolution corresponds to spatial frequency of 0.339 $\rm \mathring{A}^{-1}$



8.2 Resolution estimates (i)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
resolution estimate (A)	0.143	0.5	Half-bit
Reported by author	2.95	-	-
Author-provided FSC curve	3.68	4.36	3.79
Unmasked-calculated*	-	-	-

^{*}Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from author-provided FSC intersecting FSC 0.143 CUT-OFF 3.68 differs from the reported value 2.95 by more than 10 %




9 Map-model fit (i)


This section contains information regarding the fit between EMDB map EMD-10517 and PDB model 6TM2. Per-residue inclusion information can be found in section 3 on page 6.

9.1 Map-model overlays

9.1.1 Map-model overlay (i)




9.1.2 Map-model assembly overlay (i)



The images above show the 3D surface view of the map at the recommended contour level 0.17 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.




9.2 Q-score mapped to coordinate model (i)



The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.


9.3 Atom inclusion mapped to coordinate model (i)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.17).



9.4 Atom inclusion (i)



At the recommended contour level, 87% of all backbone atoms, 86% of all non-hydrogen atoms, are inside the map.



9.5 Map-model fit summary (i)

The table lists the average atom inclusion at the recommended contour level (0.17) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.8588	0.5490
A	0.8297	0.5300
В	0.8299	0.5300
С	0.9313	0.5890
D	0.9316	0.5880
E	0.7889	0.5140
F	0.7889	0.5130
G	0.8214	0.5480
Н	0.8214	0.5450



