

Full wwPDB X-ray Structure Validation Report (i)

Jan 14, 2024 - 10:34 am GMT

PDB ID : 6TM3

Title : Structure of methylene-tetrahydromethanopterin dehydrogenase from Methy-

lorubrum extorquens AM1 in a close conformation containing NADP+ and

methylene-H4MPT

Authors: Wagner, T.; Huang, G.; Demmer, U.; Warkentin, E.; Ermler, U.; Shima, S.

Deposited on : 2019-12-03

Resolution : 1.08 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.4, CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.36

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 1.08 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# \mathrm{Entries})$	$(\# ext{Entries}, ext{ resolution range}(ext{Å}))$
R_{free}	130704	1386 (1.12-1.04)
Clashscore	141614	1021 (1.10-1.06)
Ramachandran outliers	138981	1381 (1.12-1.04)
Sidechain outliers	138945	1379 (1.12-1.04)
RSRZ outliers	127900	1359 (1.12-1.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	А	303	91%	• 5%



2 Entry composition (i)

There are 7 unique types of molecules in this entry. The entry contains 2939 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Bifunctional protein MdtA.


Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf	Trace
1	A	289	Total 2295	C 1465	N 387	O 435	S 8	0	31	0

There are 15 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	289	VAL	-	expression tag	UNP P55818
A	290	ASP	-	expression tag	UNP P55818
A	291	LYS	-	expression tag	UNP P55818
A	292	LEU	-	expression tag	UNP P55818
A	293	ALA	-	expression tag	UNP P55818
A	294	ALA	-	expression tag	UNP P55818
A	295	ALA	-	expression tag	UNP P55818
A	296	LEU	-	expression tag	UNP P55818
A	297	GLU	-	expression tag	UNP P55818
A	298	HIS	-	expression tag	UNP P55818
A	299	HIS	-	expression tag	UNP P55818
A	300	HIS	-	expression tag	UNP P55818
A	301	HIS	-	expression tag	UNP P55818
A	302	HIS	=	expression tag	UNP P55818
A	303	HIS	-	expression tag	UNP P55818


• Molecule 2 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NAP) (formula: C₂₁H₂₈N₇O₁₇P₃) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf
2	A	1	Total 48	C 21	_	O 17	P 3	0	0


• Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 6 3 3	0	1


 \bullet Molecule 4 is 5,10-DIMETHYLENE TETRAHYDROMETHANOPTERIN (three-letter code: H4M) (formula: C $_{31}H_{45}N_6O_{16}P)$ (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	A	Aton	ns		ZeroOcc	AltConf
4	٨	1	Total	С	N	О	0	1
4	A	1	31	21	6	4	U	1

 \bullet Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $\mathrm{C_2H_6O_2}).$



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 4 2 2	0	1
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0

Continued on next page...



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 4 2 2	0	1
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	1
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	1
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	1
5	A	1	Total C O 4 2 2	0	0

• Molecule 6 is LITHIUM ION (three-letter code: LI) (formula: Li).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	2	Total Li 2 2	0	0

• Molecule 7 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	501	Total O 501 501	0	50



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Bifunctional protein MdtA





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 41 3 2	Depositor
Cell constants	125.75Å 125.75Å 125.75Å	Donogitor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	41.92 - 1.08	Depositor
Resolution (A)	41.92 - 1.08	EDS
% Data completeness	99.3 (41.92-1.08)	Depositor
(in resolution range)	99.5 (41.92-1.08)	EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.05	Depositor
$< I/\sigma(I) > 1$	1.37 (at 1.08Å)	Xtriage
Refinement program	PHENIX 1.14_3260	Depositor
D D.	0.113 , 0.127	Depositor
R, R_{free}	0.129 , 0.142	DCC
R_{free} test set	7185 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	11.8	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.36, 50.5	EDS
L-test for twinning ²	$ < L > = 0.49, < L^2> = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	2939	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 2.96% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, LI, H4M, NAP, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
		RMSZ	# Z > 5	RMSZ	# Z > 5	
	1	A	0.35	0/2426	0.64	0/3257

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2295	0	2407	13	0
2	A	48	0	25	3	0
3	A	6	0	8	0	0
4	A	31	0	27	3	0
5	A	56	0	84	4	0
6	A	2	0	0	0	0
7	A	501	0	0	2	0
All	All	2939	0	2551	14	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (14) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
1:A:279[A]:GLU:HB3	5:A:407[A]:EDO:H11	1.64	0.79
1:A:279[A]:GLU:CB	5:A:407[A]:EDO:H11	2.22	0.68
1:A:78[B]:GLU:HG2	7:A:797:HOH:O	1.96	0.64
2:A:401:NAP:C4N	4:A:403[B]:H4M:H16	2.31	0.61
1:A:221:TYR:O	2:A:401:NAP:H2N	2.05	0.56
1:A:257[B]:LEU:HD21	1:A:261[B]:ARG:NH1	2.22	0.55
1:A:279[A]:GLU:CG	5:A:407[A]:EDO:H11	2.38	0.53
1:A:4:LYS:HD2	7:A:630[B]:HOH:O	2.13	0.47
1:A:102[A]:THR:OG1	2:A:401:NAP:H4N	2.16	0.46
1:A:102[A]:THR:HG22	1:A:221:TYR:CE1	2.54	0.43
1:A:82:LYS:HD3	5:A:411[A]:EDO:O2	2.18	0.43
1:A:18:PHE:CZ	4:A:403[B]:H4M:H15	2.54	0.42
1:A:18:PHE:CE2	4:A:403[B]:H4M:H15	2.55	0.42
1:A:113:ALA:O	1:A:245[A]:ARG:NH2	2.53	0.41

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	321/303 (106%)	315 (98%)	6 (2%)	0	100 100	

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Analysed Rotameric Outlie		Percentiles	
1	A	240/217 (111%)	239 (100%)	1 (0%)	91 74	

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	256	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 19 ligands modelled in this entry, 2 are monoatomic - leaving 17 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Trino	Chain	Res	Link	Bond lengths			Bond angles		
MIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	EDO	A	417	-	3,3,3	0.06	0	2,2,2	0.18	0
5	EDO	A	415	-	3,3,3	0.41	0	2,2,2	0.42	0
5	EDO	A	406	-	3,3,3	0.44	0	2,2,2	0.55	0
3	GOL	A	402[A]	-	5,5,5	0.69	0	5,5,5	0.84	0
5	EDO	A	404[A]	-	3,3,3	0.40	0	2,2,2	0.27	0



Mal	Trino	Chain	Res	T inle	Во	ond leng	ths	В	ond ang	les
Mol	Type	Chain	nes	Link	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2
5	EDO	A	413[B]	-	3,3,3	0.48	0	2,2,2	0.30	0
5	EDO	A	414	-	3,3,3	0.47	0	2,2,2	0.41	0
5	EDO	A	412	-	3,3,3	0.48	0	2,2,2	0.23	0
5	EDO	A	405	-	3,3,3	0.42	0	2,2,2	0.63	0
5	EDO	A	416[B]	-	3,3,3	0.48	0	2,2,2	0.37	0
4	H4M	A	403[B]	-	32,34,58	1.36	1 (3%)	38,50,86	1.50	6 (15%)
5	EDO	A	408	-	3,3,3	0.38	0	2,2,2	0.57	0
5	EDO	A	411[A]	-	3,3,3	0.43	0	2,2,2	0.63	0
5	EDO	A	410	-	3,3,3	0.43	0	2,2,2	0.26	0
5	EDO	A	409	-	3,3,3	0.49	0	2,2,2	0.28	0
5	EDO	A	407[A]	-	3,3,3	0.47	0	2,2,2	0.18	0
2	NAP	A	401	-	45,52,52	1.10	3 (6%)	56,80,80	1.14	5 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	A	417	-	-	1/1/1/1	-
5	EDO	A	415	-	-	0/1/1/1	-
5	EDO	A	406	-	-	0/1/1/1	-
3	GOL	A	402[A]	-	-	0/4/4/4	-
5	EDO	A	404[A]	-	-	0/1/1/1	-
5	EDO	A	413[B]	-	-	1/1/1/1	-
5	EDO	A	414	-	-	0/1/1/1	-
5	EDO	A	412	-	-	0/1/1/1	-
5	EDO	A	405	-	-	1/1/1/1	-
5	EDO	A	416[B]	-	-	0/1/1/1	-
4	H4M	A	403[B]	_	-	1/15/43/85	0/4/4/5
5	EDO	A	408	-	-	0/1/1/1	-
5	EDO	A	411[A]	-	-	0/1/1/1	-
5	EDO	A	410	_	-	1/1/1/1	-
5	EDO	A	409	-	-	0/1/1/1	-
5	EDO	A	407[A]	_	-	0/1/1/1	_
2	NAP	A	401	_	-	5/31/67/67	0/5/5/5

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	Observed(Å)	$\operatorname{Ideal}(\text{\AA})$
4	A	403[B]	H4M	C10-N5	-5.99	1.42	1.46
2	A	401	NAP	O4D-C1D	3.77	1.46	1.41

Continued on next page...



 $Continued\ from\ previous\ page...$

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	Observed(A)	$\operatorname{Ideal}(\text{\AA})$
2	A	401	NAP	C6N-N1N	3.55	1.44	1.35
2	A	401	NAP	C6N-C5N	2.35	1.43	1.38

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$Ideal(^{o})$
2	A	401	NAP	C5N-C6N-N1N	-4.37	114.14	120.40
4	A	403[B]	H4M	CX5-CX4-CX3	-4.01	105.39	113.07
2	A	401	NAP	N3A-C2A-N1A	-3.45	123.29	128.68
4	A	403[B]	H4M	C4A-C4-N3	3.27	116.76	110.99
4	A	403[B]	H4M	C10-N5-C4A	-3.04	124.69	127.47
4	A	403[B]	H4M	CX1-CX2-CX3	-2.71	108.97	113.65
4	A	403[B]	H4M	C2-N1-C8A	2.55	118.03	113.43
2	A	401	NAP	O4D-C1D-C2D	-2.40	103.42	106.93
2	A	401	NAP	C2A-N1A-C6A	2.19	122.50	118.75
4	A	403[B]	H4M	C6-C9-N10	2.16	104.18	102.56
2	A	401	NAP	N6A-C6A-N1A	2.12	122.97	118.57

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	401	NAP	O4D-C1D-N1N-C2N
2	A	401	NAP	O4D-C1D-N1N-C6N
2	A	401	NAP	C2D-C1D-N1N-C2N
2	A	401	NAP	C2D-C1D-N1N-C6N
4	A	403[B]	H4M	CX3-CX4-CX5-OX5
5	A	417	EDO	O1-C1-C2-O2
5	A	410	EDO	O1-C1-C2-O2
5	A	405	EDO	O1-C1-C2-O2
2	A	401	NAP	C2B-O2B-P2B-O2X
5	A	413[B]	EDO	O1-C1-C2-O2


There are no ring outliers.

4 monomers are involved in 9 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	403[B]	H4M	3	0
5	A	411[A]	EDO	1	0
5	A	407[A]	EDO	3	0
2	A	401	NAP	3	0



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\# \mathrm{RSRZ}{>}2$	$OWAB(Å^2)$	Q < 0.9
1	A	289/303 (95%)	-0.03	4 (1%) 75 69	10, 13, 22, 60	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	290	ASP	7.1
1	A	289	VAL	5.3
1	A	56	LYS	2.9
1	A	180	ASP	2.3

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$ m B ext{-}factors(\AA^2)$	Q<0.9
5	EDO	A	411[A]	4/4	0.30	0.27	53,54,55,55	4
5	EDO	A	416[B]	4/4	0.56	0.19	24,25,26,27	4
5	EDO	A	414	4/4	0.60	0.14	54,54,55,55	0


Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}(\mathring{\mathbf{A}}^2)$	Q < 0.9
5	EDO	A	409	4/4	0.65	0.16	55,56,57,58	0
5	EDO	A	417	4/4	0.71	0.27	41,42,45,46	0
5	EDO	A	405	4/4	0.78	0.12	38,40,41,44	0
5	EDO	A	412	4/4	0.80	0.17	43,46,48,48	0
5	EDO	A	408	4/4	0.87	0.17	36,40,41,41	0
5	EDO	A	413[B]	4/4	0.89	0.15	47,47,48,48	4
5	EDO	A	407[A]	4/4	0.89	0.17	35,39,40,40	4
5	EDO	A	415	4/4	0.92	0.09	23,27,29,29	0
5	EDO	A	410	4/4	0.94	0.12	21,30,30,30	0
6	LI	A	418	1/1	0.94	0.16	14,14,14,14	0
3	GOL	A	402[A]	6/6	0.95	0.08	17,18,19,20	6
5	EDO	A	406	4/4	0.95	0.13	19,24,27,29	0
4	H4M	A	403[B]	31/54	0.97	0.08	10,12,33,34	0
6	LI	A	419	1/1	0.97	0.60	20,20,20,20	0
5	EDO	A	404[A]	4/4	0.99	0.06	10,10,10,10	4
2	NAP	A	401	48/48	0.99	0.06	10,11,16,18	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







6.5 Other polymers (i)

There are no such residues in this entry.

