

Full wwPDB EM Validation Report (i)

May 11, 2024 – 03:34 pm BST

PDB ID : 6TNI

EMDB ID : EMD-10534

Title : Structure of FANCD2 homodimer Authors : Alcon, P.; Shakeel, S.; Passmore, L.A.

Deposited on : 2019-12-08

Resolution : 3.40 Å(reported)

Based on initial model : 3S4W

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*A user guide is available at

https://www.wwpdb.org/validation/2017/EMValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev92

MolProbity : 4.02b-467

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $MapQ \quad : \quad 1.9.13$

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36.2

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $ELECTRON\ MICROSCOPY$

The reported resolution of this entry is 3.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	${ m EM\ structures} \ (\#{ m Entries})$
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion <40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain				
1	A	1475	71%	• 28%	_		
1	a	1475	70%	• 28%			



2 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 11625 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Uncharacterized protein.

\mathbf{Mol}	Chain	Residues	Atoms				AltConf	Trace		
1	Δ	1067	Total	С	N	О	S	0	0	İ
1	Λ	1007	5815	3543	1118	1143	11		U	
1		1067	Total	С	N	O	S	0	0	
1	a	1007	5810	3540	1116	1143	11	0	U	

There are 76 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1437	ALA	GLY	conflict	UNP F1NP22
A	1439	ARG	-	expression tag	UNP F1NP22
A	1440	LEU	-	expression tag	UNP F1NP22
A	1441	GLU	-	expression tag	UNP F1NP22
A	1442	VAL	-	expression tag	UNP F1NP22
A	1443	LEU	-	expression tag	UNP F1NP22
A	1444	PHE	-	expression tag	UNP F1NP22
A	1445	GLN	-	expression tag	UNP F1NP22
A	1446	GLY	-	expression tag	UNP F1NP22
A	1447	PRO	-	expression tag	UNP F1NP22
A	1448	TRP	-	expression tag	UNP F1NP22
A	1449	SER	-	expression tag	UNP F1NP22
A	1450	HIS	-	expression tag	UNP F1NP22
A	1451	PRO	-	expression tag	UNP F1NP22
A	1452	GLN	-	expression tag	UNP F1NP22
A	1453	PHE	-	expression tag	UNP F1NP22
A	1454	GLU	-	expression tag	UNP F1NP22
A	1455	LYS	-	expression tag	UNP F1NP22
A	1456	GLY	-	expression tag	UNP F1NP22
A	1457	SER	-	expression tag	UNP F1NP22
A	1458	ALA	-	expression tag	UNP F1NP22
A	1459	GLY		expression tag	UNP F1NP22
A	1460	SER	-	expression tag	UNP F1NP22
A	1461	ALA	-	expression tag	UNP F1NP22
A	1462	ALA		expression tag	UNP F1NP22
A	1463	GLY	-	expression tag	UNP F1NP22

Continued on next page...



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	1464	SER	rictaar	expression tag	UNP F1NP22
A	1465	GLY	_	expression tag	UNP F1NP22
A	1466	ALA	_	expression tag	UNP F1NP22
A	1467	GLY	_	expression tag	UNP F1NP22
A	1468	TRP	_	expression tag	UNP F1NP22
A	1469	SER	_	expression tag	UNP F1NP22
A	1470	HIS	_	expression tag	UNP F1NP22
A	1470	PRO	-	expression tag	UNP F1NP22
A	1472	GLN	_	expression tag	UNP F1NP22
A	1473	PHE	-	expression tag	UNP F1NP22
A	1473	GLU	-	expression tag	UNP F1NP22
A	1474	LYS	-	1 0	UNP F1NP22
			CIV	expression tag	
a	1437	ALA	GLY	conflict	UNP F1NP22
a	1439	ARG	-	expression tag	UNP F1NP22
a	1440	LEU	-	expression tag	UNP F1NP22
a	1441	GLU	-	expression tag	UNP F1NP22
a	1442	VAL	-	expression tag	UNP F1NP22
a	1443	LEU	-	expression tag	UNP F1NP22
a	1444	PHE	-	expression tag	UNP F1NP22
a	1445	GLN	-	expression tag	UNP F1NP22
a	1446	GLY	-	expression tag	UNP F1NP22
a	1447	PRO	-	expression tag	UNP F1NP22
a	1448	TRP	-	expression tag	UNP F1NP22
a	1449	SER	-	expression tag	UNP F1NP22
a	1450	HIS	-	expression tag	UNP F1NP22
a	1451	PRO	-	expression tag	UNP F1NP22
a	1452	GLN	-	expression tag	UNP F1NP22
a	1453	PHE	-	expression tag	UNP F1NP22
a	1454	GLU	-	expression tag	UNP F1NP22
a	1455	LYS	-	expression tag	UNP F1NP22
a	1456	GLY	-	expression tag	UNP F1NP22
a	1457	SER	_	expression tag	UNP F1NP22
a	1458	ALA	-	expression tag	UNP F1NP22
a	1459	GLY	_	expression tag	UNP F1NP22
a	1460	SER	-	expression tag	UNP F1NP22
a	1461	ALA		expression tag	UNP F1NP22
a	1462	ALA	_	expression tag	UNP F1NP22
a	1463	GLY	-	expression tag	UNP F1NP22
a	1464	SER		expression tag	UNP F1NP22
a	1465	GLY	-	expression tag	UNP F1NP22
a	1466	ALA	-	expression tag	UNP F1NP22
a	1467	GLY		expression tag	UNP F1NP22

Continued on next page...




Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
a	1468	TRP	-	expression tag	UNP F1NP22
a	1469	SER	-	expression tag	UNP F1NP22
a	1470	HIS	-	expression tag	UNP F1NP22
a	1471	PRO	-	expression tag	UNP F1NP22
a	1472	GLN	-	expression tag	UNP F1NP22
a	1473	PHE	-	expression tag	UNP F1NP22
a	1474	GLU	-	expression tag	UNP F1NP22
a	1475	LYS	-	expression tag	UNP F1NP22



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Uncharacterized protein Chain A: LINE SERVICE OF THE S THE STATE OF




4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	901085	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE	Depositor
	CORRECTION	
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose $(e^-/\text{Å}^2)$	70.0	Depositor
Minimum defocus (nm)	1800	Depositor
Maximum defocus (nm)	3500	Depositor
Magnification	81000	Depositor
Image detector	GATAN K3 BIOQUANTUM (6k x 4k)	Depositor
Maximum map value	0.097	Depositor
Minimum map value	-0.062	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.001	Depositor
Recommended contour level	0.0134	Depositor
Map size (Å)	476.99997, 476.99997, 476.99997	wwPDB
Map dimensions	450, 450, 450	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.06, 1.06, 1.06	Depositor



5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain		nd lengths	Bond	angles
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z >5
1	A	0.38	1/5835~(0.0%)	0.58	0/8080
1	a	0.39	$1/5829 \ (0.0\%)$	0.57	0/8072
All	All	0.38	$2/11664 \ (0.0\%)$	0.58	0/16152

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(A)	$\operatorname{Ideal}(\text{\AA})$
1	A	921	ARG	C-N	-5.56	1.23	1.34
1	a	921	ARG	C-N	-5.53	1.23	1.34

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

Due to software issues we are unable to calculate clashes - this section is therefore empty.


5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	\mathbf{ntiles}
1	A	1053/1475 (71%)	893 (85%)	155 (15%)	5 (0%)	29	61





Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	a	1053/1475 (71%)	883 (84%)	165 (16%)	5 (0%)	29	61
All	All	2106/2950 (71%)	1776 (84%)	320 (15%)	10 (0%)	32	61

All (10) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	438	ILE
1	A	567	SER
1	A	569	VAL
1	a	435	PHE
1	a	917	LEU
1	A	917	LEU
1	a	425	GLN
1	a	437	SER
1	A	761	PRO
1	a	761	PRO

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	3
1	A	170/1311 (13%)	150 (88%)	20 (12%)	5 19	
1	a	169/1311 (13%)	146 (86%)	23 (14%)	3 14	
All	All	339/2622 (13%)	296 (87%)	43 (13%)	8 16	

All (43) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	357	VAL
1	A	358	ARG
1	A	375	THR
1	A	378	SER
1	A	382	VAL
1	A	393	SER

Continued on next page...



Continued from previous page...

Continued from previous page					
Mol	Chain	Res	Type		
1	A	414	CYS		
1	A	420	MET		
1	A	429	MET		
1	A	458	SER		
1	A	461	TYR		
1	A	472	CYS		
1	A	483	HIS		
1	A	493	ASP		
1	A	511	LEU		
1	A	512	ARG		
1	A	533	ARG		
1	A	536	PHE		
1	A	572	HIS		
1	A	635	ASP		
1	a	357	VAL		
1	a	358	ARG		
1	a	375	THR		
1	a	377	VAL		
1	a	382	VAL		
1	a	388	LEU		
1	a	414	CYS		
1	a	420	MET		
1	a	421	GLN		
1	a	425	GLN		
1	a	429	MET		
1	a	458	SER		
1	a	472	CYS		
1	a	484	VAL		
1	a	493	ASP		
1	a	511	LEU		
1	a	512	ARG		
1	a	533	ARG		
1	a	536	PHE		
1	a	563	LYS		
1	a	572	HIS		
1	a	635	ASP		
1	a	653	LEU		

Sometimes side chains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such side chains are listed below:

\mathbf{Mol}	Chain	Res	Type
1	A	392	HIS

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type
1	A	418	GLN
1	A	427	HIS
1	A	447	HIS
1	A	483	HIS
1	a	392	HIS
1	a	418	GLN
1	a	483	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.




6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-10534. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections (i)

6.1.1 Primary map



6.1.2 Raw map



The images above show the map projected in three orthogonal directions.



6.2 Central slices (i)

6.2.1 Primary map







Y Index: 225




Z Index: 225


6.2.2 Raw map



X Index: 225



Y Index: 225



Z Index: 225


The images above show central slices of the map in three orthogonal directions.




6.3 Largest variance slices (i)

6.3.1 Primary map








X Index: 232

Raw map

6.3.2







X Index: 218

Y Index: 189


Z Index: 232

The images above show the largest variance slices of the map in three orthogonal directions.




6.4 Orthogonal standard-deviation projections (False-color) (i)

6.4.1 Primary map



6.4.2 Raw map




The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.




6.5 Orthogonal surface views (i)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.0134. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.

6.6 Mask visualisation (i)


This section was not generated. No masks/segmentation were deposited.



7 Map analysis (i)

This section contains the results of statistical analysis of the map.


7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.



7.2 Volume estimate (i)




The volume at the recommended contour level is $117~\mathrm{nm^3}$; this corresponds to an approximate mass of $106~\mathrm{kDa}$.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



7.3 Rotationally averaged power spectrum (i)




^{*}Reported resolution corresponds to spatial frequency of 0.294 $\rm \mathring{A}^{-1}$



8 Fourier-Shell correlation (i)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC (i)



*Reported resolution corresponds to spatial frequency of 0.294 $\rm \mathring{A}^{-1}$



8.2 Resolution estimates (i)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
resolution estimate (A)	0.143	0.5	Half-bit
Reported by author	3.40	-	-
Author-provided FSC curve	3.43	3.85	3.51
Unmasked-calculated*	3.99	5.84	4.03


^{*}Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.99 differs from the reported value 3.4 by more than 10 %



9 Map-model fit (i)

This section contains information regarding the fit between EMDB map EMD-10534 and PDB model 6TNI. Per-residue inclusion information can be found in section 3 on page 6.


9.1 Map-model overlay (i)



The images above show the 3D surface view of the map at the recommended contour level 0.0134 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.




9.2 Q-score mapped to coordinate model (i)



The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.


9.3 Atom inclusion mapped to coordinate model (i)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.0134).



9.4 Atom inclusion (i)



At the recommended contour level, 83% of all backbone atoms, 79% of all non-hydrogen atoms, are inside the map.



9.5 Map-model fit summary (i)

The table lists the average atom inclusion at the recommended contour level (0.0134) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.7920	0.3830
A	0.7940	0.3830
a	0.7890	0.3840



