

Full wwPDB X-ray Structure Validation Report (i)

Dec 15, 2020 – 10:18 am GMT

PDB ID : 6TP5

Title: Crystal structure of human Transmembrane prolyl 4-hydroxylase

Authors: Myllykoski, M.; Sutinen, A.; Koski, M.K.; Kallio, J.P.; Raasakka, A.; Mylly-

harju, J.; Wierenga, R.K.; Koivunen, P.

Deposited on : 2019-12-12

Resolution : 2.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul: 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.15.1 buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac: 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.15.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution			
Metric	$(\# ext{Entries})$	$(\# ext{Entries}, ext{resolution range}(ext{Å}))$			
R_{free}	130704	1377 (2.26-2.26)			
Clashscore	141614	1487 (2.26-2.26)			
Ramachandran outliers	138981	1449 (2.26-2.26)			
Sidechain outliers	138945	1450 (2.26-2.26)			
RSRZ outliers	127900	1356 (2.26-2.26)			

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain						
1	A	421	3%	78%	10%	12%			
1	В	421	2%	70%	15%	15%			
2	С	2	50%		50%				
2	Е	2		100%					
3	D	10	30%	70%					



Continued from previous page...

Mol	Chain	Length		Quality of chain
4	F	10	20%	80%



2 Entry composition (i)

There are 11 unique types of molecules in this entry. The entry contains 6394 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Transmembrane prolyl 4-hydroxylase.

Mol	Chain	Residues	${f Atoms}$					ZeroOcc	AltConf	Trace
1	A	370	Total 2995	C 1878	N 544	O 559	S 14	0	0	0
1	В	356	Total 2898	C 1821	N 529	O 534	S 14	0	2	0

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	82	HIS	-	expression tag	UNP Q9NXG6
A	83	HIS	-	expression tag	UNP Q9NXG6
A	84	HIS	_	expression tag	UNP Q9NXG6
A	85	HIS	_	expression tag	UNP Q9NXG6
A	86	HIS	_	expression tag	UNP Q9NXG6
A	87	HIS	_	expression tag	UNP Q9NXG6
A	99	VAL	ALA	conflict	UNP Q9NXG6
В	82	HIS	_	expression tag	UNP Q9NXG6
В	83	HIS	_	expression tag	UNP Q9NXG6
В	84	HIS	_	expression tag	UNP Q9NXG6
В	85	HIS	_	expression tag	UNP Q9NXG6
В	86	HIS	-	expression tag	UNP Q9NXG6
В	87	HIS	_	expression tag	UNP Q9NXG6
В	99	VAL	ALA	conflict	UNP Q9NXG6

• Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-a cetamido-2-deoxy-beta-D-glucopyranose.






Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	С	2	Total 28	C 16		0	0	0
2	E	2	Total		N	 0	0	0

• Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyran ose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
3	D	10	Total C N O 116 64 2 50	0	0	0


• Molecule 4 is an oligosaccharide called beta-D-mannopyranose-(1-2)-alpha-D-mannopyranose e-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
4	F	10	Total C N O 116 64 2 50	0	0	0

• Molecule 5 is N-OXALYLGLYCINE (three-letter code: OGA) (formula: C₄H₅NO₅) (labeled as "Ligand of Interest" by depositor).






Mol	Chain	Residues	A	ton	ns		ZeroOcc	AltConf	
5	Λ	1	Total	С	N	О	0	0	
0	9 A	1	10	4	1	5	0		
5	D	1	Total	С	Ν	О	0	0	
0	Б	1	10	4	1	5	0		

• Molecule 6 is FE (II) ION (three-letter code: FE2) (formula: Fe) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	В	1	Total Fe 1 1	0	0
6	A	1	Total Fe 1 1	0	0

 \bullet Molecule 7 is TERTIARY-BUTYL ALCOHOL (three-letter code: TBU) (formula: $\mathrm{C_4H_{10}O}).$






Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total C O 5 4 1	0	0
7	A	1	Total C O 5 4 1	0	0
7	A	1	Total C O 5 4 1	0	0
7	A	1	Total C O 5 4 1	0	0
7	В	1	Total C O 5 4 1	0	0
7	В	1	Total C O 5 4 1	0	0
7	В	1	Total C O 5 4 1	0	0
7	В	1	Total C O 5 4 1	0	0

 \bullet Molecule 8 is GLYCINE (three-letter code: GLY) (formula: $\mathrm{C_2H_5NO_2}).$





Mol	Chain	Residues	${f Atoms}$				ZeroOcc	AltConf
8	A	1	Total 5				0	0
8	A	1	Total 5	C 2	N 1	O 2	0	0

• Molecule 9 is CALCIUM ION (three-letter code: CA) (formula: Ca) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	${f Atoms}$	ZeroOcc	AltConf
9	В	2	Total Ca 2 2	0	0
9	A	2	Total Ca 2 2	0	0

• Molecule 10 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

\mathbf{Mol}	Chain	Residues	${f Atoms}$	ZeroOcc	AltConf
10	A	2	Total Cl 2 2	0	0

• Molecule 11 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
11	A	67	Total O 67 67	0	0
11	В	68	Total O 68 68	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Transmembrane prolyl 4-hydroxylase



• Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain C: 50% 50%





• Molecule 2:	2-acetamido-2-deoxy-beta-l	D-glucopyranose-((1-4)-2-acetamido	-2-deoxy-beta-D	-gluo
opyranose					

Chain E:

100%



• Molecule 3: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D:

30%

70%



• Molecule 4: beta-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:

20%

80%





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 31	Depositor
Cell constants	$92.08 ext{Å} 92.08 ext{Å} 129.50 ext{Å}$	Danasitan
a, b, c, α , β , γ	90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.30 - 2.25	Depositor
rtesoration (A)	79.74 - 2.25	EDS
% Data completeness	98.0 (50.30-2.25)	Depositor
(in resolution range)	98.0 (79.74-2.25)	EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$0.98~({ m at}~2.25{ m \AA})$	Xtriage
Refinement program	PHENIX 1.15.2_3472	Depositor
R, R_{free}	0.181 , 0.221	Depositor
·	0.181 , 0.221	DCC
R_{free} test set	2008 reflections (3.52%)	wwPDB-VP
Wilson B-factor (Å ²)	60.7	Xtriage
Anisotropy	0.154	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.29 , 58.2	EDS
L-test for twinning ²	$< L >=0.49, < L^2>=0.32$	Xtriage
	0.016 for -h,-k,l	
Estimated twinning fraction	0.042 for h,-h-k,-l	Xtriage
	0.032 for -k,-h,-l	
F_o, F_c correlation	0.97	EDS
Total number of atoms	6394	wwPDB-VP
Average B, all atoms (\mathring{A}^2)	87.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.47% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $< L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, NAG, CL, CA, TBU, FE2, OGA, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
IVIOI	Chain	RMSZ	# Z >5	RMSZ	# Z > 5
1	A	0.31	0/3062	0.49	0/4147
1	В	0.31	0/2964	0.50	0/4014
All	All	0.31	0/6026	0.49	0/8161

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2995	0	2910	28	0
1	В	2898	0	2820	36	0
2	С	28	0	25	0	0
2	E	28	0	25	0	0
3	D	116	0	97	0	0
4	F	116	0	97	0	0
5	A	10	0	3	0	0
5	В	10	0	3	0	0
6	A	1	0	0	0	0
6	В	1	0	0	0	0
7	A	20	0	40	2	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	В	20	0	40	1	0
8	A	10	0	4	0	0
9	A	2	0	0	0	0
9	В	2	0	0	0	0
10	A	2	0	0	0	0
11	A	67	0	0	0	0
11	В	68	0	0	0	0
All	All	6394	0	6064	64	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (64) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

A	A	Interatomic	Clash
Atom-1	Atom-2	${\rm distance}({\rm \AA})$	overlap (Å)
1:B:259:LYS:HA	1:B:262:ARG:HB2	1.70	0.72
1:A:340:CYS:HB2	1:A:343:THR:HG22	1.73	0.70
1:B:229:GLU:HG3	7:B:715:TBU:H32	1.75	0.69
1:A:349:GLU:HB3	1:A:351:VAL:HG13	1.76	0.67
1:A:345:LEU:H	1:A:345:LEU:HD23	1.64	0.62
1:A:338:THR:HA	1:A:353:PHE:HB3	1.83	0.59
1:A:115:GLU:O	1:A:123:ARG:NH2	2.31	0.57
1:B:191:LEU:H	1:B:191:LEU:HD12	1.70	0.56
1:B:249:PHE:O	1:B:252:MET:HB2	2.07	0.55
1:A:343:THR:HG21	1:A:461:ASP:OD2	2.07	0.55
1:B:203:GLY:O	1:B:204:HIS:ND1	2.41	0.54
1:B:142:PRO:HG3	1:B:439:SER:HA	1.90	0.53
1:A:467:GLN:O	1:A:471:GLN:HG3	2.09	0.53
1:A:210:VAL:HG12	1:A:222:MET:HE1	1.91	0.53
1:B:220:TRP:CZ3	1:B:266:ALA:HB3	2.44	0.53
1:B:126:GLN:NE2	1:B:132:ASP:OD2	2.39	0.52
1:B:114:LEU:O	1:B:143:LEU:HD21	2.10	0.52
1:A:222:MET:HE2	7:A:716:TBU:H23	1.91	0.51
1:B:287:HIS:HB3	1:B:290:MET:HG2	1.92	0.51
1:B:170:ARG:CZ	1:B:174:LEU:HD11	2.41	0.51
1:A:108:LEU:HD11	1:A:468:ALA:HB2	1.93	0.50
1:A:114:LEU:O	1:A:143:LEU:HD21	2.12	0.50
1:A:142:PRO:HG3	1:A:439:SER:HA	1.94	0.50
1:B:197:LEU:CD2	1:B:213:GLN:HG3	2.42	0.50
1:A:394:ASP:N	1:A:394:ASP:OD1	2.40	0.50
1:B:473:GLU:CG	1:B:476:ARG:HH11	2.25	0.49



Continued from previous page...

Continued from preo		Interatomic	Clash
Atom-1	Atom-2	${ m distance}({ m \AA})$	overlap (Å)
1:B:230:MET:HG3	1:B:260:TYR:CD1	2.49	0.48
1:B:152:THR:OG1	1:B:154:GLU:HG2	2.14	0.47
1:B:473:GLU:HG2	1:B:476:ARG:HH11	1.78	0.47
1:B:253:ASP:OD2	1:B:253:ASP:N	2.45	0.47
1:A:284:GLU:HA	1:A:291:ARG:NH1	2.29	0.47
1:B:476:ARG:HA	1:B:479:ARG:CZ	2.45	0.47
1:A:193:LEU:O	1:A:197:LEU:HG	2.16	0.46
1:B:319:TYR:HB2	1:B:451:LYS:HB3	1.96	0.46
1:B:351:VAL:N	1:B:352:PRO:CD	2.79	0.45
1:B:197:LEU:O	1:B:199:GLN:N	2.50	0.45
1:A:374:GLU:HA	1:A:413:PRO:HD3	1.98	0.44
1:B:311:SER:HB3	1:B:459:ASN:H	1.82	0.44
1:B:142:PRO:HD3	1:B:438:TYR:O	2.17	0.44
1:A:220:TRP:CZ3	1:A:266:ALA:HB3	2.52	0.44
1:B:191:LEU:O	1:B:195:ARG:HG2	2.17	0.44
1:A:359:TYR:CD2	1:A:360:MET:HG3	2.54	0.42
1:A:461:ASP:HB3	1:A:467:GLN:HB2	1.99	0.42
1:B:199:GLN:HG2	1:B:209:GLU:HG2	2.02	0.42
1:B:476:ARG:HA	1:B:479:ARG:NH1	2.34	0.42
1:B:205:LEU:HB2	1:B:244:LEU:HB2	2.02	0.42
1:B:294:ARG:HD3	1:B:309:GLU:HG2	2.01	0.42
1:B:365:TYR:HA	1:B:453:ILE:HG22	2.02	0.42
1:A:349:GLU:CB	1:A:351:VAL:HG13	2.46	0.42
1:B:196:LEU:HD22	1:B:196:LEU:O	2.20	0.42
1:B:197:LEU:HD22	1:B:213:GLN:HG3	2.02	0.42
1:B:359:TYR:CD2	1:B:360:MET:HG3	2.55	0.41
1:A:236:ALA:HB1	1:A:252:MET:HE1	2.02	0.41
1:B:138:LEU:HD11	1:B:146:GLU:HB2	2.03	0.41
1:A:283:GLY:O	1:A:291:ARG:HD3	2.20	0.41
1:A:345:LEU:HG	1:A:346:VAL:HG23	2.02	0.41
1:B:400:THR:HB	1:B:444:CYS:HA	2.02	0.41
1:A:344:LYS:HB2	1:A:344:LYS:HE2	1.65	0.41
1:A:398:ARG:NH1	7:A:717:TBU:O	2.46	0.41
1:A:108:LEU:HD13	1:A:464:ARG:HG2	2.03	0.41
1:A:346:VAL:HB	1:A:349:GLU:HG3	2.02	0.41
1:B:461:ASP:HB3	1:B:467:GLN:HB2	2.03	0.41
1:B:222:MET:HE3	1:B:227:ILE:HD11	2.04	0.40
1:A:279:TRP:CZ3	1:A:315:GLN:HB2	2.56	0.40

There are no symmetry-related clashes. $\,$



5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	${f Analysed}$	Favoured	Allowed	Outliers	Perce	\mathbf{ntiles}
1	A	$366/421 \ (87\%)$	360 (98%)	6 (2%)	0	100	100
1	В	352/421~(84%)	343 (97%)	8 (2%)	1 (0%)	41	46
All	All	718/842 (85%)	703 (98%)	14 (2%)	1 (0%)	51	60

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	В	198	ASP

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	${f Rotameric}$	Outliers	Percentiles		
1	A	325/368~(88%)	321 (99%)	4 (1%)	71 80		
1	В	314/368~(85%)	310 (99%)	4 (1%)	69 79		
All	All	639/736 (87%)	631 (99%)	8 (1%)	69 79		

All (8) residues with a non-rotameric sidechain are listed below:

\mathbf{Mol}	Chain	${f Res}$	\mathbf{Type}
1	A	189	SER
1	A	302	ARG
1	A	325	TYR
1	A	345	LEU
1	В	123	ARG



Continued from previous page...

Mol	Chain	Res	Type
1	В	189	SER
1	В	304	SER
1	В	325	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	169	GLN
1	В	190	GLN
1	В	472	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

24 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Tuna	Chain Res Link		Во	ond leng	ths	Bond angles			
MIOI	Type	Chain	nes	S LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	С	1	1,2	14,14,15	0.27	0	17,19,21	0.39	0
2	NAG	С	2	2	14,14,15	0.23	0	17,19,21	0.66	1 (5%)
3	NAG	D	1	1,3	14,14,15	0.23	0	17,19,21	0.65	1 (5%)
3	MAN	D	10	3	11,11,12	0.80	0	15,15,17	1.02	2 (13%)
3	NAG	D	2	3	14,14,15	0.33	0	17,19,21	0.41	0
3	BMA	D	3	3	11,11,12	0.64	0	15,15,17	0.73	0
3	MAN	D	4	3	11,11,12	0.94	1 (9%)	15,15,17	0.96	2 (13%)



Mol	Type	Chain	Res	Link	Вс	nd leng	ths	В	ond ang	les
WIOI	Type	Chain	res	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	MAN	D	5	3	11,11,12	0.84	0	15,15,17	0.99	0
3	MAN	D	6	3	11,11,12	0.85	0	15,15,17	0.95	1 (6%)
3	MAN	D	7	3	11,11,12	0.88	0	15,15,17	1.26	2 (13%)
3	MAN	D	8	3	11,11,12	0.78	0	15,15,17	1.06	1 (6%)
3	MAN	D	9	3	11,11,12	0.73	0	15,15,17	0.99	2 (13%)
2	NAG	E	1	1,2	14,14,15	0.36	0	17,19,21	0.47	0
2	NAG	Е	2	2	14,14,15	0.23	0	17,19,21	0.46	0
4	NAG	F	1	1,4	14,14,15	0.24	0	17,19,21	0.60	1 (5%)
4	MAN	F	10	4	11,11,12	0.81	0	15,15,17	1.13	2 (13%)
4	NAG	F	2	4	14,14,15	0.28	0	17,19,21	0.40	0
4	BMA	F	3	4	11,11,12	0.62	0	15,15,17	0.80	0
4	MAN	F	4	4	11,11,12	0.96	0	15,15,17	0.97	1 (6%)
4	MAN	F	5	4	11,11,12	0.93	1 (9%)	15,15,17	1.18	2 (13%)
4	BMA	F	6	4	11,11,12	1.00	1 (9%)	15,15,17	0.97	0
4	MAN	F	7	4	11,11,12	0.75	0	15,15,17	1.15	2 (13%)
4	MAN	F	8	4	11,11,12	0.81	0	15,15,17	1.05	1 (6%)
4	MAN	F	9	4	11,11,12	0.90	0	15,15,17	0.99	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	С	1	1,2	-	1/6/23/26	0/1/1/1
2	NAG	С	2	2	-	2/6/23/26	0/1/1/1
3	NAG	D	1	1,3	-	0/6/23/26	0/1/1/1
3	MAN	D	10	3	-	0/2/19/22	0/1/1/1
3	NAG	D	2	3	-	2/6/23/26	0/1/1/1
3	BMA	D	3	3	-	0/2/19/22	0/1/1/1
3	MAN	D	4	3	-	2/2/19/22	0/1/1/1
3	MAN	D	5	3	-	2/2/19/22	0/1/1/1
3	MAN	D	6	3	-	1/2/19/22	0/1/1/1
3	MAN	D	7	3	-	0/2/19/22	0/1/1/1
3	MAN	D	8	3	-	0/2/19/22	0/1/1/1
3	MAN	D	9	3	-	0/2/19/22	0/1/1/1
2	NAG	Ε	1	1,2	-	2/6/23/26	0/1/1/1
2	NAG	Ε	2	2	-	0/6/23/26	0/1/1/1
4	NAG	F	1	1,4	-	0/6/23/26	0/1/1/1



Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	${f Torsions}$	Rings
4	MAN	F	10	4	-	2/2/19/22	0/1/1/1
4	NAG	F	2	4	-	0/6/23/26	0/1/1/1
4	BMA	F	3	4	-	0/2/19/22	0/1/1/1
4	MAN	F	4	4	-	2/2/19/22	0/1/1/1
4	MAN	F	5	4	-	0/2/19/22	0/1/1/1
4	BMA	F	6	4	-	0/2/19/22	0/1/1/1
4	MAN	F	7	4	1	0/2/19/22	0/1/1/1
4	MAN	F	8	4	-	0/2/19/22	0/1/1/1
4	MAN	F	9	4	-	0/2/19/22	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	${f Observed(\AA)}$	$\operatorname{Ideal}(ext{\AA})$
3	D	4	MAN	O5-C1	-2.09	1.40	1.43
4	F	6	BMA	O5-C1	-2.05	1.40	1.43
4	F	5	MAN	O5-C1	-2.01	1.40	1.43

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}({}^o)$	$Ideal(^{o})$
4	F	5	MAN	O2-C2-C3	-2.89	104.35	110.14
4	F	5	MAN	C1-O5-C5	2.73	115.89	112.19
3	D	7	MAN	C1-O5-C5	2.58	115.68	112.19
4	F	4	MAN	C1-O5-C5	2.39	115.42	112.19
4	F	10	MAN	C1-O5-C5	2.37	115.41	112.19
3	D	9	MAN	O2-C2-C3	-2.37	105.39	110.14
3	D	7	MAN	O5-C1-C2	2.36	114.41	110.77
2	С	2	NAG	C1-O5-C5	2.34	115.37	112.19
4	F	7	MAN	C1-O5-C5	2.32	115.34	112.19
3	D	8	MAN	C1-O5-C5	2.29	115.30	112.19
4	F	7	MAN	O2-C2-C3	-2.29	105.56	110.14
3	D	4	MAN	O2-C2-C3	-2.26	105.60	110.14
3	D	10	MAN	C1-O5-C5	2.23	115.21	112.19
4	F	9	MAN	C1-O5-C5	2.22	115.20	112.19
3	D	1	NAG	C1-O5-C5	2.18	115.14	112.19
4	F	9	MAN	O2-C2-C3	-2.15	105.83	110.14
3	D	4	MAN	C1-O5-C5	2.14	115.10	112.19
3	D	10	MAN	O2-C2-C3	-2.14	105.84	110.14
3	D	9	MAN	C1-O5-C5	2.12	115.07	112.19
4	F	10	MAN	O5-C1-C2	2.11	114.03	110.77
3	D	6	MAN	C1-O5-C5	2.08	115.01	112.19
4	F	1	NAG	C1-O5-C5	2.06	114.98	112.19



Continued from previous page...

\mathbf{Mol}	Chain	\mathbf{Res}	Type	${f Atoms}$	\mathbf{Z}	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^o)$
4	F	8	MAN	O2-C2-C3	-2.04	106.05	110.14

There are no chirality outliers.

All (16) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	D	5	MAN	C4-C5-C6-O6
2	С	2	NAG	C4-C5-C6-O6
4	F	4	MAN	O5-C5-C6-O6
2	С	2	NAG	O5-C5-C6-O6
3	D	5	MAN	O5-C5-C6-O6
4	F	4	MAN	C4-C5-C6-O6
3	D	4	MAN	O5-C5-C6-O6
3	D	2	NAG	O5-C5-C6-O6
3	D	2	NAG	C4-C5-C6-O6
3	D	6	MAN	O5-C5-C6-O6
2	С	1	NAG	C4-C5-C6-O6
2	E	1	NAG	C4-C5-C6-O6
2	Е	1	NAG	O5-C5-C6-O6
4	F	10	MAN	O5-C5-C6-O6
4	F	10	MAN	C4-C5-C6-O6
3	D	4	MAN	C4-C5-C6-O6

There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



















5.6 Ligand geometry (i)

Of 20 ligands modelled in this entry, 8 are monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



Mal	Trans	Chain	Res	Link	В	ond leng	gths	Bond angles		
Mol	Type	Chain	nes	Lilik	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2
7	TBU	A	703	_	4,4,4	0.79	0	6,6,6	0.59	0
8	GLY	A	720	_	1,4,4	0.06	0	0,4,4	0.00	-
7	TBU	A	718	_	4,4,4	0.76	0	6,6,6	0.52	0
7	TBU	В	717	-	4,4,4	0.72	0	6,6,6	0.68	0
7	TBU	A	716	_	4,4,4	0.74	0	6,6,6	0.60	0
8	GLY	A	719	_	1,4,4	0.05	0	0,4,4	0.00	-
7	TBU	В	716	_	4,4,4	0.75	0	6,6,6	0.52	0
7	TBU	В	715	_	4,4,4	0.78	0	6,6,6	0.51	0
7	TBU	В	718	_	4,4,4	0.79	0	6,6,6	0.48	0
5	OGA	A	701	6	3,9,9	3.27	2 (66%)	4,11,11	2.54	1 (25%)
7	TBU	A	717	_	4,4,4	0.78	0	6,6,6	0.46	0
5	OGA	В	701	6	3,9,9	3.35	2 (66%)	4,11,11	2.11	1 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	OGA	A	701	6	-	1/3/9/9	-
8	GLY	A	720	_	-	0/0/2/2	_
8	GLY	A	719	_	-	0/0/2/2	-
5	OGA	В	701	6	-	0/3/9/9	_

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	$\operatorname{Ideal}(ext{\AA})$
5	В	701	OGA	C2-N1	5.42	1.45	1.33
5	A	701	OGA	C2-N1	5.23	1.45	1.33
5	A	701	OGA	O2'-C2	-2.16	1.18	1.23
5	В	701	OGA	O2'-C2	-2.04	1.19	1.23

All (2) bond angle outliers are listed below:

	Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^o)$
Ī	5	A	701	OGA	C1-C2-N1	4.53	120.10	115.60
Ī	5	В	701	OGA	C1-C2-N1	4.04	119.62	115.60

There are no chirality outliers.

All (1) torsion outliers are listed below:




Mol	Chain	Res	Type	Atoms
5	A	701	OGA	C5-C4-N1-C2

There are no ring outliers.


3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	A	716	TBU	1	0
7	В	715	TBU	1	0
7	A	717	TBU	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	${f Analysed}$	<RSRZ $>$	$\#\mathrm{RSRZ}{>}2$	$OWAB(m \AA^2)$	Q<0.9
1	A	370/421 (87%)	0.05	12 (3%) 47 50	45, 72, 153, 191	0
1	В	$356/421 \ (84\%)$	0.10	8 (2%) 62 65	49, 87, 152, 199	0
All	All	726/842 (86%)	0.08	20 (2%) 53 55	45, 77, 153, 199	0

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	345	LEU	5.1
1	В	353	PHE	4.1
1	A	179	TYR	3.9
1	A	350	SER	3.8
1	A	188	VAL	3.5
1	A	478	ALA	3.2
1	A	346	VAL	2.9
1	В	481	GLY	2.8
1	A	180	GLU	2.8
1	A	181	GLU	2.8
1	В	176	THR	2.6
1	A	187	GLN	2.6
1	A	477	LEU	2.4
1	В	262	ARG	2.3
1	A	178	GLU	2.3
1	A	347	ALA	2.3
1	В	281[A]	TYR	2.3
1	В	108	LEU	2.3
1	В	174	LEU	2.2
1	В	217	GLY	2.1



6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.


6.3 Carbohydrates (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	${f B\text{-}factors}({f \AA}^2)$	Q<0.9
4	BMA	F	6	11/12	0.62	0.26	174,178,183,183	0
3	MAN	D	5	11/12	0.62	0.23	131,159,169,181	0
3	MAN	D	6	11/12	0.65	0.31	147,179,185,188	0
4	MAN	F	9	11/12	0.69	0.18	114,132,135,136	0
4	MAN	F	8	11/12	0.79	0.14	110,115,127,137	0
3	MAN	D	4	11/12	0.80	0.11	136,146,152,160	0
2	NAG	С	2	14/15	0.84	0.21	144,155,168,170	0
3	MAN	D	9	11/12	0.84	0.28	111,145,154,157	0
2	NAG	Е	2	14/15	0.85	0.19	127,140,147,150	0
4	MAN	F	5	11/12	0.87	0.11	136,146,160,173	0
4	MAN	F	10	11/12	0.87	0.20	105,125,144,146	0
4	MAN	F	4	11/12	0.88	0.08	109,127,132,139	0
3	MAN	D	8	11/12	0.89	0.13	$120,\!125,\!137,\!145$	0
3	BMA	D	3	11/12	0.90	0.08	95,108,125,136	0
2	NAG	Е	1	14/15	0.92	0.13	83,98,124,132	0
4	BMA	F	3	11/12	0.92	0.10	86,113,119,122	0
4	MAN	F	7	11/12	0.92	0.08	102,109,117,122	0
3	MAN	D	10	11/12	0.93	0.18	123,130,143,144	0
3	MAN	D	7	11/12	0.94	0.10	90,96,112,119	0
2	NAG	С	1	14/15	0.94	0.09	84,96,120,144	0
3	NAG	D	2	14/15	0.95	0.10	69,82,88,103	0
4	NAG	F	2	14/15	0.96	0.09	69,80,86,94	0
4	NAG	F	1	14/15	0.98	0.14	52,56,69,72	0
3	NAG	D	1	14/15	0.98	0.14	50,59,65,66	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



















6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	${ m Res}$	Atoms	RSCC	RSR	${f B\text{-factors}}({f A}^2)$	Q<0.9
8	GLY	A	719	5/5	0.76	0.11	95,100,117,119	0
7	TBU	A	703	5/5	0.84	0.22	86,86,94,95	0
10	CL	A	723	1/1	0.85	0.32	102,102,102,102	0




Continued from previous page...

Mol	Type	Chain	${f Res}$	Atoms	RSCC	RSR	${f B\text{-factors}}({f A}^2)$	$\mathbf{Q}{<}0.9$
7	TBU	В	718	5/5	0.86	0.20	85,97,100,101	0
7	TBU	В	717	5/5	0.87	0.26	98,101,104,106	0
9	CA	В	720	1/1	0.87	0.11	103,103,103,103	0
7	TBU	A	718	5/5	0.88	0.24	105,106,111,116	0
7	TBU	В	715	5/5	0.89	0.14	94,102,103,105	0
8	GLY	A	720	5/5	0.89	0.19	128,137,159,163	0
5	OGA	A	701	10/10	0.91	0.18	48,82,120,123	0
7	TBU	В	716	5/5	0.92	0.41	80,84,95,102	0
5	OGA	В	701	10/10	0.92	0.16	54,85,109,111	0
7	TBU	A	717	5/5	0.93	0.19	84,91,103,112	0
10	CL	A	724	1/1	0.96	0.07	106,106,106,106	0
7	TBU	A	716	5/5	0.96	0.31	81,85,87,92	0
9	CA	В	719	1/1	0.98	0.09	100,100,100,100	0
9	CA	A	722	1/1	0.99	0.15	77,77,77,77	0
6	FE2	В	702	1/1	0.99	0.14	85,85,85,85	0
9	CA	A	721	1/1	0.99	0.12	69,69,69,69	0
6	FE2	A	702	1/1	0.99	0.13	80,80,80,80	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.










Electron density around OGA A 701: 2mF_o-DF_c (at 0.7 rmsd) in gray mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)

Electron density around OGA B 701:


 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)



























6.5 Other polymers (i)

There are no such residues in this entry.

