



Full wwPDB NMR Structure Validation Report ⓘ

Feb 20, 2022 – 01:21 AM EST

PDB ID : 1TQR
Title : NMR Structure of DNA 17-mer GGAAAATCTCTAGCAGT corresponding to the extremity of the U5 LTR of the HIV-1 genome
Authors : Renisio, J.G.; Cosquer, S.; Cherrak, I.; El Antri, S.; Mauffret, O.; Fermandjian, S.
Deposited on : 2004-06-18

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : 2.26
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.26


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR



The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	17	 12% 88%
2	B	17	 53% 47%

2 Ensemble composition and analysis

This entry contains 9 models. This entry does not contain polypeptide chains, therefore identification of well-defined residues and clustering analysis are not possible. All residues are included in the validation scores.

3 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 1079 atoms, of which 388 are hydrogens and 0 are deuteriums.

- Molecule 1 is a DNA chain called DNA HIV-1 U5 LTR extremity.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	P	
1	A	17	541	167	193	67	98	16	0

- Molecule 2 is a DNA chain called DNA HIV-1 U5 LTR extremity.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	P	
2	B	17	538	166	195	59	102	16	0

4 Residue-property plots

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: DNA HIV-1 U5 LTR extremity



- Molecule 2: DNA HIV-1 U5 LTR extremity



4.2 Scores per residue for each member of the ensemble


Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1

- Molecule 1: DNA HIV-1 U5 LTR extremity



- Molecule 2: DNA HIV-1 U5 LTR extremity




4.2.2 Score per residue for model 2

- Molecule 1: DNA HIV-1 U5 LTR extremity




- Molecule 2: DNA HIV-1 U5 LTR extremity




4.2.3 Score per residue for model 3

- Molecule 1: DNA HIV-1 U5 LTR extremity



- Molecule 2: DNA HIV-1 U5 LTR extremity



4.2.4 Score per residue for model 4

- Molecule 1: DNA HIV-1 U5 LTR extremity




- Molecule 2: DNA HIV-1 U5 LTR extremity



4.2.5 Score per residue for model 5

- Molecule 1: DNA HIV-1 U5 LTR extremity



- Molecule 2: DNA HIV-1 U5 LTR extremity



4.2.6 Score per residue for model 6

- Molecule 1: DNA HIV-1 U5 LTR extremity



- Molecule 2: DNA HIV-1 U5 LTR extremity



4.2.7 Score per residue for model 7

- Molecule 1: DNA HIV-1 U5 LTR extremity




- Molecule 2: DNA HIV-1 U5 LTR extremity



4.2.8 Score per residue for model 8

- Molecule 1: DNA HIV-1 U5 LTR extremity



- Molecule 2: DNA HIV-1 U5 LTR extremity



4.2.9 Score per residue for model 9

- Molecule 1: DNA HIV-1 U5 LTR extremity



- Molecule 2: DNA HIV-1 U5 LTR extremity



5 Refinement protocol and experimental data overview

The models were refined using the following method: *torsion angle dynamics including classical and RDC constraints*.

Of the 50 calculated structures, 9 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CNS	refinement	0.9a

No chemical shift data was provided.

6 Model quality i

6.1 Standard geometry i

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	348	193	193	10±2
2	B	343	195	195	10±2
All	All	6219	3492	3492	174

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:5:DA:C2	1:A:6:DA:C4	0.55	2.95	2	6
1:A:7:DT:C4	1:A:8:DC:C4	0.53	2.96	9	8
1:A:16:DG:H2 [?]	1:A:17:DT:C6	0.53	2.39	8	9
1:A:3:DA:C2	1:A:4:DA:C2	0.53	2.96	7	4
2:B:27:DG:H2 [?]	2:B:28:DA:C8	0.52	2.39	7	7
1:A:16:DG:C2 [?]	1:A:17:DT:H71	0.52	2.34	5	6
2:B:28:DA:C2	2:B:29:DT:C2	0.52	2.98	2	8
2:B:26:DA:C6	2:B:27:DG:C6	0.51	2.98	2	6
2:B:32:DT:C4	2:B:33:DC:N4	0.51	2.79	7	5
2:B:24:DA:C2	2:B:25:DG:C4	0.50	2.99	4	7
2:B:29:DT:C6	2:B:30:DT:H72	0.50	2.41	2	1
1:A:3:DA:C2	1:A:4:DA:C4	0.49	3.01	6	1
1:A:13:DG:H2 [?]	1:A:14:DC:C6	0.49	2.43	2	5

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
2:B:27:DG:N2	2:B:28:DA:C2	0.48	2.82	5	6
1:A:4:DA:C5	1:A:5:DA:C5	0.48	3.02	7	3
2:B:25:DG:C6	2:B:26:DA:N6	0.47	2.82	1	5
1:A:16:DG:C4	1:A:17:DT:C4	0.47	3.02	8	1
1:A:9:DT:C4	1:A:10:DC:N4	0.47	2.83	2	7
1:A:2:DG:N2	2:B:34:DC:C2	0.46	2.83	3	3
2:B:25:DG:C2	2:B:26:DA:C5	0.46	3.03	2	5
2:B:23:DT:C2	2:B:24:DA:C8	0.45	3.04	4	3
1:A:15:DA:C4	2:B:21:DG:N2	0.45	2.85	1	2
2:B:27:DG:C2	2:B:28:DA:C2	0.45	3.04	7	7
1:A:8:DC:C6	1:A:9:DT:H72	0.45	2.47	7	2
2:B:30:DT:C2'	2:B:31:DT:H72	0.44	2.42	7	2
2:B:23:DT:C2	2:B:24:DA:N7	0.44	2.85	2	3
2:B:25:DG:C2	2:B:26:DA:C6	0.44	3.05	1	1
2:B:30:DT:H2''	2:B:31:DT:H72	0.44	1.88	1	2
1:A:13:DG:C2	1:A:14:DC:C2	0.44	3.06	1	1
1:A:14:DC:C2	1:A:15:DA:C8	0.44	3.05	2	6
1:A:10:DC:C6	1:A:11:DT:C7	0.43	3.01	4	9
1:A:9:DT:C4	1:A:10:DC:C4	0.43	3.05	3	6
2:B:25:DG:N1	2:B:26:DA:C6	0.42	2.87	4	4
1:A:1:DG:C6	1:A:2:DG:C6	0.42	3.07	7	1
1:A:16:DG:H2''	1:A:17:DT:H71	0.42	1.91	5	3
1:A:1:DG:H2''	1:A:2:DG:C8	0.42	2.50	4	1
1:A:4:DA:C6	1:A:5:DA:C6	0.42	3.08	7	3
1:A:10:DC:C2	1:A:11:DT:C5	0.42	3.08	6	2
1:A:9:DT:C2	2:B:27:DG:N2	0.41	2.88	6	1
1:A:8:DC:H2''	1:A:9:DT:OP2	0.41	2.15	9	1
1:A:8:DC:C2	1:A:9:DT:C5	0.41	3.08	5	1
2:B:28:DA:C2'	2:B:29:DT:H72	0.41	2.46	9	1
2:B:29:DT:C2	2:B:30:DT:C5	0.41	3.09	5	1
2:B:26:DA:C2	2:B:27:DG:C2	0.41	3.09	8	1
2:B:18:DA:H2''	2:B:19:DC:C6	0.41	2.50	7	2
1:A:16:DG:C2'	1:A:17:DT:H72	0.41	2.45	2	1
2:B:30:DT:C2'	2:B:31:DT:H71	0.41	2.46	2	1
2:B:21:DG:H2''	2:B:22:DC:C6	0.40	2.50	4	1
1:A:4:DA:N6	1:A:5:DA:C6	0.40	2.90	5	1
2:B:30:DT:H2''	2:B:31:DT:C7	0.40	2.47	7	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

6.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided