



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 24, 2024 – 10:25 AM EDT

PDB ID : 6TQ5
Title : Alcohol dehydrogenase from *Candida magnoliae* DSMZ 70638 (ADHA): complex with NADP+
Authors : Rovida, S.; Aalbers, F.S.; Fraaije, M.W.; Mattevi, A.
Deposited on : 2019-12-16
Resolution : 1.60 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 2.37.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION





The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3398 (1.60-1.60)
Clashscore	141614	3665 (1.60-1.60)
Ramachandran outliers	138981	3564 (1.60-1.60)
Sidechain outliers	138945	3563 (1.60-1.60)
RSRZ outliers	127900	3321 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	246	
1	B	246	
1	C	246	
1	D	246	

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 7848 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Enzyme subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	240	Total 1797	C 1133	N 307	O 349	S 8	0	5	0
1	B	240	Total 1781	C 1125	N 303	O 345	S 8	0	4	0
1	C	240	Total 1773	C 1118	N 301	O 346	S 8	0	3	0
1	D	240	Total 1761	C 1110	N 301	O 342	S 8	0	0	0

- Molecule 2 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NAP) (formula: $C_{21}H_{28}N_7O_{17}P_3$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	Total 40	C 15	N 6	O 16	P 3	0	0
2	B	1	Total 39	C 15	N 5	O 16	P 3	0	0

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	C	1	Total	C	N	O	P	0	0
			39	15	5	16	3		
2	D	1	Total	C	N	O	P	0	0
			39	15	5	16	3		

- Molecule 3 is ISOPROPYL ALCOHOL (three-letter code: IPA) (formula: C₃H₈O).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	3	1		
3	C	1	Total	C	O	0	0
			4	3	1		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	C	1	Total C O 6 3 3	0	0
4	D	1	Total C O 6 3 3	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	137	Total O 137 137	0	0
5	B	143	Total O 143 143	0	0
5	C	140	Total O 140 140	0	0
5	D	139	Total O 139 139	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Enzyme subunit




- Molecule 1: Enzyme subunit



- Molecule 1: Enzyme subunit



- Molecule 1: Enzyme subunit



4 Data and refinement statistics

Property	Value	Source
Space group	I 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	76.11Å 113.14Å 124.21Å 90.00° 94.04° 90.00°	Depositor
Resolution (Å)	83.55 – 1.60 45.54 – 1.60	Depositor EDS
% Data completeness (in resolution range)	97.3 (83.55-1.60) 97.3 (45.54-1.60)	Depositor EDS
R_{merge}	0.17	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.06 (at 1.60Å)	Xtrriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.141 , 0.197 0.146 , 0.195	Depositor DCC
R_{free} test set	6542 reflections (4.88%)	wwPDB-VP
Wilson B-factor (Å ²)	14.8	Xtrriage
Anisotropy	0.453	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 54.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	7848	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.33% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: NAP, GOL, IPA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.40	3/1830 (0.2%)	1.24	9/2487 (0.4%)
1	B	1.27	6/1823 (0.3%)	1.23	10/2477 (0.4%)
1	C	1.26	5/1815 (0.3%)	1.24	15/2468 (0.6%)
1	D	1.29	6/1794 (0.3%)	1.22	17/2439 (0.7%)
All	All	1.31	20/7262 (0.3%)	1.24	51/9871 (0.5%)

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	69	SER	CB-OG	-16.73	1.20	1.42
1	D	97	SER	CB-OG	-9.51	1.29	1.42
1	D	246	ALA	C-O	8.91	1.40	1.23
1	B	72	GLU	CD-OE1	8.69	1.35	1.25
1	A	157	MET	CB-CG	7.76	1.76	1.51
1	B	72	GLU	CD-OE2	7.37	1.33	1.25
1	B	88	ASP	CG-OD2	7.17	1.41	1.25
1	C	133	SER	CB-OG	-7.10	1.33	1.42
1	D	133	SER	CB-OG	-6.96	1.33	1.42
1	B	72	GLU	CG-CD	6.88	1.62	1.51
1	D	69	SER	CB-OG	-6.63	1.33	1.42
1	A	97	SER	CB-OG	-6.61	1.33	1.42
1	C	39	SER	CB-OG	-6.34	1.34	1.42
1	C	97	SER	CB-OG	-6.13	1.34	1.42
1	D	17	SER	CB-OG	5.30	1.49	1.42
1	C	7	PRO	N-CA	-5.27	1.38	1.47
1	D	88	ASP	CG-OD1	5.18	1.37	1.25
1	B	46	GLU	CG-CD	-5.11	1.44	1.51
1	B	226	TYR	CE2-CZ	-5.01	1.32	1.38
1	C	218	ASP	CG-OD1	5.00	1.36	1.25

All (51) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	88	ASP	CB-CG-OD1	-13.55	106.10	118.30
1	B	88	ASP	CB-CG-OD2	12.81	129.83	118.30
1	D	157	MET	CG-SD-CE	-9.66	84.74	100.20
1	A	153	ARG	NE-CZ-NH2	-9.61	115.50	120.30
1	B	18	ARG	NE-CZ-NH1	-9.59	115.51	120.30
1	A	157	MET	CG-SD-CE	9.09	114.73	100.20
1	D	246	ALA	CA-C-O	-8.96	101.28	120.10
1	D	88	ASP	CB-CG-OD2	-8.78	110.40	118.30
1	C	169	ASP	CB-CG-OD1	8.34	125.81	118.30
1	C	218	ASP	CB-CG-OD2	-8.22	110.91	118.30
1	A	88	ASP	CB-CG-OD1	-8.18	110.94	118.30
1	B	18	ARG	NE-CZ-NH2	8.07	124.33	120.30
1	B	177	ARG	NE-CZ-NH2	-8.01	116.29	120.30
1	A	169	ASP	CB-CG-OD1	7.92	125.43	118.30
1	C	88[A]	ASP	CB-CG-OD2	-7.92	111.17	118.30
1	C	88[B]	ASP	CB-CG-OD2	-7.92	111.17	118.30
1	C	137	ARG	NE-CZ-NH1	-7.89	116.35	120.30
1	C	153	ARG	NE-CZ-NH2	-7.54	116.53	120.30
1	D	222	ASP	CB-CG-OD2	-7.44	111.61	118.30
1	C	171	PHE	CB-CG-CD2	-7.19	115.77	120.80
1	D	177	ARG	NE-CZ-NH2	-7.14	116.73	120.30
1	C	173	ARG	NE-CZ-NH1	6.98	123.79	120.30
1	D	208	ASP	CB-CG-OD2	-6.96	112.04	118.30
1	A	195	ARG	NE-CZ-NH1	6.92	123.76	120.30
1	C	60	THR	CA-CB-CG2	-6.75	102.95	112.40
1	D	195	ARG	NE-CZ-NH2	-6.56	117.02	120.30
1	C	67	ASP	CB-CG-OD1	6.29	123.96	118.30
1	D	195	ARG	NE-CZ-NH1	6.08	123.34	120.30
1	A	107	ASP	CB-CG-OD1	6.03	123.72	118.30
1	D	67	ASP	CB-CG-OD1	5.87	123.58	118.30
1	B	195	ARG	NE-CZ-NH2	-5.75	117.42	120.30
1	B	195	ARG	NE-CZ-NH1	5.59	123.09	120.30
1	C	137	ARG	NE-CZ-NH2	5.59	123.09	120.30
1	B	56	LYS	CA-CB-CG	5.54	125.59	113.40
1	C	40	ARG	NE-CZ-NH2	-5.54	117.53	120.30
1	A	70	ASP	CB-CG-OD2	-5.49	113.36	118.30
1	D	137	ARG	NE-CZ-NH2	-5.45	117.58	120.30
1	B	56	LYS	CB-CA-C	-5.43	99.53	110.40
1	D	137	ARG	NE-CZ-NH1	5.43	123.01	120.30
1	D	107	ASP	CB-CG-OD1	5.42	123.18	118.30
1	A	130	LYS	CD-CE-NZ	5.41	124.13	111.70
1	A	157	MET	CB-CG-SD	-5.36	96.31	112.40
1	B	157	MET	CG-SD-CE	-5.36	91.63	100.20

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	134	ASP	CB-CG-OD1	5.26	123.03	118.30
1	D	18	ARG	NE-CZ-NH1	-5.26	117.67	120.30
1	D	153	ARG	NE-CZ-NH1	5.22	122.91	120.30
1	C	218	ASP	CB-CG-OD1	5.18	122.96	118.30
1	C	177	ARG	NE-CZ-NH2	-5.17	117.72	120.30
1	C	133	SER	CA-CB-OG	-5.09	97.45	111.20
1	D	229	LYS	CD-CE-NZ	-5.07	100.03	111.70
1	D	169	ASP	CB-CG-OD2	-5.05	113.75	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1797	0	1813	12	0
1	B	1781	0	1808	9	0
1	C	1773	0	1791	19	0
1	D	1761	0	1775	11	0
2	A	40	0	19	0	0
2	B	39	0	18	0	0
2	C	39	0	18	0	0
2	D	39	0	18	0	0
3	A	4	0	8	0	0
3	C	4	0	8	3	0
4	C	6	0	7	3	0
4	D	6	0	8	2	0
5	A	137	0	0	1	1
5	B	143	0	0	3	0
5	C	140	0	0	2	0
5	D	139	0	0	3	0
All	All	7848	0	7291	49	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:157:MET:CG	1:A:157:MET:CB	1.76	1.63
1:A:153:ARG:HD2	5:D:409:HOH:O	1.56	1.05
5:B:521:HOH:O	1:C:153:ARG:HD2	1.64	0.95
1:A:35:VAL:H	1:A:61:HIS:HD2	1.19	0.91
1:D:35:VAL:H	1:D:61:HIS:HD2	1.23	0.86
1:B:35:VAL:H	1:B:61:HIS:HD2	1.25	0.84
1:A:157:MET:CB	1:A:157:MET:SD	2.66	0.83
1:C:35:VAL:H	1:C:61:HIS:HD2	1.24	0.80
1:A:135:LYS:HE2	5:A:438:HOH:O	1.90	0.71
1:C:35:VAL:H	1:C:61:HIS:CD2	2.09	0.68
1:D:35:VAL:H	1:D:61:HIS:CD2	2.11	0.61
1:D:96:ILE:HD11	5:D:462:HOH:O	2.01	0.59
1:D:56:LYS:HE3	4:D:302:GOL:O2	2.04	0.58
1:C:62:HIS:HD1	4:C:303:GOL:C3	2.17	0.57
1:A:35:VAL:H	1:A:61:HIS:CD2	2.11	0.56
1:B:35:VAL:H	1:B:61:HIS:CD2	2.15	0.55
1:C:31:GLU:O	3:C:302:IPA:H32	2.09	0.53
1:C:49:LYS:HD3	1:C:63:VAL:HG13	1.91	0.53
1:C:192:GLY:O	1:C:194[A]:THR:HG23	2.08	0.53
1:C:31:GLU:O	3:C:302:IPA:C3	2.58	0.52
1:C:53:PRO:O	1:C:61:HIS:HE1	1.92	0.51
1:C:80:ALA:HB1	5:C:410:HOH:O	2.10	0.51
1:A:157:MET:CG	1:A:157:MET:CA	2.81	0.49
1:B:153:ARG:HG3	1:C:153:ARG:NH2	2.29	0.47
1:C:154:GLY:HA2	5:C:423:HOH:O	2.15	0.47
1:C:203:ASP:OD1	1:C:205:SER:OG	2.17	0.47
1:B:212:ASN:ND2	5:B:401:HOH:O	2.48	0.46
1:C:194[B]:THR:HG23	1:C:196:THR:HG23	1.97	0.46
1:B:96[B]:ILE:HD13	1:B:96[B]:ILE:HA	1.56	0.46
1:B:153:ARG:HG3	1:C:153:ARG:CZ	2.45	0.46
1:C:62:HIS:HD1	4:C:303:GOL:H31	1.82	0.45
1:A:157:MET:HB3	1:A:157:MET:CE	2.47	0.45
1:A:140:PRO:HD2	1:A:231[B]:LYS:HG2	1.99	0.44
1:C:56:LYS:HD2	3:C:302:IPA:H33	1.98	0.44
1:A:53:PRO:O	1:A:61:HIS:HE1	2.00	0.44
1:C:81:PRO:HB2	4:C:303:GOL:H31	1.99	0.43
1:D:171:PHE:CE2	1:D:175:LEU:HD11	2.53	0.43
1:D:192:GLY:O	1:D:194:THR:HG23	2.18	0.43
1:B:153:ARG:HD3	1:B:244:TYR:CZ	2.54	0.43
1:C:20:ILE:HD11	1:C:194[B]:THR:HG21	2.00	0.42
1:A:57:GLN:HG2	1:A:58:GLY:N	2.35	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:53:PRO:O	1:D:61:HIS:HE1	2.03	0.42
1:B:96[B]:ILE:HG23	1:B:96[B]:ILE:HD12	1.70	0.42
1:D:139:THR:HG23	5:D:432:HOH:O	2.20	0.41
1:B:96[B]:ILE:HD13	5:B:434:HOH:O	2.20	0.41
1:D:180:GLY:N	1:D:181:PRO:CD	2.84	0.41
1:A:143:ILE:O	1:A:186:VAL:HA	2.21	0.40
1:D:156:PRO:C	1:D:157:MET:HG2	2.42	0.40
1:D:8:SER:HA	4:D:302:GOL:O3	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:536:HOH:O	5:A:536:HOH:O[2_555]	0.68	1.52

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	243/246 (99%)	235 (97%)	8 (3%)	0	100	100
1	B	242/246 (98%)	237 (98%)	5 (2%)	0	100	100
1	C	241/246 (98%)	234 (97%)	7 (3%)	0	100	100
1	D	238/246 (97%)	231 (97%)	7 (3%)	0	100	100
All	All	964/984 (98%)	937 (97%)	27 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	194/195 (100%)	192 (99%)	2 (1%)	76	61
1	B	193/195 (99%)	190 (98%)	3 (2%)	62	41
1	C	192/195 (98%)	190 (99%)	2 (1%)	76	61
1	D	189/195 (97%)	187 (99%)	2 (1%)	73	57
All	All	768/780 (98%)	759 (99%)	9 (1%)	71	54

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	56	LYS
1	A	57	GLN
1	B	57	GLN
1	B	157	MET
1	B	242	ASN
1	C	7	PRO
1	C	203	ASP
1	D	200	GLU
1	D	242	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (21) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	61	HIS
1	A	142	HIS
1	A	232	ASN
1	A	242	ASN
1	B	61	HIS
1	B	142	HIS
1	B	212	ASN
1	B	232	ASN
1	B	242	ASN
1	C	57	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	61	HIS
1	C	142	HIS
1	C	212	ASN
1	C	232	ASN
1	C	242	ASN
1	D	57	GLN
1	D	61	HIS
1	D	65	GLN
1	D	212	ASN
1	D	232	ASN
1	D	242	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAP	A	301	-	38,43,52	1.34	3 (7%)	52,67,80	2.55	12 (23%)
3	IPA	A	302	-	3,3,3	0.98	0	3,3,3	1.18	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	GOL	D	302	-	5,5,5	0.50	0	5,5,5	1.87	2 (40%)
2	NAP	C	301	-	38,42,52	2.03	7 (18%)	50,65,80	1.80	8 (16%)
2	NAP	B	301	-	38,42,52	1.83	7 (18%)	50,65,80	2.15	16 (32%)
3	IPA	C	302	-	3,3,3	0.84	0	3,3,3	1.91	1 (33%)
4	GOL	C	303	-	5,5,5	1.95	2 (40%)	5,5,5	2.85	2 (40%)
2	NAP	D	301	-	38,42,52	1.48	6 (15%)	50,65,80	2.09	16 (32%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAP	A	301	-	-	3/23/59/67	0/4/4/5
4	GOL	D	302	-	-	2/4/4/4	-
2	NAP	C	301	-	-	4/23/56/67	0/4/4/5
2	NAP	B	301	-	-	3/23/56/67	0/4/4/5
4	GOL	C	303	-	-	1/4/4/4	-
2	NAP	D	301	-	-	4/23/56/67	0/4/4/5

All (25) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	301	NAP	PN-O3	7.88	1.68	1.59
2	C	301	NAP	PA-O3	5.89	1.65	1.59
2	A	301	NAP	PA-O3	4.99	1.64	1.59
2	B	301	NAP	C2A-N3A	4.95	1.39	1.32
2	B	301	NAP	P2B-O2B	4.83	1.68	1.59
2	D	301	NAP	P2B-O2B	4.76	1.67	1.59
2	B	301	NAP	PN-O3	4.03	1.63	1.59
2	D	301	NAP	O4D-C1D	3.97	1.52	1.43
2	C	301	NAP	P2B-O2B	3.95	1.66	1.59
2	A	301	NAP	C2A-N3A	3.66	1.37	1.32
2	B	301	NAP	O4D-C1D	3.28	1.50	1.43
2	D	301	NAP	PA-O3	3.26	1.63	1.59
2	B	301	NAP	PA-O3	3.12	1.62	1.59
4	C	303	GOL	O1-C1	2.95	1.54	1.42
2	B	301	NAP	O4D-C4D	2.82	1.49	1.44
2	D	301	NAP	C2A-N3A	2.72	1.36	1.32
2	C	301	NAP	C4A-N3A	2.64	1.39	1.35

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	301	NAP	C4A-N3A	-2.53	1.32	1.35
2	C	301	NAP	O4B-C1B	2.52	1.44	1.40
2	A	301	NAP	P2B-O2B	2.28	1.63	1.59
2	D	301	NAP	C6A-C5A	2.28	1.51	1.43
2	C	301	NAP	PN-O1N	2.23	1.58	1.50
4	C	303	GOL	O3-C3	-2.22	1.33	1.42
2	D	301	NAP	C1D-C2D	2.16	1.55	1.51
2	C	301	NAP	O4D-C1D	2.11	1.48	1.43

All (57) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	301	NAP	O4D-C1D-N1N	-10.95	93.13	110.57
2	D	301	NAP	O4D-C1D-C2D	-9.84	88.81	105.76
2	A	301	NAP	C4B-O4B-C1B	-8.25	102.37	109.92
2	B	301	NAP	O4D-C1D-C2D	-7.45	92.92	105.76
2	B	301	NAP	N3A-C2A-N1A	-6.74	119.52	128.67
2	C	301	NAP	C4B-O4B-C1B	-6.68	103.81	109.92
4	C	303	GOL	O2-C2-C3	-5.10	88.08	109.18
2	A	301	NAP	O4B-C4B-C3B	5.07	115.22	105.15
2	C	301	NAP	O4D-C1D-C2D	-4.94	97.25	105.76
2	C	301	NAP	O5D-C5D-C4D	-4.00	95.38	108.99
2	B	301	NAP	O3-PA-O1A	-3.72	99.52	110.70
2	A	301	NAP	O2D-C2D-C1D	3.60	122.47	110.10
2	A	301	NAP	O2A-PA-O1A	3.31	127.85	112.44
2	B	301	NAP	O2N-PN-O3	-3.22	98.58	107.27
4	D	302	GOL	O1-C1-C2	-3.11	96.39	110.38
2	C	301	NAP	O3-PN-O1N	-3.04	101.57	110.70
2	B	301	NAP	O2B-P2B-O1X	-3.03	98.53	109.33
2	A	301	NAP	C4A-C5A-N7A	-3.02	106.15	109.34
2	D	301	NAP	O3-PN-O1N	-2.93	101.89	110.70
2	A	301	NAP	O4D-C4D-C3D	-2.92	99.36	105.15
2	A	301	NAP	O2A-PA-O3	-2.82	99.64	107.27
2	D	301	NAP	O2A-PA-O3	-2.80	99.70	107.27
4	C	303	GOL	O2-C2-C1	2.76	120.61	109.18
2	B	301	NAP	C6A-C5A-C4A	-2.63	112.78	117.90
2	D	301	NAP	O4D-C4D-C3D	-2.60	102.23	104.63
2	C	301	NAP	N3A-C2A-N1A	-2.59	125.15	128.67
2	B	301	NAP	O2N-PN-O1N	2.56	124.34	112.44
2	C	301	NAP	O4B-C4B-C3B	2.55	110.21	105.15
3	C	302	IPA	O2-C2-C1	2.51	126.44	110.28
2	D	301	NAP	N3A-C2A-N1A	-2.50	125.28	128.67

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	301	NAP	O2A-PA-O3	2.48	113.97	107.27
2	C	301	NAP	O2N-PN-O1N	2.47	123.95	112.44
2	D	301	NAP	O3-PA-O1A	2.45	118.08	110.70
2	A	301	NAP	O5B-PA-O1A	-2.45	99.23	108.94
2	D	301	NAP	O2N-PN-O5D	2.42	118.54	107.57
2	D	301	NAP	O2B-P2B-O1X	-2.38	100.86	109.33
2	D	301	NAP	O3X-P2B-O2B	-2.28	96.95	105.85
2	C	301	NAP	C4A-C5A-N7A	-2.28	106.93	109.34
2	D	301	NAP	C3B-C2B-C1B	-2.28	98.45	102.81
2	D	301	NAP	O2N-PN-O3	2.24	113.33	107.27
2	A	301	NAP	C4D-O4D-C1D	2.22	113.24	108.63
2	B	301	NAP	O4D-C4D-C3D	-2.22	102.58	104.63
2	A	301	NAP	N3A-C2A-N1A	-2.22	125.67	128.67
2	D	301	NAP	O2D-C2D-C3D	2.22	115.79	111.43
2	B	301	NAP	N6A-C6A-N1A	2.20	123.03	118.33
2	B	301	NAP	O5D-C5D-C4D	-2.17	101.59	108.99
2	D	301	NAP	O2A-PA-O1A	2.17	122.54	112.44
2	B	301	NAP	C1B-N9A-C4A	-2.16	122.85	126.64
2	B	301	NAP	C4A-C5A-N7A	2.14	111.59	109.34
2	D	301	NAP	O3X-P2B-O2X	2.12	115.75	107.80
2	D	301	NAP	C1B-N9A-C4A	-2.09	122.97	126.64
2	D	301	NAP	C4B-O4B-C1B	-2.09	108.01	109.92
4	D	302	GOL	O2-C2-C1	-2.07	100.59	109.18
2	B	301	NAP	C2B-C1B-N9A	-2.07	107.97	112.56
2	A	301	NAP	O2B-C2B-C1B	2.05	117.27	110.05
2	B	301	NAP	O2X-P2B-O2B	-2.02	97.98	105.85
2	B	301	NAP	O2X-P2B-O1X	2.01	118.67	110.83

There are no chirality outliers.

All (17) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	301	NAP	C5D-O5D-PN-O3
2	A	301	NAP	C5D-O5D-PN-O2N
2	B	301	NAP	C5D-O5D-PN-O3
2	B	301	NAP	C5D-O5D-PN-O1N
2	C	301	NAP	C5D-O5D-PN-O3
2	C	301	NAP	C5D-O5D-PN-O1N
2	C	301	NAP	C5D-O5D-PN-O2N
2	D	301	NAP	C5D-O5D-PN-O3
2	D	301	NAP	C5D-O5D-PN-O2N
4	D	302	GOL	O1-C1-C2-C3

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
4	C	303	GOL	O1-C1-C2-O2
4	D	302	GOL	O1-C1-C2-O2
2	A	301	NAP	C5D-O5D-PN-O1N
2	B	301	NAP	C5D-O5D-PN-O2N
2	D	301	NAP	C5D-O5D-PN-O1N
2	C	301	NAP	C2B-O2B-P2B-O3X
2	D	301	NAP	C2B-O2B-P2B-O2X


There are no ring outliers.


3 monomers are involved in 8 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	302	GOL	2	0
3	C	302	IPA	3	0
4	C	303	GOL	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	240/246 (97%)	0.36	10 (4%) 36 33	9, 13, 32, 53	0
1	B	240/246 (97%)	0.38	13 (5%) 25 23	10, 16, 35, 52	0
1	C	240/246 (97%)	0.50	17 (7%) 16 14	9, 15, 42, 83	0
1	D	240/246 (97%)	0.41	11 (4%) 32 29	10, 16, 38, 54	0
All	All	960/984 (97%)	0.41	51 (5%) 26 24	9, 15, 36, 83	0

All (51) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	201	GLY	7.8
1	C	197	SER	7.8
1	C	202	ILE	6.8
1	D	7	PRO	5.3
1	D	206	MET	5.1
1	C	199	ALA	5.1
1	A	206	MET	5.0
1	C	200	GLU	4.9
1	B	206	MET	4.6
1	B	208	ASP	4.6
1	C	206	MET	4.6
1	C	205	SER	4.5
1	C	207	PHE	4.4
1	B	202	ILE	4.4
1	A	208	ASP	4.3
1	D	138	GLN	4.1
1	C	198	MET	4.1
1	A	7	PRO	4.0
1	B	7	PRO	4.0
1	A	207	PHE	3.9
1	D	134	ASP	3.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	207	PHE	3.7
1	D	208	ASP	3.6
1	A	202	ILE	3.5
1	C	203	ASP	3.5
1	C	204	PRO	3.4
1	D	207	PHE	3.3
1	A	199	ALA	3.2
1	D	202	ILE	3.1
1	B	203	ASP	3.1
1	B	205	SER	3.1
1	D	205	SER	3.1
1	B	134	ASP	3.0
1	D	99	PHE	2.9
1	C	43	GLU	2.9
1	B	57	GLN	2.9
1	C	208	ASP	2.8
1	B	201	GLY	2.6
1	B	136	PRO	2.5
1	A	201	GLY	2.4
1	A	58	GLY	2.4
1	D	201	GLY	2.3
1	B	199	ALA	2.3
1	B	212	ASN	2.3
1	C	246	ALA	2.3
1	C	212	ASN	2.1
1	A	205	SER	2.1
1	C	194[A]	THR	2.1
1	D	203	ASP	2.1
1	C	7	PRO	2.0
1	A	57	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)


There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	IPA	C	302	4/4	0.73	0.13	27,29,29,36	0
4	GOL	D	302	6/6	0.81	0.18	27,37,46,51	0
2	NAP	C	301	39/48	0.87	0.16	17,24,41,44	0
4	GOL	C	303	6/6	0.90	0.18	26,27,34,48	0
2	NAP	B	301	39/48	0.94	0.11	16,18,23,25	0
2	NAP	A	301	40/48	0.95	0.09	10,15,20,34	0
2	NAP	D	301	39/48	0.95	0.09	14,19,25,27	0
3	IPA	A	302	4/4	0.95	0.14	29,29,31,38	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around NAP B 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around NAP A 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around NAP D 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.