



## wwPDB EM Validation Summary Report ⓘ

Dec 8, 2022 – 03:05 PM EST

PDB ID : 7TQL  
EMDB ID : EMD-26067  
Title : CryoEM structure of the human 40S small ribosomal subunit in complex with translation initiation factors eIF1A and eIF5B.  
Authors : Lapointe, C.P.; Grosely, R.; Sokabe, M.; Alvarado, C.; Wang, J.; Montabana, E.; Villa, N.; Shin, B.; Dever, T.; Fraser, C.; Fernandez, I.S.; Puglisi, J.D.  
Deposited on : 2022-01-26  
Resolution : 3.20 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev43  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
MolProbity : 4.02b-467  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
MapQ : 1.9.9  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.31.2


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*ELECTRON MICROSCOPY*

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826
RNA backbone	4643	859

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion  $< 40\%$ ). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	1	619	19% (red), 97% (green), 0% (yellow), 0% (grey), 0% (orange)
2	2	1656	5% (red), 77% (green), 22% (yellow), 0% (grey), 0% (orange)
3	3	75	25% (red), 68% (green), 31% (yellow), 0% (grey), 0% (orange)
4	4	90	11% (red), 91% (green), 9% (yellow), 0% (grey), 0% (orange)
5	B	419	0% (red), 49% (green), 51% (grey), 0% (yellow), 0% (orange)
5	C	419	0% (red), 51% (green), 49% (grey), 0% (yellow), 0% (orange)
6	D	218	0% (red), 98% (green), 0% (yellow), 0% (grey), 0% (orange)
7	E	258	0% (red), 98% (green), 0% (yellow), 0% (grey), 0% (orange)






Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
8	F	225	99%
9	G	226	96%
10	H	185	99%
11	I	205	97%
12	J	180	98%
13	K	189	97%
14	L	147	98%
15	M	97	99%
16	N	141	100%
17	O	120	99%
18	P	134	99%
19	Q	115	100%
20	R	139	100%
21	S	132	97%
22	T	140	98%
23	U	139	99%
24	V	101	96%
25	W	129	99%
26	X	141	99%
27	Y	123	98%
28	Z	82	100%
29	a	72	94%
30	b	82	96%
31	c	99	100%
32	d	58	98%

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
33	e	51	 10% 94% 6%
34	f	54	 100%
35	g	65	 35% 98%
36	h	19	 5% 100%
37	j	314	 96%

## 2 Entry composition [i](#)

There are 40 unique types of molecules in this entry. The entry contains 80976 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Eukaryotic translation initiation factor 5B.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	1	619	4901	3123	844	912	22	0	0

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
1	624	THR	ASN	engineered mutation	UNP O60841
1	625	GLU	-	insertion	UNP O60841
1	626	LYS	THR	engineered mutation	UNP O60841
1	627	ASP	GLU	engineered mutation	UNP O60841
1	906	THR	LEU	engineered mutation	UNP O60841

- Molecule 2 is a RNA chain called 18S ribosomal RNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	P		
2	2	1656	35361	15784	6351	11570	1656	0	0

- Molecule 3 is a RNA chain called Human Met-tRNA<sup>i</sup>Met.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	P		
3	3	75	1607	717	298	517	75	0	0

- Molecule 4 is a protein called Translation initiation factor eIF1A.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
4	4	90	727	459	132	132	4	0	0

- Molecule 5 is a protein called ribosomal protein uS2.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
5	B	206	1624	1035	287	294	8	0	0
5	C	213	1729	1098	309	308	14	0	0

- Molecule 6 is a protein called 40S ribosomal protein S2.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
6	D	218	1682	1090	289	293	10	0	0

- Molecule 7 is a protein called 40S ribosomal protein S4, X isoform.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
7	E	258	2050	1311	381	350	8	0	0

- Molecule 8 is a protein called 40S ribosomal protein S3.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
8	F	225	1748	1115	315	311	7	0	0

- Molecule 9 is a protein called 40S ribosomal protein S6.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
9	G	226	1831	1144	365	315	7	0	0

- Molecule 10 is a protein called 40S ribosomal protein S7.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
10	H	185	1492	952	274	265	1	0	0

- Molecule 11 is a protein called 40S ribosomal protein S8.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
11	I	205	1682	1056	331	290	5	0	0

- Molecule 12 is a protein called 40S ribosomal protein S9.

Mol	Chain	Residues	Atoms					AltConf	Trace
12	J	180	Total	C	N	O	S	0	0
			1499	955	300	242	2		

- Molecule 13 is a protein called 40S ribosomal protein S5.

Mol	Chain	Residues	Atoms					AltConf	Trace
13	K	189	Total	C	N	O	S	0	0
			1495	934	284	270	7		

- Molecule 14 is a protein called 40S ribosomal protein S11.

Mol	Chain	Residues	Atoms					AltConf	Trace
14	L	147	Total	C	N	O	S	0	0
			1204	767	225	206	6		

- Molecule 15 is a protein called 40S ribosomal protein S10.

Mol	Chain	Residues	Atoms					AltConf	Trace
15	M	97	Total	C	N	O	S	0	0
			816	533	144	133	6		

- Molecule 16 is a protein called ribosomal protein uS15.

Mol	Chain	Residues	Atoms					AltConf	Trace
16	N	141	Total	C	N	O	S	0	0
			1152	740	220	191	1		

- Molecule 17 is a protein called ribosomal protein eS12.

Mol	Chain	Residues	Atoms					AltConf	Trace
17	O	120	Total	C	N	O	S	0	0
			929	583	164	174	8		

- Molecule 18 is a protein called ribosomal protein uS11.

Mol	Chain	Residues	Atoms					AltConf	Trace
18	P	134	Total	C	N	O	S	0	0
			998	610	197	185	6		

- Molecule 19 is a protein called 40S ribosomal protein S15.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
19	Q	115	950	604	176	163	7	0	0

- Molecule 20 is a protein called 40S ribosomal protein S16.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
20	R	139	1109	704	210	192	3	0	0

- Molecule 21 is a protein called ribosomal protein eS17.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
21	S	132	1066	669	199	194	4	0	0

- Molecule 22 is a protein called ribosomal protein uS13.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
22	T	140	1162	731	234	196	1	0	0

- Molecule 23 is a protein called 40S ribosomal protein S19.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
23	U	139	1083	678	208	194	3	0	0

- Molecule 24 is a protein called ribosomal protein uS10.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
24	V	101	803	504	153	142	4	0	0

- Molecule 25 is a protein called 40S ribosomal protein S15a.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
25	W	129	1034	659	193	176	6	0	0

- Molecule 26 is a protein called 40S ribosomal protein S23.



Mol	Chain	Residues	Atoms					AltConf	Trace
26	X	141	Total	C	N	O	S	0	0
			1098	693	219	183	3		

- Molecule 27 is a protein called Isoform 3 of 40S ribosomal protein S24.

Mol	Chain	Residues	Atoms					AltConf	Trace
27	Y	123	Total	C	N	O	S	0	0
			1006	637	197	167	5		

- Molecule 28 is a protein called 40S ribosomal protein S21.

Mol	Chain	Residues	Atoms					AltConf	Trace
28	Z	82	Total	C	N	O	S	0	0
			625	384	116	120	5		

- Molecule 29 is a protein called ribosomal protein eS25.

Mol	Chain	Residues	Atoms					AltConf	Trace
29	a	72	Total	C	N	O	S	0	0
			574	368	104	101	1		

- Molecule 30 is a protein called 40S ribosomal protein S27.

Mol	Chain	Residues	Atoms					AltConf	Trace
30	b	82	Total	C	N	O	S	0	0
			640	402	118	113	7		

- Molecule 31 is a protein called 40S ribosomal protein S26.

Mol	Chain	Residues	Atoms					AltConf	Trace
31	c	99	Total	C	N	O	S	0	0
			792	492	165	130	5		

- Molecule 32 is a protein called ribosomal protein eS28.

Mol	Chain	Residues	Atoms					AltConf	Trace
32	d	58	Total	C	N	O	S	0	0
			449	274	86	87	2		

- Molecule 33 is a protein called ribosomal protein eS30.

Mol	Chain	Residues	Atoms					AltConf	Trace
33	e	51	Total	C	N	O	S	0	0
			391	237	88	65	1		

- Molecule 34 is a protein called ribosomal protein uS14.

Mol	Chain	Residues	Atoms					AltConf	Trace
34	f	54	Total	C	N	O	S	0	0
			455	284	93	73	5		

- Molecule 35 is a protein called 40S ribosomal protein S27a.

Mol	Chain	Residues	Atoms					AltConf	Trace
35	g	65	Total	C	N	O	S	0	0
			529	333	99	90	7		


- Molecule 36 is a protein called ribosomal protein eL41.

Mol	Chain	Residues	Atoms					AltConf	Trace
36	h	19	Total	C	N	O	S	0	0
			185	114	50	19	2		

- Molecule 37 is a protein called Receptor of activated protein C kinase 1.


Mol	Chain	Residues	Atoms					AltConf	Trace
37	j	314	Total	C	N	O	S	0	0
			2440	1537	425	466	12		

- Molecule 38 is PHOSPHOAMINOPHOSPHONIC ACID-GUANYLATE ESTER (three-letter code: GNP) (formula:  $C_{10}H_{17}N_6O_{13}P_3$ ) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					AltConf
			Total	C	N	O	P	
38	1	1	32	10	6	13	3	0

- Molecule 39 is GUANOSINE-5'-MONOPHOSPHATE (three-letter code: 5GP) (formula:  $C_{10}H_{14}N_5O_8P$ ).



Mol	Chain	Residues	Atoms					AltConf
			Total	C	N	O	P	
39	3	1	23	10	5	7	1	0


- Molecule 40 is ZINC ION (three-letter code: ZN) (formula: Zn).

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>AltConf</b>
40	c	1	Total 1	Zn 1	0
40	f	1	Total 1	Zn 1	0
40	g	1	Total 1	Zn 1	0


### 3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Eukaryotic translation initiation factor 5B




- Molecule 2: 18S ribosomal RNA








• Molecule 3: Human Met-tRNAiMet




• Molecule 4: Translation initiation factor eIF1A





• Molecule 5: ribosomal protein uS2







- Molecule 10: 40S ribosomal protein S7




- Molecule 11: 40S ribosomal protein S8




- Molecule 12: 40S ribosomal protein S9



- Molecule 13: 40S ribosomal protein S9




- Molecule 14: 40S ribosomal protein S5



- Molecule 15: 40S ribosomal protein S10









- Molecule 16: ribosomal protein uS15

Chain N: 100%




- Molecule 17: ribosomal protein eS12

Chain O: 26% 99%



- Molecule 18: ribosomal protein uS11

Chain P: 99%




- Molecule 19: 40S ribosomal protein S15

Chain Q: 100%




- Molecule 20: 40S ribosomal protein S16

Chain R: 100%




- Molecule 21: ribosomal protein eS17

Chain S: 17% 97%



- Molecule 22: ribosomal protein uS13

Chain T:  98%




- Molecule 23: 40S ribosomal protein S19

Chain U:  99%



- Molecule 24: ribosomal protein uS10

Chain V:  96%




- Molecule 25: 40S ribosomal protein S15a

Chain W:  99%




- Molecule 26: 40S ribosomal protein S23

Chain X:  99%




- Molecule 27: Isoform 3 of 40S ribosomal protein S24

Chain Y:  98%




- Molecule 28: 40S ribosomal protein S21

Chain Z:  100%



- Molecule 29: ribosomal protein eS25

Chain a:  94% 6%




- Molecule 30: 40S ribosomal protein S27

Chain b:  96% .



- Molecule 31: 40S ribosomal protein S26

Chain c:  100%





- Molecule 32: ribosomal protein eS28

Chain d:  98% .



- Molecule 33: ribosomal protein eS30

Chain e:  10% 94% 6%




- Molecule 34: ribosomal protein uS14

Chain f:  100%




- Molecule 35: 40S ribosomal protein S27a


Chain g:  35% 98% .



- Molecule 36: ribosomal protein eL41



- Molecule 37: Receptor of activated protein C kinase 1



## 4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	190000	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	70	Depositor
Minimum defocus (nm)	500	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	2.534	Depositor
Minimum map value	-0.031	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.023	Depositor
Recommended contour level	0.1	Depositor
Map size ( $\text{\AA}$ )	628.8, 628.8, 628.8	wwPDB
Map dimensions	480, 480, 480	wwPDB
Map angles ( $^\circ$ )	90.0, 90.0, 90.0	wwPDB
Pixel spacing ( $\text{\AA}$ )	1.31, 1.31, 1.31	Depositor

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: 5GP, ZN, GNP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	1	0.71	0/4983	0.81	1/6718 (0.0%)
2	2	0.51	0/39542	0.78	9/61622 (0.0%)
3	3	0.53	0/1798	0.79	1/2802 (0.0%)
4	4	0.75	0/737	0.93	1/986 (0.1%)
5	B	0.69	0/1661	0.76	0/2259
5	C	0.69	0/1756	0.76	0/2350
6	D	0.72	0/1718	0.78	0/2322
7	E	0.76	0/2092	0.81	3/2816 (0.1%)
8	F	0.73	1/1776 (0.1%)	0.79	0/2392
9	G	0.72	0/1854	0.89	2/2469 (0.1%)
10	H	0.71	0/1515	0.76	0/2030
11	I	0.68	0/1711	0.78	1/2282 (0.0%)
12	J	0.73	0/1524	0.81	0/2035
13	K	0.77	0/1516	0.79	0/2037
14	L	0.68	0/1225	0.81	1/1640 (0.1%)
15	M	0.67	0/840	0.75	0/1133
16	N	0.69	0/1176	0.77	0/1580
17	O	0.73	0/939	0.80	0/1261
18	P	0.75	0/1011	0.85	0/1356
19	Q	0.68	0/968	0.75	0/1294
20	R	0.70	0/1126	0.77	0/1506
21	S	0.75	0/1080	0.79	1/1449 (0.1%)
22	T	0.71	0/1180	0.82	0/1581
23	U	0.70	0/1101	0.77	0/1477
24	V	0.71	0/813	0.82	0/1092
25	W	0.69	0/1051	0.77	0/1406
26	X	0.69	0/1116	0.80	0/1490
27	Y	0.71	0/1023	0.80	0/1359
28	Z	0.71	0/631	0.78	0/844
29	a	0.74	0/580	0.84	0/780
30	b	0.76	0/653	0.87	1/876 (0.1%)
31	c	0.69	0/805	0.82	0/1079

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
32	d	0.70	0/451	0.82	0/604
33	e	0.71	0/393	0.91	0/516
34	f	0.71	0/466	0.81	0/618
35	g	0.73	0/540	0.80	0/718
36	h	0.64	0/186	0.89	0/236
37	j	0.73	0/2497	0.81	2/3399 (0.1%)
All	All	0.62	1/86034 (0.0%)	0.79	23/124414 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	1	0	2
33	e	0	1
All	All	0	3

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	F	192	TRP	CB-CG	-5.60	1.40	1.50

The worst 5 of 23 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
9	G	134	GLY	C-N-CD	-11.96	94.30	120.60
37	j	103	GLY	N-CA-C	7.19	131.07	113.10
2	2	1832	A	C2'-C3'-O3'	6.07	123.42	113.70
2	2	1231	C	C2'-C3'-O3'	6.02	123.34	113.70
7	E	139	LEU	CA-CB-CG	5.89	128.86	115.30

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	1	807	LEU	Peptide
1	1	921	LYS	Mainchain
33	e	3	ALA	Peptide

## 5.2 Too-close contacts [i](#)

Due to software issues we are unable to calculate clashes - this section is therefore empty.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	1	617/619 (100%)	518 (84%)	87 (14%)	12 (2%)	8	39
4	4	88/90 (98%)	72 (82%)	14 (16%)	2 (2%)	6	34
5	B	204/419 (49%)	196 (96%)	7 (3%)	1 (0%)	29	67
5	C	211/419 (50%)	198 (94%)	13 (6%)	0	100	100
6	D	216/218 (99%)	207 (96%)	8 (4%)	1 (0%)	29	67
7	E	256/258 (99%)	240 (94%)	16 (6%)	0	100	100
8	F	223/225 (99%)	214 (96%)	9 (4%)	0	100	100
9	G	224/226 (99%)	211 (94%)	10 (4%)	3 (1%)	12	47
10	H	183/185 (99%)	174 (95%)	8 (4%)	1 (0%)	29	67
11	I	203/205 (99%)	186 (92%)	15 (7%)	2 (1%)	15	54
12	J	178/180 (99%)	168 (94%)	9 (5%)	1 (1%)	25	64
13	K	187/189 (99%)	173 (92%)	13 (7%)	1 (0%)	29	67
14	L	145/147 (99%)	140 (97%)	5 (3%)	0	100	100
15	M	95/97 (98%)	86 (90%)	8 (8%)	1 (1%)	14	51
16	N	139/141 (99%)	130 (94%)	9 (6%)	0	100	100
17	O	118/120 (98%)	111 (94%)	6 (5%)	1 (1%)	19	58
18	P	132/134 (98%)	123 (93%)	8 (6%)	1 (1%)	19	58
19	Q	113/115 (98%)	108 (96%)	5 (4%)	0	100	100
20	R	137/139 (99%)	131 (96%)	6 (4%)	0	100	100
21	S	130/132 (98%)	123 (95%)	6 (5%)	1 (1%)	19	58
22	T	138/140 (99%)	130 (94%)	7 (5%)	1 (1%)	22	61

*Continued on next page...*



Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
23	U	137/139 (99%)	132 (96%)	5 (4%)	0	100	100
24	V	99/101 (98%)	94 (95%)	5 (5%)	0	100	100
25	W	127/129 (98%)	122 (96%)	5 (4%)	0	100	100
26	X	139/141 (99%)	128 (92%)	10 (7%)	1 (1%)	22	61
27	Y	121/123 (98%)	117 (97%)	4 (3%)	0	100	100
28	Z	80/82 (98%)	73 (91%)	7 (9%)	0	100	100
29	a	70/72 (97%)	66 (94%)	3 (4%)	1 (1%)	11	46
30	b	80/82 (98%)	75 (94%)	5 (6%)	0	100	100
31	c	97/99 (98%)	94 (97%)	3 (3%)	0	100	100
32	d	56/58 (97%)	56 (100%)	0	0	100	100
33	e	47/51 (92%)	45 (96%)	2 (4%)	0	100	100
34	f	52/54 (96%)	48 (92%)	4 (8%)	0	100	100
35	g	63/65 (97%)	54 (86%)	9 (14%)	0	100	100
36	h	17/19 (90%)	17 (100%)	0	0	100	100
37	j	312/314 (99%)	284 (91%)	25 (8%)	3 (1%)	15	54
All	All	5434/5927 (92%)	5044 (93%)	356 (7%)	34 (1%)	29	64

5 of 34 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	1	675	LEU
1	1	729	ILE
9	G	130	PRO
9	G	135	PRO
26	X	86	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	1	544/544 (100%)	535 (98%)	9 (2%)	60	83

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
4	4	76/76 (100%)	71 (93%)	5 (7%)	16	51
5	B	172/366 (47%)	171 (99%)	1 (1%)	86	94
5	C	194/366 (53%)	193 (100%)	1 (0%)	88	95
6	D	182/184 (99%)	179 (98%)	3 (2%)	62	84
7	E	221/221 (100%)	218 (99%)	3 (1%)	67	86
8	F	188/189 (100%)	187 (100%)	1 (0%)	88	95
9	G	197/197 (100%)	193 (98%)	4 (2%)	55	80
10	H	166/166 (100%)	165 (99%)	1 (1%)	86	94
11	I	178/178 (100%)	175 (98%)	3 (2%)	60	83
12	J	160/160 (100%)	158 (99%)	2 (1%)	69	87
13	K	159/159 (100%)	155 (98%)	4 (2%)	47	77
14	L	133/133 (100%)	131 (98%)	2 (2%)	65	85
15	M	88/88 (100%)	88 (100%)	0	100	100
16	N	124/124 (100%)	124 (100%)	0	100	100
17	O	101/101 (100%)	101 (100%)	0	100	100
18	P	103/104 (99%)	103 (100%)	0	100	100
19	Q	104/104 (100%)	104 (100%)	0	100	100
20	R	115/115 (100%)	115 (100%)	0	100	100
21	S	118/119 (99%)	116 (98%)	2 (2%)	60	83
22	T	122/122 (100%)	120 (98%)	2 (2%)	62	84
23	U	110/110 (100%)	109 (99%)	1 (1%)	78	91
24	V	93/93 (100%)	89 (96%)	4 (4%)	29	64
25	W	112/112 (100%)	111 (99%)	1 (1%)	78	91
26	X	113/113 (100%)	113 (100%)	0	100	100
27	Y	107/107 (100%)	104 (97%)	3 (3%)	43	74
28	Z	66/66 (100%)	66 (100%)	0	100	100
29	a	64/64 (100%)	61 (95%)	3 (5%)	26	62
30	b	74/74 (100%)	72 (97%)	2 (3%)	44	75
31	c	86/86 (100%)	86 (100%)	0	100	100
32	d	51/51 (100%)	50 (98%)	1 (2%)	55	80
33	e	36/37 (97%)	34 (94%)	2 (6%)	21	57

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
34	f	48/48 (100%)	48 (100%)	0	100	100
35	g	58/58 (100%)	57 (98%)	1 (2%)	60	83
36	h	18/18 (100%)	18 (100%)	0	100	100
37	j	272/272 (100%)	263 (97%)	9 (3%)	38	71
All	All	4753/5125 (93%)	4683 (98%)	70 (2%)	66	85

5 of 70 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
32	d	57	THR
33	e	25	LYS
37	j	62	HIS
9	G	133	LEU
9	G	116	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 19 such sidechains are listed below:

Mol	Chain	Res	Type
20	R	97	GLN
30	b	51	GLN
33	e	22	GLN
27	Y	85	ASN
9	G	56	ASN

### 5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	2	1646/1656 (99%)	359 (21%)	36 (2%)
3	3	74/75 (98%)	22 (29%)	2 (2%)
All	All	1720/1731 (99%)	381 (22%)	38 (2%)

5 of 381 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	2	2	A
2	2	3	C
2	2	4	C
2	2	11	A
2	2	17	C

5 of 38 RNA pucker outliers are listed below:

Mol	Chain	Res	Type
2	2	1502	C
2	2	1832	A
2	2	1519	U
2	2	1565	C
3	3	64	C

#### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

#### 5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 3 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
38	GNP	1	1301	-	29,34,34	1.45	6 (20%)	33,54,54	2.28	6 (18%)
39	5GP	3	101	-	18,25,26	1.00	2 (11%)	19,37,40	0.65	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
38	GNP	1	1301	-	-	5/14/38/38	0/3/3/3
39	5GP	3	101	-	-	2/3/25/26	0/3/3/3

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
38	1	1301	GNP	C6-N1	3.96	1.39	1.33
38	1	1301	GNP	PG-O1G	3.24	1.51	1.46
38	1	1301	GNP	PB-O1B	2.86	1.50	1.46
39	3	101	5GP	C5-C6	-2.53	1.42	1.47
38	1	1301	GNP	PB-O2B	-2.24	1.50	1.56

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
38	1	1301	GNP	C5-C6-N1	-8.74	111.48	123.43
38	1	1301	GNP	C2-N1-C6	5.98	125.43	115.93
38	1	1301	GNP	O2B-PB-O1B	3.91	118.12	109.92
38	1	1301	GNP	C2-N3-C4	-3.25	111.65	115.36
38	1	1301	GNP	N3-C2-N1	-2.59	123.77	127.22

There are no chirality outliers.



5 of 7 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
38	1	1301	GNP	PB-N3B-PG-O1G
38	1	1301	GNP	PG-N3B-PB-O1B
38	1	1301	GNP	C5'-O5'-PA-O3A
38	1	1301	GNP	C5'-O5'-PA-O2A
39	3	101	5GP	O4'-C4'-C5'-O5'

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
2	2	10
33	e	1

The worst 5 of 11 chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	2	834:C	O3'	841:G	P	18.33
1	2	130:G	O3'	140:U	P	17.28
1	2	1416:C	O3'	1424:G	P	17.27
1	2	747:U	O3'	795:A	P	16.72
1	2	1761:U	O3'	1771:G	P	15.57


## 6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-26067. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

### 6.1 Orthogonal projections [i](#)


#### 6.1.1 Primary map



X




Y




Z


#### 6.1.2 Raw map



X



Y



Z

The images above show the map projected in three orthogonal directions.



## 6.2 Central slices [i](#)

### 6.2.1 Primary map



X Index: 240




Y Index: 240




Z Index: 240


### 6.2.2 Raw map



X Index: 240



Y Index: 240



Z Index: 240

The images above show central slices of the map in three orthogonal directions.

## 6.3 Largest variance slices [i](#)

### 6.3.1 Primary map



X Index: 259




Y Index: 238




Z Index: 252


### 6.3.2 Raw map



X Index: 259



Y Index: 238




Z Index: 223

The images above show the largest variance slices of the map in three orthogonal directions.


## 6.4 Orthogonal surface views [i](#)

### 6.4.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.1. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

### 6.4.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


## 6.5 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


### 6.5.1 emd\_26067\_msk\_1.map [i](#)



X



Y




Z

## 7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

### 7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


## 7.2 Volume estimate [i](#)



The volume at the recommended contour level is 478  $\text{nm}^3$ ; this corresponds to an approximate mass of 432 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

### 7.3 Rotationally averaged power spectrum i




\*Reported resolution corresponds to spatial frequency of 0.312 Å<sup>-1</sup>

## 8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

### 8.1 FSC [i](#)



\*Reported resolution corresponds to spatial frequency of 0.312  $\text{\AA}^{-1}$



## 8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.20	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	6.92	10.32	7.44

\*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 6.92 differs from the reported value 3.2 by more than 10 %

## 9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-26067 and PDB model 7TQL. Per-residue inclusion information can be found in section 3 on page 13.

### 9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.1 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

## 9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

## 9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.1).







































































## 9.4 Atom inclusion [i](#)



At the recommended contour level, 91% of all backbone atoms, 80% of all non-hydrogen atoms, are inside the map.

## 9.5 Map-model fit summary









The table lists the average atom inclusion at the recommended contour level (0.1) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.7992	 0.4120
1	 0.5782	 0.2370
2	 0.8907	 0.4520
3	 0.6061	 0.2690
4	 0.6201	 0.3420
B	 0.7922	 0.4240
C	 0.7604	 0.4190
D	 0.8067	 0.4500
E	 0.7914	 0.4410
F	 0.7736	 0.4140
G	 0.7032	 0.3630
H	 0.5188	 0.2560
I	 0.7751	 0.4120
J	 0.7728	 0.4210
K	 0.8006	 0.4380
L	 0.7382	 0.4230
M	 0.7701	 0.3900
N	 0.7769	 0.4200
O	 0.5202	 0.2170
P	 0.7977	 0.4340
Q	 0.7885	 0.4150
R	 0.8295	 0.4440
S	 0.6525	 0.3360
T	 0.7650	 0.4100
U	 0.8025	 0.4380
V	 0.7481	 0.4080
W	 0.8139	 0.4520
X	 0.8067	 0.4540
Y	 0.7613	 0.4110
Z	 0.7545	 0.4130
a	 0.7273	 0.4050
b	 0.7978	 0.4320
c	 0.8110	 0.4510
d	 0.7816	 0.4320
e	 0.7420	 0.4190



*Continued on next page...*

*Continued from previous page...*

Chain	Atom inclusion	Q-score
f	 0.8402	 0.4590
g	 0.4893	 0.2410
h	 0.7321	 0.4340
j	 0.7562	 0.3800