



wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 23, 2023 – 07:18 PM EDT

PDB ID : 5TRU
Title : Structure of the first-in-class checkpoint inhibitor Ipilimumab bound to human CTLA-4
Authors : Ramagopal, U.A.; Liu, W.; Garrett-Thomson, S.C.; Yan, Q.; Srinivasan, M.; Wong, S.C.; Bell, A.; Mankikar, S.; Rangan, V.S.; Deshpande, S.; Bonanno, J.B.; Korman, A.J.; Almo, S.C.
Deposited on : 2016-10-27
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.35.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2092 (3.00-3.00)
Ramachandran outliers	138981	2333 (3.00-3.00)
Sidechain outliers	138945	2336 (3.00-3.00)
RSRZ outliers	127900	1990 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	L	215	91% (green), 8% (yellow), 1% (orange), 0% (red), 0% (grey)
1	l	215	94% (green), 4% (yellow), 2% (orange), 0% (red), 0% (grey)
2	H	225	91% (green), 7% (yellow), 2% (orange), 0% (red), 0% (grey)
2	h	225	84% (green), 10% (yellow), 6% (grey)
3	C	118	90% (green), 7% (yellow), 3% (orange), 0% (red), 0% (grey)
3	c	118	94% (green), 4% (yellow), 2% (orange), 0% (red), 0% (grey)

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 8036 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ipilimumab Fab light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	L	213	Total 1627	C 1023	N 275	O 325	S 4	0	0	0
1	l	212	Total 1579	C 987	N 271	O 317	S 4	0	0	0

- Molecule 2 is a protein called Ipilimumab Fab heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	H	215	Total 1619	C 1030	N 270	O 313	S 6	0	0	0
2	h	212	Total 1589	C 1011	N 266	O 306	S 6	0	0	0


- Molecule 3 is a protein called Cytotoxic T-lymphocyte protein 4.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	c	115	Total 816	C 515	N 134	O 158	S 9	0	0	0
3	C	114	Total 806	C 510	N 132	O 155	S 9	0	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Ipilimumab Fab light chain

Chain L:  91% 8%



- Molecule 1: Ipilimumab Fab light chain

Chain l:  94%





- Molecule 2: Ipilimumab Fab heavy chain

Chain H:  91%




- Molecule 2: Ipilimumab Fab heavy chain

Chain h:  84% 10% 6%





- Molecule 3: Cytotoxic T-lymphocyte protein 4

Chain c:  94%



- Molecule 3: Cytotoxic T-lymphocyte protein 4

Chain C:  90% 7%



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	95.84Å 197.50Å 148.12Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	34.91 – 3.00 34.91 – 3.00	Depositor EDS
% Data completeness (in resolution range)	99.2 (34.91-3.00) 99.3 (34.91-3.00)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.67 (at 3.00Å)	Xtrriage
Refinement program	REFMAC 5	Depositor
R, R_{free}	0.203 , 0.268 0.206 , 0.270	Depositor DCC
R_{free} test set	1447 reflections (5.10%)	wwPDB-VP
Wilson B-factor (Å ²)	62.1	Xtrriage
Anisotropy	0.607	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.27 , 13.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	8036	wwPDB-VP
Average B, all atoms (Å ²)	71.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.45% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	L	0.51	0/1665	0.73	0/2263
1	l	0.47	0/1616	0.68	0/2200
2	H	0.54	0/1661	0.75	0/2268
2	h	0.53	0/1631	0.70	0/2229
3	C	0.49	0/819	0.67	0/1118
3	c	0.47	0/828	0.65	0/1129
All	All	0.51	0/8220	0.71	0/11207

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

Due to software issues we are unable to calculate clashes - this section is therefore empty.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	L	211/215 (98%)	199 (94%)	11 (5%)	1 (0%)	29 68

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	l	210/215 (98%)	201 (96%)	7 (3%)	2 (1%)	15	53
2	H	211/225 (94%)	196 (93%)	14 (7%)	1 (0%)	29	68
2	h	208/225 (92%)	196 (94%)	9 (4%)	3 (1%)	11	43
3	C	110/118 (93%)	104 (94%)	6 (6%)	0	100	100
3	c	111/118 (94%)	100 (90%)	11 (10%)	0	100	100
All	All	1061/1116 (95%)	996 (94%)	58 (6%)	7 (1%)	22	60

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	H	101	TRP
2	h	101	TRP
1	l	139	ASN
2	h	41	PRO
1	L	139	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	180/185 (97%)	164 (91%)	16 (9%)	9	35
1	l	166/185 (90%)	159 (96%)	7 (4%)	30	66
2	H	174/191 (91%)	163 (94%)	11 (6%)	18	51
2	h	173/191 (91%)	152 (88%)	21 (12%)	5	21
3	C	78/97 (80%)	70 (90%)	8 (10%)	7	28
3	c	82/97 (84%)	78 (95%)	4 (5%)	25	61
All	All	853/946 (90%)	786 (92%)	67 (8%)	12	41

5 of 67 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	c	68	CYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	C	4	HIS
3	C	87	MET
2	H	186	VAL
2	H	184	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
3	c	4	HIS
3	c	75	ASN
2	H	204	ASN
1	l	90	GLN
1	l	138	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	L	213/215 (99%)	-0.55	0	100 100	45, 59, 73, 81	0
1	l	212/215 (98%)	-0.28	0	100 100	45, 68, 132, 149	0
2	H	215/225 (95%)	-0.43	0	100 100	48, 63, 91, 109	0
2	h	212/225 (94%)	-0.32	1 (0%)	91 75	46, 65, 108, 125	0
3	C	114/118 (96%)	-0.28	1 (0%)	84 63	60, 84, 108, 126	0
3	c	115/118 (97%)	-0.17	0	100 100	57, 84, 112, 134	0
All	All	1081/1116 (96%)	-0.36	2 (0%)	95 87	45, 66, 115, 149	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	h	133	SER	2.9
3	C	23	CYS	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.