



# Full wwPDB NMR Structure Validation Report ⓘ

Feb 19, 2022 – 10:56 PM EST

PDB ID : 1TUQ  
Title : NMR Structure Analysis of the B-DNA Dodecamer CTCtCACGTGGAG with a tricyclic cytosin base analogue  
Authors : Engman, K.C.; Sandin, P.; Osborne, S.; Brown, T.; Billeter, M.; Lincoln, P.; Norden, B.; Albinsson, B.; Wilhelmsson, L.M.  
Deposited on : 2004-06-25

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
RCI : v\_1n\_11\_5\_13\_A (Berjanski et al., 2005)  
PANAV : Wang et al. (2010)  
ShiftChecker : 2.26  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.26


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*SOLUTION NMR*



The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	12	
1	B	12	

## 2 Ensemble composition and analysis

This entry contains 10 models. This entry does not contain polypeptide chains, therefore identification of well-defined residues and clustering analysis are not possible. All residues are included in the validation scores.

### 3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 778 atoms, of which 272 are hydrogens and 0 are deuteriums.

- Molecule 1 is a DNA chain called 5'-D(P\*CP\*TP\*CP\*(TC1)P\*AP\*CP\*GP\*TP\*GP\*GP\*AP\*G)-3'.

Mol	Chain	Residues	Atoms							Trace
			Total	C	H	N	O	P	S	
1	A	12	Total 389	C 122	H 136	N 46	O 72	P 12	S 1	0
1	B	12	Total 389	C 122	H 136	N 46	O 72	P 12	S 1	0


## 4 Residue-property plots

### 4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: 5'-D(P\*CP\*TP\*CP\*(TC1)P\*AP\*CP\*GP\*TP\*GP\*GP\*AP\*G)-3'

Chain A: 



- Molecule 1: 5'-D(P\*CP\*TP\*CP\*(TC1)P\*AP\*CP\*GP\*TP\*GP\*GP\*AP\*G)-3'

Chain B: 




### 4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.

#### 4.2.1 Score per residue for model 1


- Molecule 1: 5'-D(P\*CP\*TP\*CP\*(TC1)P\*AP\*CP\*GP\*TP\*GP\*GP\*AP\*G)-3'

Chain A: 



- Molecule 1: 5'-D(P\*CP\*TP\*CP\*(TC1)P\*AP\*CP\*GP\*TP\*GP\*GP\*AP\*G)-3'

Chain B: 



### 4.2.2 Score per residue for model 2

- Molecule 1: 5'-D(P\*CP\*TP\*CP\*(TC1)P\*AP\*CP\*GP\*TP\*GP\*GP\*AP\*G)-3'

Chain A:  17% 83%

C1 T2 C3 C4 A5 C6 G7 T8 G9 G10 A11 G12

- Molecule 1: 5'-D(P\*CP\*TP\*CP\*(TC1)P\*AP\*CP\*GP\*TP\*GP\*GP\*AP\*G)-3'

Chain B:  8% 92%

C101 T102 C103 C104 A105 C106 G107 T108 G109 G110 A111 G112

### 4.2.3 Score per residue for model 3

- Molecule 1: 5'-D(P\*CP\*TP\*CP\*(TC1)P\*AP\*CP\*GP\*TP\*GP\*GP\*AP\*G)-3'

Chain A:  17% 83%

C1 T2 C3 C4 A5 C6 G7 T8 G9 G10 A11 G12

- Molecule 1: 5'-D(P\*CP\*TP\*CP\*(TC1)P\*AP\*CP\*GP\*TP\*GP\*GP\*AP\*G)-3'

Chain B:  25% 75%

C101 T102 C103 C104 A105 C106 G107 T108 G109 G110 A111 G112

### 4.2.4 Score per residue for model 4

- Molecule 1: 5'-D(P\*CP\*TP\*CP\*(TC1)P\*AP\*CP\*GP\*TP\*GP\*GP\*AP\*G)-3'

Chain A:  25% 75%

C1 T2 C3 C4 A5 C6 G7 T8 G9 G10 A11 G12

- Molecule 1: 5'-D(P\*CP\*TP\*CP\*(TC1)P\*AP\*CP\*GP\*TP\*GP\*GP\*AP\*G)-3'


Chain B:  8% 92%

C101 T102 C103 C104 A105 C106 G107 T108 G109 G110 A111 G112

#### 4.2.5 Score per residue for model 5


- Molecule 1: 5'-D(P\*CP\*TP\*CP\*(TC1)P\*AP\*CP\*GP\*TP\*GP\*GP\*AP\*G)-3'

Chain A:  25% 75%

 C1 T2 C3 C4 A5 C6 G7 T8 G9 G10 A11 G12

- Molecule 1: 5'-D(P\*CP\*TP\*CP\*(TC1)P\*AP\*CP\*GP\*TP\*GP\*GP\*AP\*G)-3'


Chain B:  33% 67%

 C101 T102 C103 C104 A105 C106 G107 T108 G109 G110 A111 G112

#### 4.2.6 Score per residue for model 6

- Molecule 1: 5'-D(P\*CP\*TP\*CP\*(TC1)P\*AP\*CP\*GP\*TP\*GP\*GP\*AP\*G)-3'

Chain A:  8% 92%

 C1 T2 C3 C4 A5 C6 G7 T8 G9 G10 A11 G12

- Molecule 1: 5'-D(P\*CP\*TP\*CP\*(TC1)P\*AP\*CP\*GP\*TP\*GP\*GP\*AP\*G)-3'


Chain B:  8% 92%

 C101 T102 C103 C104 A105 C106 G107 T108 G109 G110 A111 G112

#### 4.2.7 Score per residue for model 7

- Molecule 1: 5'-D(P\*CP\*TP\*CP\*(TC1)P\*AP\*CP\*GP\*TP\*GP\*GP\*AP\*G)-3'

Chain A:  100%

 C1 T2 C3 C4 A5 C6 G7 T8 G9 G10 A11 G12

- Molecule 1: 5'-D(P\*CP\*TP\*CP\*(TC1)P\*AP\*CP\*GP\*TP\*GP\*GP\*AP\*G)-3'

Chain B:  8% 92%

 C101 T102 C103 C104 A105 C106 G107 T108 G109 G110 A111 G112

#### 4.2.8 Score per residue for model 8

- Molecule 1: 5'-D(P\*CP\*TP\*CP\*(TC1)P\*AP\*CP\*GP\*TP\*GP\*GP\*AP\*G)-3'

Chain A:  33% 67%

C1 T2 C3 C4 A5 C6 C7 T8 G9 G10 A11 G12

- Molecule 1: 5'-D(P\*CP\*TP\*CP\*(TC1)P\*AP\*CP\*GP\*TP\*GP\*GP\*AP\*G)-3'

Chain B:  25% 75%

C101 T102 C103 C104 A105 C106 G107 T108 G109 G110 A111 G112

#### 4.2.9 Score per residue for model 9

- Molecule 1: 5'-D(P\*CP\*TP\*CP\*(TC1)P\*AP\*CP\*GP\*TP\*GP\*GP\*AP\*G)-3'

Chain A:  17% 83%

C1 T2 C3 C4 A5 C6 C7 T8 G9 G10 A11 G12

- Molecule 1: 5'-D(P\*CP\*TP\*CP\*(TC1)P\*AP\*CP\*GP\*TP\*GP\*GP\*AP\*G)-3'

Chain B:  25% 75%

C101 T102 C103 C104 A105 C106 G107 T108 G109 G110 A111 G112

#### 4.2.10 Score per residue for model 10

- Molecule 1: 5'-D(P\*CP\*TP\*CP\*(TC1)P\*AP\*CP\*GP\*TP\*GP\*GP\*AP\*G)-3'

Chain A:  17% 83%

C1 T2 C3 C4 A5 C6 C7 T8 G9 G10 A11 G12

- Molecule 1: 5'-D(P\*CP\*TP\*CP\*(TC1)P\*AP\*CP\*GP\*TP\*GP\*GP\*AP\*G)-3'

Chain B:  8% 92%

C101 T102 C103 C104 A105 C106 G107 T108 G109 G110 A111 G112



## 5 Refinement protocol and experimental data overview

The models were refined using the following method: *torsion angle dynamics*.

Of the 100 calculated structures, 10 were deposited, based on the following criterion: *lowest CYANA target function*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	structure solution	1.1
CYANA	refinement	1.1

No chemical shift data was provided.

## 6 Model quality i

### 6.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: TC1

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	1.58±0.01	0±0/253 ( 0.0± 0.0%)	3.15±0.00	33±0/386 ( 8.5± 0.0%)
1	B	1.58±0.01	0±0/253 ( 0.0± 0.0%)	3.14±0.01	33±0/386 ( 8.5± 0.0%)
All	All	1.58	0/5060 ( 0.0%)	3.15	660/7720 ( 8.5%)

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	10	DG	C5-C6-N1	11.97	117.48	111.50	5	10
1	A	9	DG	C5-C6-N1	11.94	117.47	111.50	7	10
1	B	112	DG	C5-C6-N1	11.94	117.47	111.50	1	10
1	B	110	DG	C5-C6-N1	11.91	117.46	111.50	7	10
1	B	109	DG	C5-C6-N1	11.91	117.45	111.50	7	10
1	A	12	DG	C5-C6-N1	11.89	117.45	111.50	8	10
1	A	7	DG	C5-C6-N1	11.88	117.44	111.50	6	10
1	B	107	DG	C5-C6-N1	11.88	117.44	111.50	6	10
1	A	9	DG	C6-N1-C2	-11.17	118.40	125.10	7	10
1	B	112	DG	C6-N1-C2	-11.16	118.40	125.10	1	10
1	A	12	DG	C6-N1-C2	-11.12	118.43	125.10	4	10
1	A	7	DG	C6-N1-C2	-11.10	118.44	125.10	6	10
1	A	10	DG	C6-N1-C2	-11.06	118.46	125.10	5	10
1	B	110	DG	C6-N1-C2	-11.06	118.46	125.10	7	10
1	B	109	DG	C6-N1-C2	-11.05	118.47	125.10	7	10
1	B	107	DG	C6-N1-C2	-11.02	118.49	125.10	8	10
1	A	9	DG	N1-C2-N3	8.91	129.25	123.90	7	10
1	B	112	DG	N1-C2-N3	8.87	129.22	123.90	1	10
1	B	110	DG	N1-C2-N3	8.82	129.19	123.90	1	10
1	A	12	DG	N1-C2-N3	8.82	129.19	123.90	4	10

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	7	DG	N1-C2-N3	8.79	129.17	123.90	6	10
1	B	107	DG	N1-C2-N3	8.72	129.13	123.90	1	10
1	A	10	DG	N1-C2-N3	8.69	129.11	123.90	9	10
1	B	109	DG	N1-C2-N3	8.69	129.11	123.90	1	10
1	A	2	DT	O4'-C4'-C3'	-7.42	101.53	104.50	2	10
1	B	102	DT	O4'-C4'-C3'	-7.36	101.56	104.50	5	10
1	B	111	DA	O4'-C4'-C3'	-7.36	101.56	104.50	4	10
1	B	108	DT	O4'-C4'-C3'	-7.32	101.57	104.50	6	10
1	A	5	DA	O4'-C4'-C3'	-7.32	101.57	104.50	2	10
1	B	105	DA	O4'-C4'-C3'	-7.32	101.57	104.50	8	10
1	A	12	DG	O4'-C4'-C3'	-7.32	101.57	104.50	2	10
1	A	3	DC	O4'-C4'-C3'	-7.31	101.58	104.50	9	10
1	B	103	DC	O4'-C4'-C3'	-7.30	101.58	104.50	9	10
1	A	10	DG	O4'-C4'-C3'	-7.30	101.58	104.50	3	10
1	B	106	DC	O4'-C4'-C3'	-7.30	101.58	104.50	4	10
1	A	11	DA	O4'-C4'-C3'	-7.29	101.58	104.50	2	10
1	B	112	DG	O4'-C4'-C3'	-7.28	101.59	104.50	10	10
1	B	107	DG	O4'-C4'-C3'	-7.27	101.59	104.50	7	10
1	A	1	DC	O4'-C4'-C3'	-7.27	101.59	104.50	2	10
1	B	110	DG	O4'-C4'-C3'	-7.27	101.59	104.50	1	10
1	A	6	DC	O4'-C4'-C3'	-7.27	101.59	104.50	5	10
1	A	9	DG	O4'-C4'-C3'	-7.27	101.59	104.50	4	10
1	B	101	DC	O4'-C4'-C3'	-7.26	101.60	104.50	5	10
1	A	7	DG	O4'-C4'-C3'	-7.24	101.60	104.50	2	10
1	B	109	DG	O4'-C4'-C3'	-7.24	101.60	104.50	9	10
1	A	8	DT	O4'-C4'-C3'	-7.23	101.61	104.50	10	10
1	A	8	DT	C2-N3-C4	-6.17	123.50	127.20	6	10
1	A	2	DT	C2-N3-C4	-6.17	123.50	127.20	8	10
1	B	108	DT	C2-N3-C4	-6.15	123.51	127.20	2	10
1	B	102	DT	C2-N3-C4	-6.13	123.52	127.20	9	10
1	A	12	DG	N1-C2-N2	-5.57	111.19	116.20	4	10
1	A	7	DG	N1-C2-N2	-5.55	111.20	116.20	6	10
1	B	107	DG	N1-C2-N2	-5.54	111.22	116.20	10	10
1	A	10	DG	N1-C2-N2	-5.52	111.23	116.20	5	10
1	A	9	DG	N1-C2-N2	-5.51	111.24	116.20	7	10
1	B	110	DG	N1-C2-N2	-5.51	111.24	116.20	1	10
1	B	109	DG	N1-C2-N2	-5.49	111.26	116.20	9	10
1	B	112	DG	N1-C2-N2	-5.49	111.26	116.20	1	10
1	A	10	DG	C5-C6-O6	-5.48	125.31	128.60	5	10
1	B	112	DG	C5-C6-O6	-5.45	125.33	128.60	5	10
1	A	12	DG	C5-C6-O6	-5.45	125.33	128.60	7	10

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	9	DG	C5-C6-O6	-5.44	125.33	128.60	7	10
1	B	107	DG	C5-C6-O6	-5.43	125.34	128.60	2	10
1	A	7	DG	C5-C6-O6	-5.41	125.36	128.60	4	10
1	B	109	DG	C5-C6-O6	-5.41	125.36	128.60	9	10
1	B	110	DG	C5-C6-O6	-5.39	125.36	128.60	5	10

There are no chirality outliers.

There are no planarity outliers.

## 6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	253	136	137	15±3
1	B	253	136	137	14±4
All	All	5060	2720	2740	264

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 34.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:2:DT:C2	1:A:3:DC:C5	0.79	2.70	10	9
1:B:102:DT:C2	1:B:103:DC:C5	0.78	2.72	2	10
1:B:108:DT:C4	1:B:109:DG:C6	0.73	2.77	9	8
1:A:8:DT:C4	1:A:9:DG:C6	0.73	2.76	2	5
1:A:9:DG:C6	1:A:10:DG:C6	0.67	2.81	9	8
1:B:109:DG:C6	1:B:110:DG:C6	0.65	2.84	6	8
1:B:111:DA:C4	1:B:112:DG:N7	0.64	2.65	10	5
1:A:7:DG:C8	1:A:8:DT:C7	0.63	2.82	7	6
1:A:11:DA:C4	1:A:12:DG:N7	0.62	2.68	1	4
1:A:2:DT:C4	1:A:3:DC:N4	0.61	2.67	2	4
1:A:9:DG:C6	1:A:10:DG:O6	0.60	2.55	6	9
1:B:108:DT:C4	1:B:109:DG:O6	0.60	2.55	4	6
1:B:109:DG:C6	1:B:110:DG:O6	0.59	2.55	9	8

Continued on next page...

*Continued from previous page...*

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:B:108:DT:N3	1:B:109:DG:C6	0.59	2.71	2	5
1:A:8:DT:C4	1:A:9:DG:O6	0.56	2.58	4	4
1:B:111:DA:C6	1:B:112:DG:O6	0.55	2.59	10	2
1:A:7:DG:C8	1:A:8:DT:H73	0.54	2.38	7	4
1:A:8:DT:N3	1:A:9:DG:C6	0.54	2.76	10	4
1:A:4:TC1:N5	1:A:5:DA:N6	0.53	2.57	3	10
1:B:105:DA:C4	1:B:106:DC:C5	0.52	2.96	2	3
1:B:103:DC:C2	1:B:104:TC1:C14	0.52	2.92	4	1
1:A:2:DT:N3	1:A:3:DC:C5	0.52	2.77	5	3
1:B:104:TC1:N5	1:B:105:DA:N6	0.52	2.57	2	10
1:B:102:DT:C4	1:B:103:DC:N4	0.51	2.79	6	2
1:B:106:DC:N3	1:B:107:DG:C5	0.51	2.79	6	1
1:B:106:DC:N3	1:B:107:DG:C6	0.50	2.80	6	2
1:B:102:DT:N3	1:B:103:DC:C5	0.50	2.80	2	4
1:A:7:DG:C8	1:A:8:DT:H71	0.49	2.42	7	1
1:A:6:DC:N3	1:A:7:DG:C6	0.48	2.81	6	4
1:A:3:DC:C5	1:A:4:TC1:S12	0.48	3.06	4	1
1:A:11:DA:C6	1:A:12:DG:O6	0.48	2.66	1	1
1:A:10:DG:C6	1:A:11:DA:C6	0.48	3.01	1	7
1:B:106:DC:C2	1:B:107:DG:C5	0.47	3.03	6	1
1:A:8:DT:O4	1:B:105:DA:N1	0.47	2.48	9	7
1:A:6:DC:N3	1:A:7:DG:C5	0.47	2.83	6	2
1:A:4:TC1:C6	1:A:5:DA:N6	0.47	2.78	10	10
1:B:110:DG:C6	1:B:111:DA:C6	0.46	3.03	4	4
1:B:105:DA:C6	1:B:106:DC:N4	0.46	2.83	2	2
1:A:1:DC:C6	1:A:2:DT:H72	0.46	2.45	9	3
1:B:106:DC:N4	1:B:107:DG:O6	0.46	2.49	5	8
1:B:107:DG:C2'	1:B:108:DT:H71	0.46	2.41	10	4
1:A:6:DC:C2	1:A:7:DG:C5	0.46	3.04	6	3
1:A:5:DA:N1	1:B:108:DT:O4	0.46	2.49	8	4
1:A:5:DA:N6	1:B:107:DG:O6	0.45	2.50	1	5
1:A:6:DC:N4	1:A:7:DG:O6	0.45	2.49	4	9
1:A:10:DG:O6	1:A:11:DA:N6	0.45	2.50	1	4
1:A:7:DG:O6	1:B:105:DA:N6	0.45	2.49	2	2
1:A:2:DT:C2	1:A:3:DC:C6	0.45	3.04	5	1
1:B:104:TC1:C6	1:B:105:DA:N6	0.45	2.80	7	10
1:B:110:DG:C6	1:B:111:DA:N6	0.45	2.85	6	3
1:B:110:DG:O6	1:B:111:DA:N6	0.45	2.50	4	4
1:A:2:DT:N3	1:A:3:DC:C4	0.44	2.86	6	2
1:B:101:DC:C6	1:B:102:DT:H72	0.43	2.48	2	2
1:A:10:DG:C6	1:A:11:DA:N6	0.42	2.87	1	1

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:3:DC:C4	1:A:4:TC1:C4	0.42	3.02	4	1
1:A:5:DA:C4	1:A:6:DC:C5	0.42	3.07	1	3
1:A:7:DG:C2'	1:A:8:DT:H71	0.42	2.44	6	4
1:A:3:DC:N4	1:A:4:TC1:N5	0.41	2.67	4	1
1:A:11:DA:C6	1:A:12:DG:C6	0.41	3.07	1	1
1:B:108:DT:O4	1:B:109:DG:O6	0.41	2.39	3	2
1:B:102:DT:C2	1:B:103:DC:C6	0.40	3.07	2	1
1:B:107:DG:H2''	1:B:108:DT:H71	0.40	1.94	9	1

## 6.3 Torsion angles [i](#)

### 6.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

### 6.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

### 6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

Mol	Type	Chain	Res	Link	Bond lengths		
					Counts	RMSZ	#Z>2
1	TC1	A	4	1	24,29,30	1.29±0.01	4±0 (15±2%)
1	TC1	B	104	1	24,29,30	1.29±0.01	4±0 (15±1%)

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.

Mol	Type	Chain	Res	Link	Counts	Bond angles	
						RMSZ	#Z>2
1	TC1	A	4	1	30,42,45	2.36±0.01	8±0 (28±1%)
1	TC1	B	104	1	30,42,45	2.36±0.01	8±0 (27±1%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TC1	B	104	1	-	0±0,4,21,22	0±0,4,4,4
1	TC1	A	4	1	-	0±0,4,21,22	0±0,4,4,4

All unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
1	A	4	TC1	C4-C13	3.77	1.47	1.41	3	10
1	B	104	TC1	C4-C13	3.76	1.47	1.41	10	10
1	B	104	TC1	C4-N5	2.39	1.39	1.35	4	10
1	A	4	TC1	C4-N5	2.36	1.39	1.35	8	10
1	A	4	TC1	C6-C11	2.17	1.40	1.43	10	10
1	B	104	TC1	C6-C11	2.16	1.40	1.43	9	10
1	A	4	TC1	C6-N5	2.07	1.38	1.35	8	6
1	B	104	TC1	C6-N5	2.06	1.38	1.35	10	7

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	4	TC1	C6-N5-C4	7.70	127.67	116.85	3	10
1	B	104	TC1	C6-N5-C4	7.70	127.67	116.85	7	10
1	A	4	TC1	C4-N3-C2	4.57	119.30	114.90	2	10

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	B	104	TC1	C4-N3-C2	4.57	119.30	114.90	9	10
1	A	4	TC1	C4-C13-S12	4.54	123.94	120.87	8	10
1	B	104	TC1	C11-C6-N5	4.54	122.12	125.87	6	10
1	B	104	TC1	C4-C13-S12	4.53	123.93	120.87	4	10
1	A	4	TC1	C11-C6-N5	4.50	122.15	125.87	9	10
1	A	4	TC1	C13-C4-N5	3.31	119.45	121.33	2	10
1	B	104	TC1	C13-C4-N5	3.31	119.45	121.33	4	10
1	B	104	TC1	C7-C6-C11	2.82	119.80	116.04	6	10
1	A	4	TC1	C7-C6-C11	2.80	119.78	116.04	10	10
1	A	4	TC1	C6-C11-S12	2.21	122.36	120.87	10	10
1	B	104	TC1	C6-C11-S12	2.20	122.36	120.87	6	10
1	A	4	TC1	O4'-C4'-C5'	2.18	116.55	109.37	3	10
1	B	104	TC1	O4'-C4'-C5'	2.18	116.55	109.37	2	10
1	B	104	TC1	O4'-C1'-C2'	2.03	102.41	106.25	6	1
1	A	4	TC1	O4'-C1'-C2'	2.02	102.44	106.25	5	5

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 6.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 6.7 Other polymers [i](#)

There are no such molecules in this entry.

## 6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 7 Chemical shift validation

No chemical shift data were provided