

Full wwPDB X-ray Structure Validation Report (i)

May 16, 2020 – 01:40 am BST

PDB ID : 5TVA

Title : A. aeolicus BioW with AMP and CoA

Authors: Estrada, P.; Manandhar, M.; Dong, S.-H.; Deveryshetty, J.; Agarwal, V.;

Cronan, J.E.; Nair, S.K.

Deposited on : 2016-11-08

Resolution : 2.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

 $Mol Probity \quad : \quad 4.02b\text{--}467$

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13 EDS : 2.11

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac: 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.11

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \text{Whole archive} \\ (\#\text{Entries}) \end{array}$	$\begin{array}{c} {\rm Similar \; resolution} \\ (\#{\rm Entries, \; resolution \; range(\AA)}) \end{array}$
R_{free}	130704	1377 (2.26-2.26)
Clashscore	141614	1487 (2.26-2.26)
Ramachandran outliers	138981	1449 (2.26-2.26)
Sidechain outliers	138945	1450 (2.26-2.26)
RSRZ outliers	127900	1356 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	240	86%	12%	•
1	В	240	78%	21%	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:



Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	COA	A	302	X	-	-	-



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 4034 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called 6-carboxyhexanoate--CoA ligase.


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	A	239	Total 1922	C 1226	N 328	O 361	S 7	0	1	0
1	В	238	Total 1909	C 1217		O 358	S 7	0	0	0

• Molecule 2 is ADENOSINE MONOPHOSPHATE (three-letter code: AMP) (formula: C₁₀H₁₄N₅O₇P).

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf		
2	Λ	1	Total	С	Ν	О	Р	0	0
2	2 A	1	23	10	5	7	1	U	0
9	D	1	Total	С	N	О	Р	0	0
2	Б	1	23	10	5	7	1	U	0

• Molecule 3 is COENZYME A (three-letter code: COA) (formula: $C_{21}H_{36}N_7O_{16}P_3S$).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf			
3	A	1	Total 48	C 21	N 7	O 16	P 3	S 1	0	0

• Molecule 4 is water.


\mathbf{Mol}	Chain	Residues	${f Atoms}$	ZeroOcc	AltConf
4	A	73	Total O 73 73	0	0
4	В	36	Total O 36 36	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: 6-carboxyhexanoate--CoA ligase





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	55.66Å 68.86Å 156.57Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.07 - 2.25	Depositor
resolution (A)	39.14 - 2.25	EDS
% Data completeness	99.9 (38.07-2.25)	Depositor
(in resolution range)	$100.0 \ (39.14-2.25)$	EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	2.04 (at 2.24Å)	Xtriage
Refinement program	PHENIX (1.10_2155: ???)	Depositor
P. P.	0.201 , 0.245	Depositor
R, R_{free}	0.202 , 0.247	DCC
R_{free} test set	1467 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	45.1	Xtriage
Anisotropy	0.497	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.36, 46.2	EDS
L-test for twinning ²	$ < L > = 0.47, < L^2> = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4034	wwPDB-VP
Average B, all atoms (Å ²)	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 5.23% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $< L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: COA, AMP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Chain		Bond	lengths	Bond angles		
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.47	0/1953	0.61	1/2616 (0.0%)	
1	В	0.45	0/1937	0.58	0/2594	
All	All	0.46	0/3890	0.60	1/5210 (0.0%)	

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	${f Atoms}$	Z	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^o)$
1	A	166	LEU	CA-CB-CG	6.62	130.53	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1922	0	1987	22	0
1	В	1909	0	1972	44	0
2	A	23	0	12	1	0
2	В	23	0	12	0	0
3	A	48	0	31	3	0
4	A	73	0	0	3	0
4	В	36	0	0	4	0
All	All	4034	0	4014	66	0



The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (66) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:91:GLU:HG2	1:B:95:LYS:NZ	1.48	1.25
1:A:132:ARG:NH2	4:A:401:HOH:O	1.95	0.98
1:B:91:GLU:HG2	1:B:95:LYS:HZ2	1.19	0.97
1:B:91:GLU:CG	1:B:95:LYS:NZ	2.36	0.88
1:B:180:CYS:SG	4:B:619:HOH:O	2.33	0.86
1:A:2:ASP:N	4:A:402:HOH:O	2.17	0.78
1:B:64:LYS:NZ	4:B:603:HOH:O	2.17	0.78
1:B:2:ASP:O	1:B:25:LYS:HD2	1.89	0.72
1:B:91:GLU:CG	1:B:95:LYS:HZ2	2.00	0.70
1:B:74:GLU:O	1:B:78:GLU:HG2	1.91	0.70
1:B:91:GLU:HG2	1:B:95:LYS:HZ3	1.53	0.66
1:B:120:ILE:HG23	1:B:121:GLU:HG3	1.80	0.63
1:A:192:VAL:HG12	1:A:200:VAL:HB	1.80	0.63
1:A:206:LYS:HD2	1:A:210:ASP:HB3	1.82	0.62
1:B:124:GLU:OE1	1:B:221:ARG:NH2	2.31	0.62
1:B:108:LYS:NZ	1:B:131:GLU:OE1	2.30	0.61
1:B:91:GLU:HG2	1:B:95:LYS:HZ1	1.63	0.59
1:B:54:LYS:N	4:B:602:HOH:O	2.16	0.58
1:B:119:ASP:HB3	1:B:123:GLY:H	1.68	0.58
1:B:124:GLU:N	1:B:124:GLU:OE2	2.37	0.56
1:A:215:ARG:NH2	2:A:301:AMP:O1P	2.38	0.56
1:B:2:ASP:N	1:B:25:LYS:HZ3	2.04	0.56
1:A:124:GLU:OE2	4:A:403:HOH:O	2.19	0.54
1:A:190:GLY:HA3	1:A:205:LEU:HD12	1.90	0.53
1:B:82:LYS:O	1:B:85:THR:HG22	2.07	0.53
1:A:183:ASP:N	1:A:183:ASP:OD1	2.40	0.52
1:B:89:ILE:HG12	1:B:120:ILE:HD11	1.90	0.52
1:B:59:VAL:HG11	1:B:172:PHE:CE1	2.45	0.51
1:B:113:ARG:HH21	1:B:132:ARG:CZ	2.23	0.50
1:A:135:ARG:HH22	3:A:302:COA:P3B	2.34	0.50
1:B:178:GLU:OE2	1:B:215:ARG:HD3	2.12	0.50
1:B:143:ASP:OD2	1:B:146:LYS:HG3	2.12	0.49
1:B:141:TRP:CD2	1:B:147:VAL:HG11	2.48	0.49
1:B:108:LYS:HE2	1:B:108:LYS:HB2	1.55	0.48
1:B:27:GLU:H	1:B:27:GLU:CD	2.17	0.47
1:A:135:ARG:NH2	3:A:302:COA:O9A	2.47	0.47
1:B:83:LYS:HA	1:B:83:LYS:HD3	1.71	0.47

Continued on next page...



Continued from previous page...

A		Interatomic	Clash
Atom-1	Atom-2	${ m distance} \; ({ m \AA})$	overlap (Å)
1:A:178:GLU:OE2	1:A:215:ARG:HD3	2.14	0.47
1:B:9:ARG:NE	4:B:601:HOH:O	2.14	0.46
1:A:92:GLU:H	1:A:92:GLU:CD	2.17	0.46
1:B:91:GLU:CG	1:B:95:LYS:HZ1	2.24	0.46
1:B:15:LYS:O	1:B:17:VAL:HG13	2.16	0.46
1:B:73:PRO:HA	1:B:204:PRO:O	2.15	0.45
1:B:73:PRO:O	1:B:77:ARG:HG3	2.17	0.45
1:B:125:ARG:NH2	1:B:130:LYS:HB3	2.31	0.45
1:B:95:LYS:HE2	1:B:95:LYS:HB2	1.47	0.45
1:A:167:THR:HG22	1:A:178:GLU:OE1	2.17	0.45
1:B:91:GLU:C	1:B:95:LYS:HZ3	2.20	0.44
1:B:178:GLU:O	1:B:192:VAL:HA	2.18	0.44
1:A:41:LYS:HD2	1:A:41:LYS:N	2.32	0.44
1:A:137:ILE:HG22	3:A:302:COA:H10	1.99	0.43
1:A:180:CYS:HB2	1:A:215:ARG:HG2	2.00	0.43
1:A:12:LYS:HG2	1:A:44[A]:ASP:OD2	2.19	0.43
1:B:199:TYR:CE2	1:B:201:ARG:HG3	2.54	0.42
1:A:141:TRP:CD2	1:A:147:VAL:HG11	2.55	0.42
1:A:192:VAL:HG13	1:A:197:ILE:HD12	2.01	0.41
1:A:74:GLU:CD	1:A:74:GLU:H	2.23	0.41
1:B:81:VAL:HG11	1:B:95:LYS:HD3	2.02	0.41
1:B:20:ALA:HB3	1:B:36:LEU:HD22	2.03	0.41
1:B:143:ASP:CG	1:B:146:LYS:HG3	2.42	0.41
1:B:175:VAL:HA	1:B:219:VAL:HG12	2.03	0.40
1:B:24:VAL:HB	1:B:28:GLU:HB2	2.03	0.40
1:B:78:GLU:H	1:B:78:GLU:HG2	1.75	0.40
1:A:73:PRO:O	1:A:77:ARG:HG3	2.22	0.40
1:B:2:ASP:HB3	1:B:3:LEU:H	1.69	0.40
1:A:203:THR:HA	1:A:204:PRO:HA	1.82	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	A	$238/240 \ (99\%)$	227 (95%)	10 (4%)	1 (0%)	34	37
1	В	$236/240 \ (98\%)$	232 (98%)	4 (2%)	0	100	100
All	All	474/480 (99%)	459 (97%)	14 (3%)	1 (0%)	47	55

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	208	LYS

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	${f Analysed}$	Analysed Rotameric O		Percentiles
1	A	211/211 (100%)	203 (96%)	8 (4%)	33 39
1	В	$209/211 \; (99\%)$	206 (99%)	3 (1%)	67 76
All	All	$420/422 \ (100\%)$	409 (97%)	11 (3%)	46 55

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	ASP
1	A	96	LYS
1	A	145	LYS
1	A	153	LYS
1	A	166	LEU
1	A	167	THR
1	A	192	VAL
1	A	203	THR
1	В	71	LYS
1	В	95	LYS
1	В	215	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.



5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Tuno	Chain	Res	Link	В	ond leng	gths	В	ond ang	les
WIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	$\mid \# Z > 2 \mid$
2	AMP	В	500	-	22,25,25	1.03	2 (9%)	25,38,38	1.26	3 (12%)
3	COA	A	302	-	41,50,50	4.27	16 (39%)	52,75,75	1.54	7 (13%)
2	AMP	A	301	-	22,25,25	0.90	1 (4%)	25,38,38	1.45	4 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	${f Chirals}$	Torsions	Rings
3	COA	A	302	-	3/3/11/13	16/44/64/64	0/3/3/3
2	AMP	В	500	-	-	0/6/26/26	0/3/3/3
2	AMP	A	301	-	-	1/6/26/26	0/3/3/3

All (19) bond length outliers are listed below:



Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(ext{\AA})$	$Ideal(\AA)$
3	A	302	COA	C2B-C1B	-17.48	1.27	1.53
3	A	302	COA	O4B-C1B	9.98	1.55	1.41
3	A	302	COA	C3B-C4B	-8.24	1.30	1.52
3	A	302	COA	C5P-N4P	6.86	1.49	1.33
3	A	302	COA	C2B-C3B	6.74	1.68	1.52
3	A	302	COA	C9P-N8P	5.90	1.46	1.33
3	A	302	COA	O4B-C4B	5.33	1.56	1.45
3	A	302	COA	C6A-N6A	5.01	1.52	1.34
3	A	302	COA	P3B-O3B	4.85	1.68	1.59
3	A	302	COA	C6P-C5P	3.19	1.57	1.51
3	A	302	COA	C5B-C4B	2.95	1.60	1.51
3	A	302	COA	C3P-N4P	2.75	1.52	1.46
3	A	302	COA	O5P-C5P	-2.75	1.17	1.23
2	В	500	AMP	C5-C4	2.74	1.48	1.40
2	A	301	AMP	C5-C4	2.72	1.48	1.40
3	A	302	COA	P2A-O6A	2.49	1.69	1.59
2	В	500	AMP	C2-N3	2.40	1.36	1.32
3	A	302	COA	OAP-CAP	-2.18	1.38	1.42
3	A	302	COA	P1A-O5B	2.12	1.67	1.59

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
3	A	302	COA	N3A-C2A-N1A	-4.57	121.54	128.68
3	A	302	COA	C3B-C2B-C1B	-4.09	90.84	99.89
2	A	301	AMP	C4-C5-N7	-3.58	105.67	109.40
2	A	301	AMP	N3-C2-N1	-3.33	123.47	128.68
3	A	302	COA	O4B-C1B-C2B	-3.23	102.21	106.93
2	В	500	AMP	N3-C2-N1	-2.93	124.10	128.68
2	В	500	AMP	C4-C5-N7	-2.83	106.45	109.40
3	A	302	COA	CEP-CBP-CDP	-2.62	103.82	109.17
3	A	302	COA	C1B-N9A-C4A	2.47	130.97	126.64
3	A	302	COA	O9P-C9P-N8P	-2.39	117.86	122.99
2	A	301	AMP	O3P-P-O2P	2.31	116.48	107.64
2	В	500	AMP	O3P-P-O2P	2.31	116.47	107.64
2	A	301	AMP	O3P-P-O5'	-2.27	100.68	106.73
3	A	302	COA	O6A-CCP-CBP	2.27	114.20	110.55

All (3) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	A	302	COA	C1B
3	A	302	COA	C2B

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type	Atom
3	A	302	COA	СЗВ

All (17) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	302	COA	C3B-O3B-P3B-O7A
3	A	302	COA	C5B-O5B-P1A-O1A
3	A	302	COA	C5B-O5B-P1A-O2A
3	A	302	COA	OAP-CAP-CBP-CCP
3	A	302	COA	C9P-CAP-CBP-CCP
3	A	302	COA	OAP-CAP-CBP-CDP
3	A	302	COA	C9P-CAP-CBP-CDP
3	A	302	COA	OAP-CAP-CBP-CEP
3	A	302	COA	C9P-CAP-CBP-CEP
3	A	302	COA	CEP-CBP-CCP-O6A
2	A	301	AMP	C5'-O5'-P-O1P
3	A	302	COA	O4B-C4B-C5B-O5B
3	A	302	COA	O9P-C9P-CAP-OAP
3	A	302	COA	N8P-C9P-CAP-OAP
3	A	302	COA	C3B-O3B-P3B-O8A
3	A	302	COA	C5B-O5B-P1A-O3A
3	A	302	COA	C3B-C4B-C5B-O5B


There are no ring outliers.


2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	302	COA	3	0
2	A	301	AMP	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$\#\mathrm{RSRZ}{>}2$	$OWAB(A^2)$	Q<0.9
1	A	239/240 (99%)	0.32	8 (3%) 46 48	32, 46, 75, 86	0
1	В	238/240 (99%)	0.25	8 (3%) 45 47	35, 57, 81, 87	0
All	All	477/480 (99%)	0.28	16 (3%) 45 47	32, 52, 78, 87	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	122	THR	3.9
1	A	100	ILE	2.9
1	В	107	PRO	2.9
1	В	222	LYS	2.8
1	A	213	GLY	2.7
1	В	223	GLU	2.7
1	A	208	LYS	2.7
1	В	71	LYS	2.7
1	A	186	ASP	2.6
1	В	224	LEU	2.6
1	A	188	VAL	2.5
1	A	72	SER	2.5
1	A	209	GLY	2.5
1	A	212	LEU	2.3
1	В	124	GLU	2.3
1	В	221	ARG	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



6.3 Carbohydrates (i)


There are no carbohydrates in this entry.

6.4 Ligands (i)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	${f Res}$	Atoms	RSCC	RSR	${f B-factors}({f A}^2)$	Q<0.9
3	COA	A	302	48/48	0.93	0.15	43,53,65,68	0
2	AMP	A	301	23/23	0.93	0.14	35,52,59,70	0
2	AMP	В	500	23/23	0.95	0.14	43,56,74,75	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.











6.5 Other polymers (i)

There are no such residues in this entry.

