



# Full wwPDB X-ray Structure Validation Report i

Jul 25, 2024 – 08:02 AM EDT

PDB ID : 8TV2  
Title : Structure of apo FabS1C\_L1  
Authors : Singer, A.U.; Bruce, H.A.; Blazer, L.; Adams, J.J.; Sicheri, F.; Sidhu, S.S.  
Deposited on : 2023-08-17  
Resolution : 2.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>  
with specific help available everywhere you see the i symbol.

The types of validation reports are described at  
<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*


The reported resolution of this entry is 2.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1332 (2.68-2.64)
Clashscore	141614	1374 (2.68-2.64)
Ramachandran outliers	138981	1349 (2.68-2.64)
Sidechain outliers	138945	1349 (2.68-2.64)
RSRZ outliers	127900	1318 (2.68-2.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



## 2 Entry composition [\(i\)](#)

There are 5 unique types of molecules in this entry. The entry contains 6551 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called S1C variant of Fab\_L1 heavy chain.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	D	232	Total C	N	O	S	0	1	0
			1710	1087	278	339	6		

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	A	225	Total C	N	O	S	0	0	0
			1673	1066	274	328	5		

- Molecule 2 is a protein called S1C variant of Fab\_L1 light chain.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	E	210	Total C	N	O	S	0	0	0
			1573	984	258	325	6		

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	209	Total C	N	O	S	0	0	0
			1553	972	252	324	5		

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	D	2	Total Cl		0	0
			2	2		
3	E	3	Total Cl		0	0
			3	3		
3	A	2	Total Cl		0	0
			2	2		
3	B	1	Total Cl		0	0
			1	1		

- Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	D	1	Total Na		0	0
			1	1		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total Na 1 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	D	11	Total O 11 11	0	0
5	E	8	Total O 9 9	0	1
5	A	7	Total O 7 7	0	0
5	B	5	Total O 5 5	0	0

### 3 Residue-property plots [\(i\)](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: S1C variant of Fab\_L1 heavy chain




- Molecule 1: S1C variant of Fab\_L1 heavy chain



- Molecule 2: S1C variant of Fab\_L1 light chain



- Molecule 2: S1C variant of Fab\_L1 light chain





## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	74.26 Å    74.26 Å    370.84 Å 90.00°    90.00°    120.00°	Depositor
Resolution (Å)	64.31 – 2.65 64.31 – 2.65	Depositor EDS
% Data completeness (in resolution range)	93.9 (64.31-2.65) 93.9 (64.31-2.65)	Depositor EDS
$R_{merge}$	0.13	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle^1$	1.75 (at 2.65 Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
$R$ , $R_{free}$	0.204 , 0.247 0.198 , 0.241	Depositor DCC
$R_{free}$ test set	2002 reflections (5.94%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	67.2	Xtriage
Anisotropy	0.304	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 42.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.043 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	6551	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	78.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.24% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [\(i\)](#)

### 5.1 Standard geometry [\(i\)](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.48	0/1721	0.68	0/2357
1	D	0.54	0/1758	0.74	1/2409 (0.0%)
2	B	0.50	0/1586	0.69	0/2161
2	E	0.52	0/1606	0.70	0/2187
All	All	0.51	0/6671	0.71	1/9114 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	4

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	30	ILE	CG1-CB-CG2	-5.34	99.66	111.40

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	174	VAL	Mainchain
1	D	175[A]	THR	Mainchain
1	D	175[B]	THR	Mainchain

## 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1673	0	1576	37	0
1	D	1710	0	1610	28	0
2	B	1553	0	1449	29	0
2	E	1573	0	1485	22	0
3	A	2	0	0	1	0
3	B	1	0	0	0	0
3	D	2	0	0	0	0
3	E	3	0	0	0	0
4	B	1	0	0	0	0
4	D	1	0	0	0	0
5	A	7	0	0	0	0
5	B	5	0	0	0	0
5	D	11	0	0	1	0
5	E	9	0	0	0	0
All	All	6551	0	6120	107	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (107) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:145:VAL:HG12	1:A:233:LYS:HD2	1.57	0.85
2:E:160:ARG:HD2	2:E:181:VAL:HG11	1.62	0.82
1:D:91:MET:HB3	1:D:94:LEU:HD21	1.60	0.81
2:B:182:THR:HG22	2:B:192:SER:H	1.47	0.78
2:E:182:THR:HG22	2:E:192:SER:H	1.56	0.71
1:D:240:CYS:N	2:E:232:CYS:SG	2.65	0.70
2:E:19:VAL:HG21	2:E:94:LEU:HD13	1.78	0.66
2:B:125:LYS:HA	2:B:158:TYR:OH	1.95	0.66
1:A:183:LEU:HD21	1:A:206:VAL:HG21	1.77	0.66
1:A:190:PHE:HE1	1:A:205:VAL:CG2	2.09	0.66
2:B:207:HIS:O	2:B:229:ARG:NH1	2.31	0.63
1:A:143:PRO:HB3	1:A:169:TYR:HB3	1.80	0.62
2:B:198:THR:O	2:B:199:LEU:HD23	1.99	0.62

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:168:ASP:HB3	1:A:199:LEU:HD13	1.82	0.61
2:B:105:GLN:HE21	2:B:111:PHE:HB3	1.65	0.60
2:B:29:VAL:HG11	2:B:106:GLN:HE21	1.65	0.60
2:E:126:ARG:HH21	2:E:127:THR:HB	1.67	0.59
2:B:177:SER:HA	2:B:196:THR:O	2.03	0.58
1:D:113:TRP:CE3	1:A:118:VAL:HG12	2.37	0.58
1:A:106:ARG:NH2	1:A:125:ASP:OD2	2.31	0.58
1:D:158:GLY:O	1:D:210:SER:HB2	2.04	0.57
1:D:195:GLN:NE2	5:D:401:HOH:O	2.36	0.56
2:E:124:ILE:HG23	2:E:184:GLN:NE2	2.20	0.56
2:B:181:VAL:HG22	2:B:193:LEU:HD12	1.87	0.56
2:E:184:GLN:HG3	2:E:191:TYR:CE2	2.40	0.56
2:B:21:ILE:HG12	2:B:120:THR:HG21	1.88	0.55
2:B:158:TYR:CD1	2:B:159:PRO:HA	2.41	0.55
1:D:13:VAL:HG11	1:D:94:LEU:HD12	1.88	0.55
1:A:195:GLN:HA	2:B:178:GLN:HE22	1.72	0.53
1:D:175[A]:THR:HG23	1:D:223:ASN:HB3	1.89	0.53
2:E:185:ASP:HB3	2:E:188:ASP:OD1	2.09	0.53
1:A:146:PHE:CE2	2:B:142:GLN:HG3	2.42	0.53
1:D:146:PHE:CE2	2:E:142:GLN:HG3	2.44	0.52
2:E:182:THR:CG2	2:E:192:SER:H	2.23	0.52
2:E:29:VAL:O	2:E:30:SER:OG	2.25	0.52
1:A:14:GLN:H	1:A:14:GLN:CD	2.12	0.52
1:A:30:ILE:HD11	1:A:87:ALA:HB2	1.92	0.52
1:A:71:VAL:HG13	1:A:76:PHE:HB2	1.92	0.52
1:D:7:SER:OG	1:D:22:SER:HB2	2.10	0.51
1:D:111:VAL:HG23	1:D:111:VAL:O	2.10	0.51
1:A:60:TYR:CE2	1:A:111:VAL:HG22	2.46	0.51
1:A:106:ARG:HH21	1:A:125:ASP:CG	2.14	0.51
2:B:6:GLN:NE2	2:B:119:GLY:H	2.09	0.51
1:A:41:TRP:HD1	1:A:78:ILE:HD12	1.77	0.50
2:B:150:VAL:HG13	2:B:197:LEU:HB3	1.94	0.50
2:E:184:GLN:HG3	2:E:191:TYR:CZ	2.48	0.49
2:B:200:SER:OG	2:B:203:ASP:OD1	2.30	0.49
2:B:182:THR:CG2	2:B:192:SER:H	2.21	0.48
1:D:30:ILE:CD1	1:D:80:ALA:HB1	2.43	0.48
1:A:113:TRP:HB3	1:A:115:TRP:CE2	2.49	0.48
1:D:76:PHE:CZ	1:D:91:MET:HG2	2.50	0.47
1:A:32:SER:HB2	1:A:112:TRP:CZ3	2.49	0.47
2:B:176:ASN:N	2:B:176:ASN:OD1	2.47	0.47
1:D:192:ALA:HA	1:D:202:LEU:HB3	1.96	0.47

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:30:ILE:HD11	1:D:80:ALA:HB1	1.97	0.47
2:E:12:SER:HA	2:E:123:GLU:O	2.15	0.47
1:D:41:TRP:HD1	1:D:78:ILE:HD12	1.80	0.46
2:E:53:LEU:HA	2:E:71:VAL:HG21	1.95	0.46
1:A:190:PHE:HE1	1:A:205:VAL:HG23	1.81	0.46
2:E:126:ARG:HD2	2:E:188:ASP:O	2.16	0.46
1:D:9:GLY:HA3	1:D:131:THR:OG1	2.17	0.45
1:D:41:TRP:NE1	1:D:89:LEU:HB2	2.32	0.45
1:A:14:GLN:H	1:A:14:GLN:NE2	2.14	0.45
2:E:151:VAL:HG22	2:E:196:THR:HG23	1.99	0.45
1:A:213:LEU:HD23	1:A:213:LEU:HA	1.74	0.45
1:D:67:TYR:CE1	1:D:78:ILE:HG22	2.52	0.45
1:D:76:PHE:CE2	1:D:91:MET:HG2	2.52	0.45
2:E:30:SER:HB3	1:A:79:SER:HB2	1.99	0.45
1:A:189:THR:HA	1:A:204:SER:HA	1.99	0.45
1:A:205:VAL:HG11	2:B:153:LEU:HD22	1.99	0.45
1:A:208:VAL:HG21	1:A:218:TYR:CZ	2.52	0.45
2:B:136:PHE:O	2:B:150:VAL:HG23	2.17	0.45
2:E:181:VAL:HG22	2:E:193:LEU:HD12	1.98	0.44
1:A:13:VAL:HG11	1:A:94:LEU:HD13	2.00	0.43
1:D:178:TRP:CH2	1:D:220:CYS:HB3	2.53	0.43
1:D:213:LEU:HA	1:D:213:LEU:HD23	1.73	0.43
1:A:205:VAL:HG21	2:B:153:LEU:HD13	2.01	0.43
1:D:150:PRO:HB3	1:D:161:ALA:O	2.19	0.42
1:A:150:PRO:HB2	1:A:213:LEU:HD11	2.01	0.42
1:A:208:VAL:HG21	1:A:218:TYR:OH	2.18	0.42
1:A:67:TYR:CZ	1:A:78:ILE:HG22	2.54	0.42
1:A:99:THR:HG23	1:A:134:THR:HA	2.00	0.42
2:B:19:VAL:HG21	2:B:94:LEU:HD22	2.01	0.42
2:B:167:TYR:HB2	2:B:211:ALA:HB3	2.01	0.42
1:D:13:VAL:HG11	1:D:94:LEU:CD1	2.49	0.42
2:B:163:LYS:HB3	2:B:215:THR:HB	2.02	0.42
2:E:42:TYR:O	2:E:102:TYR:HA	2.19	0.42
1:A:195:GLN:HA	2:B:178:GLN:NE2	2.35	0.42
1:A:106:ARG:NH1	3:A:302:CL:CL	2.90	0.42
2:B:160:ARG:HB2	2:B:191:TYR:CE2	2.55	0.41
1:D:216:GLN:HA	1:D:216:GLN:OE1	2.20	0.41
1:D:187:VAL:HG22	1:D:206:VAL:HB	2.02	0.41
1:A:199:LEU:HD23	1:A:199:LEU:HA	1.88	0.41
2:B:14:SER:OG	2:B:125:LYS:HD2	2.20	0.41
2:E:2:ILE:CD1	2:E:27:GLN:H	2.33	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:158:TYR:CD1	2:B:159:PRO:CA	3.04	0.41
2:B:167:TYR:CZ	2:B:172:LEU:HD13	2.56	0.41
2:E:12:SER:HB3	2:E:125:LYS:HG3	2.01	0.41
1:A:7:SER:OG	1:A:22:SER:HB2	2.21	0.41
1:D:212:SER:HA	1:D:215:THR:OG1	2.20	0.41
2:E:158:TYR:CG	2:E:159:PRO:HA	2.56	0.40
1:A:197:SER:OG	1:A:199:LEU:HB2	2.21	0.40
1:D:30:ILE:HG23	1:D:85:ASN:OD1	2.22	0.40
1:A:190:PHE:HE1	1:A:205:VAL:HG21	1.85	0.40
1:D:67:TYR:CZ	1:D:78:ILE:HG22	2.57	0.40
1:A:170:PHE:HB2	1:A:199:LEU:HD23	2.03	0.40
2:B:118:GLN:HA	2:B:118:GLN:OE1	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [\(i\)](#)

### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	221/237 (93%)	213 (96%)	8 (4%)	0	100 100
1	D	231/237 (98%)	222 (96%)	9 (4%)	0	100 100
2	B	207/211 (98%)	196 (95%)	11 (5%)	0	100 100
2	E	208/211 (99%)	200 (96%)	8 (4%)	0	100 100
All	All	867/896 (97%)	831 (96%)	36 (4%)	0	100 100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	179/198 (90%)	166 (93%)	13 (7%)	14   21
1	D	184/198 (93%)	170 (92%)	14 (8%)	13   21
2	B	169/184 (92%)	157 (93%)	12 (7%)	14   22
2	E	174/184 (95%)	166 (95%)	8 (5%)	27   41
All	All	706/764 (92%)	659 (93%)	47 (7%)	17   25

All (47) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	26	SER
1	D	69	ASP
1	D	82	THR
1	D	83	SER
1	D	110	SER
1	D	136	SER
1	D	137	SER
1	D	173	PRO
1	D	175[A]	THR
1	D	175[B]	THR
1	D	203	SER
1	D	206	VAL
1	D	207	THR
1	D	210	SER
2	E	7	SER
2	E	10	SER
2	E	74	SER
2	E	83	SER
2	E	108	SER
2	E	174	SER
2	E	189	SER
2	E	232	CYS
1	A	14	GLN
1	A	26	SER
1	A	34	TYR
1	A	71	VAL
1	A	82	THR
1	A	137	SER

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	175	THR
1	A	189	THR
1	A	205	VAL
1	A	206	VAL
1	A	211	SER
1	A	221	ASN
1	A	223	ASN
2	B	7	SER
2	B	69	SER
2	B	79	SER
2	B	83	SER
2	B	85	THR
2	B	93	SER
2	B	106	GLN
2	B	126	ARG
2	B	158	TYR
2	B	189	SER
2	B	198	THR
2	B	225	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	6	GLN
2	B	106	GLN

### 5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [\(i\)](#)

Of 10 ligands modelled in this entry, 10 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data i

### 6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	225/237 (94%)	0.02	2 (0%)	84	83	56, 83, 105, 117
1	D	232/237 (97%)	-0.12	2 (0%)	84	83	45, 70, 97, 125
2	B	209/211 (99%)	-0.05	1 (0%)	91	91	60, 90, 113, 133
2	E	210/211 (99%)	-0.07	1 (0%)	91	91	46, 71, 105, 117
All	All	876/896 (97%)	-0.05	6 (0%)	87	87	45, 79, 108, 133

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	239	SER	3.2
1	D	240	CYS	3.1
1	A	120	TYR	2.9
2	E	2	ILE	2.7
1	A	150	PRO	2.3
2	B	4	MET	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains i

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates i

There are no monosaccharides in this entry.


### 6.4 Ligands i


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,


median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	CL	E	303	1/1	0.68	0.14	92,92,92,92	0
3	CL	E	302	1/1	0.74	0.19	80,80,80,80	0
3	CL	A	302	1/1	0.83	0.09	85,85,85,85	0
4	NA	B	302	1/1	0.83	0.17	68,68,68,68	0
3	CL	D	301	1/1	0.84	0.12	82,82,82,82	0
3	CL	E	301	1/1	0.85	0.15	79,79,79,79	0
3	CL	D	302	1/1	0.88	0.16	86,86,86,86	0
3	CL	A	301	1/1	0.89	0.11	68,68,68,68	0
3	CL	B	301	1/1	0.91	0.16	90,90,90,90	0
4	NA	D	303	1/1	0.92	0.13	48,48,48,48	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.


























## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.