



wwPDB EM Validation Summary Report ⓘ

Nov 2, 2022 – 09:46 AM EDT

PDB ID : 5TWV
EMDB ID : EMD-8470
Title : Cryo-EM structure of the pancreatic ATP-sensitive K⁺ channel SUR1/Kir6.2
in the presence of ATP and glibenclamide
Authors : Martin, G.M.; Yoshioka, C.; Chen, J.Z.; Shyng, S.L.
Deposited on : 2016-11-14
Resolution : 6.30 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev43
Mogul : 1.8.5 (274361), CSD as541be (2020)
MolProbity : 4.02b-467
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : 1.9.9
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.31.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 6.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	390	
1	C	390	
1	E	390	
1	G	390	
2	B	1590	
2	D	1590	
2	F	1590	
2	H	1590	

2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 41080 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ATP-sensitive inward rectifier potassium channel 11.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	325	2479	1598	422	444	15	0	0
1	C	325	2479	1598	422	444	15	0	0
1	E	325	2479	1598	422	444	15	0	0
1	G	325	2479	1598	422	444	15	0	0

- Molecule 2 is a protein called ATP-binding cassette sub-family C member 8.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	B	1375	7760	4866	1450	1439	5	0	0
2	D	1375	7760	4866	1450	1439	5	0	0
2	F	1375	7760	4866	1450	1439	5	0	0
2	H	1375	7760	4866	1450	1439	5	0	0

There are 32 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	-7	ASP	-	expression tag	UNP Q09427
B	-6	TYR	-	expression tag	UNP Q09427
B	-5	LYS	-	expression tag	UNP Q09427
B	-4	ASP	-	expression tag	UNP Q09427
B	-3	ASP	-	expression tag	UNP Q09427
B	-2	ASP	-	expression tag	UNP Q09427
B	-1	ASP	-	expression tag	UNP Q09427
B	0	LYS	-	expression tag	UNP Q09427
D	-7	ASP	-	expression tag	UNP Q09427

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	-6	TYR	-	expression tag	UNP Q09427
D	-5	LYS	-	expression tag	UNP Q09427
D	-4	ASP	-	expression tag	UNP Q09427
D	-3	ASP	-	expression tag	UNP Q09427
D	-2	ASP	-	expression tag	UNP Q09427
D	-1	ASP	-	expression tag	UNP Q09427
D	0	LYS	-	expression tag	UNP Q09427
F	-7	ASP	-	expression tag	UNP Q09427
F	-6	TYR	-	expression tag	UNP Q09427
F	-5	LYS	-	expression tag	UNP Q09427
F	-4	ASP	-	expression tag	UNP Q09427
F	-3	ASP	-	expression tag	UNP Q09427
F	-2	ASP	-	expression tag	UNP Q09427
F	-1	ASP	-	expression tag	UNP Q09427
F	0	LYS	-	expression tag	UNP Q09427
H	-7	ASP	-	expression tag	UNP Q09427
H	-6	TYR	-	expression tag	UNP Q09427
H	-5	LYS	-	expression tag	UNP Q09427
H	-4	ASP	-	expression tag	UNP Q09427
H	-3	ASP	-	expression tag	UNP Q09427
H	-2	ASP	-	expression tag	UNP Q09427
H	-1	ASP	-	expression tag	UNP Q09427
H	0	LYS	-	expression tag	UNP Q09427

- Molecule 3 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: $C_{10}H_{16}N_5O_{13}P_3$).




Mol	Chain	Residues	Atoms					AltConf
3	A	1	Total 31	C 10	N 5	O 13	P 3	0
3	C	1	Total 31	C 10	N 5	O 13	P 3	0
3	E	1	Total 31	C 10	N 5	O 13	P 3	0
3	G	1	Total 31	C 10	N 5	O 13	P 3	0


3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: ATP-sensitive inward rectifier potassium channel 11




- Molecule 1: ATP-sensitive inward rectifier potassium channel 11







• Molecule 1: ATP-sensitive inward rectifier potassium channel 11




• Molecule 1: ATP-sensitive inward rectifier potassium channel 11





• Molecule 2: ATP-binding cassette sub-family C member 8




ASP	TYR	L253	P254	I255	M256	V257	V258	F259	F260	P261	P262	L263	L264	L265	L266	L267	L268	L269	L270	L271	L272	L273	L274	L275	L276	L277	L278	L279	S280	R281	L301	L302	S303	S304	F305	M306	L315	L316	L317	L318	L319	L320	L321	L322	L323	L324	L325	L326	L327	L328	L329	L330	L331	L332	L333	L334	L335	L336	L337	L338	L339	L340	L341	L342	L343	L344	L345	L346	L347	L348	L349	L350	L351	L352	L353	L354	L355	L356	L357	L358	L359	L360	L361	L362	L363	L364	L365	L366	L367	L368	L369	L370	L371	L372	L373	L374	L375	L376	L377	L378	L379	L380	L381	L382	L383	L384	L385	L386	L387	L388	L389	L390	L391	L392	L393	L394	L395	L396	L397	L398	L399	E400	G401	G402	I403	I404	I405	D406	G407	L408	L409	L410	L411	K412	R421	L422	S423	I424	I425	L426	Q427	D428	F429
-----	-----	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------






● Molecule 2: ATP-binding cassette sub-family C member 8








● Molecule 2: ATP-binding cassette sub-family C member 8





• Molecule 2: ATP-binding cassette sub-family C member 8





4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C4	Depositor
Number of particles used	20000	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	40	Depositor
Minimum defocus (nm)	1500	Depositor
Maximum defocus (nm)	3000	Depositor
Magnification	81000	Depositor
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	0.095	Depositor
Minimum map value	-0.038	Depositor
Average map value	0.003	Depositor
Map value standard deviation	0.010	Depositor
Recommended contour level	0.027736	Depositor
Map size (Å)	233.92, 233.92, 233.92	wwPDB
Map dimensions	272, 272, 272	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.86, 0.86, 0.86	Depositor

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: ATP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.32	0/2536	0.66	1/3459 (0.0%)
1	C	0.32	0/2536	0.66	1/3459 (0.0%)
1	E	0.32	0/2536	0.66	1/3459 (0.0%)
1	G	0.32	0/2536	0.66	1/3459 (0.0%)
2	B	0.29	0/7912	0.51	0/10995
2	D	0.29	0/7912	0.51	0/10995
2	F	0.29	0/7912	0.51	0/10995
2	H	0.29	0/7912	0.51	0/10995
All	All	0.30	0/41792	0.55	4/57816 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	1
2	D	0	1
2	F	0	1
2	H	0	1
All	All	0	4

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	246	GLY	N-CA-C	5.18	126.05	113.10
1	C	246	GLY	N-CA-C	5.18	126.05	113.10
1	E	246	GLY	N-CA-C	5.18	126.05	113.10
1	G	246	GLY	N-CA-C	5.18	126.05	113.10

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	1384	GLY	Peptide
2	D	1384	GLY	Peptide
2	F	1384	GLY	Peptide
2	H	1384	GLY	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2479	0	2442	89	0
1	C	2479	0	2442	91	0
1	E	2479	0	2442	92	0
1	G	2479	0	2442	94	0
2	B	7760	0	4559	115	0
2	D	7760	0	4559	115	0
2	F	7760	0	4559	114	0
2	H	7760	0	4559	119	0
3	A	31	0	12	7	0
3	C	31	0	12	7	0
3	E	31	0	12	7	0
3	G	31	0	12	7	0
All	All	41080	0	28052	784	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 784 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:F:206:PRO:HB2	2:F:207:PRO:HD3	1.64	0.80
2:H:206:PRO:HB2	2:H:207:PRO:HD3	1.64	0.80
2:B:206:PRO:HB2	2:B:207:PRO:HD3	1.64	0.78
2:D:206:PRO:HB2	2:D:207:PRO:HD3	1.64	0.78
2:F:233:MET:O	2:F:237:ILE:CB	2.37	0.73

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	323/390 (83%)	310 (96%)	12 (4%)	1 (0%)	41	76
1	C	323/390 (83%)	310 (96%)	12 (4%)	1 (0%)	41	76
1	E	323/390 (83%)	310 (96%)	12 (4%)	1 (0%)	41	76
1	G	323/390 (83%)	310 (96%)	12 (4%)	1 (0%)	41	76
2	B	1363/1590 (86%)	1303 (96%)	56 (4%)	4 (0%)	41	76
2	D	1363/1590 (86%)	1302 (96%)	57 (4%)	4 (0%)	41	76
2	F	1363/1590 (86%)	1303 (96%)	56 (4%)	4 (0%)	41	76
2	H	1363/1590 (86%)	1303 (96%)	56 (4%)	4 (0%)	41	76
All	All	6744/7920 (85%)	6451 (96%)	273 (4%)	20 (0%)	44	76

5 of 20 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	98	VAL
2	B	277	ASP
2	B	316	GLY
2	D	98	VAL
2	D	277	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	264/341 (77%)	262 (99%)	2 (1%)	81	89
1	C	264/341 (77%)	262 (99%)	2 (1%)	81	89
1	E	264/341 (77%)	262 (99%)	2 (1%)	81	89
1	G	264/341 (77%)	262 (99%)	2 (1%)	81	89
2	B	224/1381 (16%)	222 (99%)	2 (1%)	78	87
2	D	224/1381 (16%)	222 (99%)	2 (1%)	78	87
2	F	224/1381 (16%)	222 (99%)	2 (1%)	78	87
2	H	224/1381 (16%)	222 (99%)	2 (1%)	78	87
All	All	1952/6888 (28%)	1936 (99%)	16 (1%)	82	89

5 of 16 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	H	6	CYS
1	G	133	PHE
1	E	95	PHE
1	G	95	PHE
2	D	123	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	ATP	G	401	-	26,33,33	0.92	0	31,52,52	1.56	5 (16%)
3	ATP	A	401	-	26,33,33	0.92	0	31,52,52	1.56	5 (16%)
3	ATP	C	401	-	26,33,33	0.92	0	31,52,52	1.56	5 (16%)
3	ATP	E	401	-	26,33,33	0.91	0	31,52,52	1.56	5 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ATP	G	401	-	-	6/18/38/38	0/3/3/3
3	ATP	A	401	-	-	6/18/38/38	0/3/3/3
3	ATP	C	401	-	-	6/18/38/38	0/3/3/3
3	ATP	E	401	-	-	6/18/38/38	0/3/3/3

There are no bond length outliers.

The worst 5 of 20 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	401	ATP	PB-O3B-PG	-3.84	119.65	132.83
3	C	401	ATP	PB-O3B-PG	-3.84	119.65	132.83
3	E	401	ATP	PB-O3B-PG	-3.84	119.65	132.83
3	G	401	ATP	PB-O3B-PG	-3.84	119.67	132.83
3	E	401	ATP	C3'-C2'-C1'	3.26	105.89	100.98

There are no chirality outliers.

5 of 24 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	401	ATP	C5'-O5'-PA-O2A
3	C	401	ATP	C5'-O5'-PA-O2A

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	E	401	ATP	C5'-O5'-PA-O2A
3	G	401	ATP	C5'-O5'-PA-O2A
3	A	401	ATP	O4'-C4'-C5'-O5'


There are no ring outliers.


4 monomers are involved in 28 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	G	401	ATP	7	0
3	A	401	ATP	7	0
3	C	401	ATP	7	0
3	E	401	ATP	7	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-8470. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 136



Y Index: 136




Z Index: 136


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 118



Y Index: 118




Z Index: 112

The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal surface views [i](#)


6.4.1 Primary map



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.027736. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.5 Mask visualisation

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 428 nm^3 ; this corresponds to an approximate mass of 387 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [i](#)




*Reported resolution corresponds to spatial frequency of 0.159 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.159 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	6.30	-	-
Author-provided FSC curve	6.64	7.54	6.72
Unmasked-calculated*	-	-	-

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-8470 and PDB model 5TWV. Per-residue inclusion information can be found in section 3 on page 6.

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.027736 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [\(i\)](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [\(i\)](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.027736).



















9.4 Atom inclusion [i](#)



At the recommended contour level, 81% of all backbone atoms, 67% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.027736) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.6651	 0.2310
A	 0.6482	 0.2570
B	 0.6711	 0.2220
C	 0.6478	 0.2550
D	 0.6692	 0.2230
E	 0.6486	 0.2560
F	 0.6731	 0.2230
G	 0.6494	 0.2570
H	 0.6681	 0.2230

