



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 21, 2024 – 05:20 PM EST

PDB ID : 8TWO  
Title : AvrB bound with UDP and RIN4\_T166-Rha  
Authors : Peng, W.; Orth, K.  
Deposited on : 2023-08-21  
Resolution : 2.09 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.09 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	323	
2	B	30	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	RAM	A	402	-	-	X	-

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4974 atoms, of which 2424 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Avirulence protein B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	294	4544	1441	2228	419	449	7	0	0	0


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	ALA	-	expression tag	UNP P13835
A	0	HIS	-	expression tag	UNP P13835

- Molecule 2 is a protein called RPM1-interacting protein 4.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	H	N	O			
2	B	25	373	126	174	33	40	0	0	0

- Molecule 3 is URIDINE-5'-DIPHOSPHATE (three-letter code: UDP) (formula: C<sub>9</sub>H<sub>14</sub>N<sub>2</sub>O<sub>12</sub>P<sub>2</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	H	N	O			P
3	A	1	36	9	11	2	12	2	0	0

- Molecule 4 is alpha-L-rhamnopyranose (three-letter code: RAM) (formula: C<sub>6</sub>H<sub>12</sub>O<sub>5</sub>) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
4	A	1	21	6	11	4	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Avirulence protein B



- Molecule 2: RPM1-interacting protein 4



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	57.99Å 119.24Å 39.98Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	52.15 – 2.09 52.15 – 2.09	Depositor EDS
% Data completeness (in resolution range)	97.7 (52.15-2.09) 97.7 (52.15-2.09)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	0.99 (at 2.10Å)	Xtrriage
Refinement program	PHENIX (1.20.1_4487: ???)	Depositor
R, $R_{free}$	0.221 , 0.290 0.219 , 0.287	Depositor DCC
$R_{free}$ test set	1661 reflections (10.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	49.7	Xtrriage
Anisotropy	0.273	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.43 , 64.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	4974	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	83.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.06% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: RAM, UDP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.81	3/2366 (0.1%)	0.89	3/3213 (0.1%)
2	B	0.77	0/206	0.71	0/280
All	All	0.81	3/2572 (0.1%)	0.88	3/3493 (0.1%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	79	GLU	CG-CD	6.25	1.61	1.51
1	A	98	TRP	CB-CG	-6.00	1.39	1.50
1	A	79	GLU	CD-OE2	5.27	1.31	1.25

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	117	ARG	NE-CZ-NH1	5.56	123.08	120.30
1	A	314	SER	CA-CB-OG	-5.43	96.53	111.20
1	A	302	LEU	CB-CG-CD1	-5.12	102.29	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2316	2228	2226	82	5
2	B	199	174	175	14	1
3	A	25	11	11	4	0
4	A	10	11	10	8	0
All	All	2550	2424	2422	88	5

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

All (88) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:402:RAM:C1	2:B:166:THR:OG1	1.92	1.17
4:A:402:RAM:C1	2:B:166:THR:HG1	1.94	0.78
1:A:107:GLU:O	1:A:111:SER:OG	2.05	0.74
1:A:44:LEU:HD13	1:A:49:ARG:NH1	2.04	0.73
1:A:54:SER:O	1:A:58:ASN:ND2	2.23	0.71
1:A:146:ASP:OD1	1:A:160:LYS:HD3	1.90	0.71
3:A:401:UDP:O3B	4:A:402:RAM:O2	2.13	0.65
1:A:120:ASP:OD2	1:A:121:THR:HG23	1.97	0.64
1:A:88:PHE:N	1:A:231:ASN:OD1	2.31	0.64
2:B:155:ASP:HA	2:B:158:ASN:OD1	1.97	0.63
1:A:131:TYR:OH	4:A:402:RAM:C6	2.46	0.62
1:A:242:LEU:HD11	1:A:250:LEU:HD12	1.81	0.60
1:A:131:TYR:OH	4:A:402:RAM:H63	2.00	0.60
1:A:122:ASP:OD2	2:B:171:LYS:HG2	2.01	0.60
1:A:208:GLN:HA	1:A:208:GLN:OE1	2.02	0.59
1:A:277:ARG:NH1	1:A:287:LEU:HB2	2.18	0.58
1:A:285:ILE:HD12	1:A:285:ILE:N	2.19	0.58
1:A:151:ASP:N	1:A:155:GLY:O	2.21	0.57
1:A:37:LEU:H	1:A:37:LEU:HD22	1.69	0.57
1:A:207:LEU:HD12	2:B:160:SER:O	2.05	0.56
1:A:58:ASN:O	1:A:63:ARG:NH1	2.38	0.56
1:A:131:TYR:CZ	4:A:402:RAM:H61	2.40	0.56
1:A:204:LYS:HE2	2:B:160:SER:HA	1.88	0.55
1:A:139:VAL:HG22	1:A:160:LYS:HG3	1.88	0.55
1:A:56:LYS:O	1:A:59:THR:HG23	2.07	0.55
1:A:75:SER:O	1:A:79:GLU:HG2	2.08	0.54
1:A:183:ASP:O	1:A:183:ASP:OD2	2.26	0.53
1:A:193:HIS:HD1	1:A:193:HIS:C	2.11	0.53
1:A:35:ARG:NE	1:A:35:ARG:HA	2.24	0.53
1:A:214:ARG:HH11	1:A:214:ARG:HG2	1.75	0.52

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:239:SER:O	1:A:283:HIS:HE1	1.92	0.52
1:A:208:GLN:OE1	2:B:167:HIS:NE2	2.42	0.52
1:A:121:THR:HB	2:B:168:ILE:HG22	1.92	0.51
1:A:187:LEU:O	1:A:190:ARG:N	2.45	0.51
1:A:198:LEU:HB3	1:A:202:ALA:HB3	1.92	0.50
1:A:267:GLY:HA2	3:A:401:UDP:O2A	2.11	0.50
1:A:131:TYR:CE1	4:A:402:RAM:H61	2.46	0.50
1:A:223:LEU:N	1:A:224:PRO:CD	2.76	0.48
1:A:179:ARG:O	1:A:213:ASP:HA	2.14	0.48
1:A:242:LEU:CD1	1:A:250:LEU:HD12	2.43	0.48
1:A:150:GLN:HG2	1:A:156:ASP:OD1	2.14	0.48
1:A:119:SER:HA	1:A:220:ALA:HB3	1.95	0.48
1:A:253:ARG:HG2	1:A:305:GLU:OE1	2.14	0.48
1:A:191:VAL:HG13	1:A:192:ALA:N	2.29	0.47
1:A:201:LYS:O	1:A:205:GLN:HG3	2.13	0.47
1:A:267:GLY:HA2	3:A:401:UDP:PA	2.55	0.47
1:A:179:ARG:NH1	1:A:216:ASP:OD1	2.48	0.47
1:A:196:THR:OG1	1:A:197:GLU:OE1	2.25	0.47
1:A:198:LEU:HD12	1:A:198:LEU:N	2.30	0.47
1:A:39:VAL:HA	1:A:42:GLN:HG3	1.96	0.46
1:A:208:GLN:OE1	2:B:167:HIS:CD2	2.68	0.46
2:B:156:GLU:O	2:B:158:ASN:ND2	2.48	0.46
1:A:186:TYR:CD1	1:A:186:TYR:C	2.89	0.46
1:A:308:PHE:HA	1:A:311:LYS:HG2	1.97	0.46
1:A:204:LYS:HG3	2:B:160:SER:HA	1.97	0.46
1:A:181:ALA:HB2	1:A:212:PRO:HB3	1.98	0.46
1:A:204:LYS:HE2	2:B:159:PRO:O	2.15	0.46
1:A:108:PRO:O	1:A:109:LEU:HB2	2.16	0.45
1:A:193:HIS:C	1:A:193:HIS:ND1	2.69	0.45
1:A:108:PRO:O	1:A:109:LEU:CB	2.65	0.45
1:A:247:LEU:HD21	1:A:283:HIS:CD2	2.51	0.45
1:A:33:SER:O	1:A:74:ARG:NH2	2.41	0.44
1:A:151:ASP:OD2	1:A:154:PHE:HD1	2.00	0.44
1:A:46:GLY:O	3:A:401:UDP:H1'	2.17	0.44
1:A:184:THR:OG1	2:B:149:PRO:HD2	2.17	0.44
1:A:200:ALA:O	1:A:201:LYS:C	2.56	0.44
1:A:151:ASP:OD1	1:A:152:ASN:N	2.51	0.44
1:A:188:ARG:HG2	1:A:188:ARG:HH11	1.83	0.44
1:A:203:LEU:O	1:A:207:LEU:HG	2.19	0.43
1:A:291:ARG:HG3	1:A:315:LEU:O	2.19	0.43
1:A:204:LYS:O	1:A:205:GLN:C	2.56	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:150:LYS:HB2	2:B:153:ASP:OD2	2.20	0.42
1:A:242:LEU:HD11	1:A:250:LEU:CD1	2.47	0.42
1:A:29:LEU:HD23	1:A:81:SER:HA	2.02	0.42
1:A:45:ILE:HD13	1:A:277:ARG:CZ	2.49	0.42
1:A:283:HIS:HB2	1:A:285:ILE:HD13	2.01	0.42
1:A:197:GLU:HB2	1:A:198:LEU:HD12	2.00	0.42
1:A:131:TYR:CZ	4:A:402:RAM:C6	3.03	0.42
1:A:137:ARG:O	1:A:141:HIS:HB2	2.20	0.42
1:A:92:TRP:CH2	1:A:271:LYS:HB2	2.55	0.42
1:A:202:ALA:HA	1:A:205:GLN:HG3	2.01	0.42
1:A:45:ILE:CD1	1:A:277:ARG:CZ	2.97	0.41
1:A:94:ARG:HD3	1:A:94:ARG:HA	1.90	0.41
1:A:37:LEU:H	1:A:37:LEU:CD2	2.33	0.41
1:A:107:GLU:H	1:A:107:GLU:CD	2.24	0.41
1:A:232:ASP:O	1:A:236:GLN:HG3	2.21	0.40
1:A:211:ASN:O	1:A:214:ARG:NH1	2.50	0.40
1:A:306:GLU:O	1:A:307:GLU:C	2.59	0.40

All (5) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:228:ASP:OD2	1:A:292:ASN:ND2[1_556]	1.81	0.39
1:A:171:ASN:OD1	2:B:165:TYR:OH[4_456]	2.07	0.13
1:A:79:GLU:OE2	1:A:79:GLU:OE2[2_655]	2.08	0.12
1:A:27:ARG:NH2	1:A:152:ASN:OD1[3_546]	2.14	0.06
1:A:228:ASP:OD2	1:A:292:ASN:HD21[1_556]	1.56	0.04

## 5.3 Torsion angles [\(i\)](#)

### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	292/323 (90%)	264 (90%)	26 (9%)	2 (1%)	22	18
2	B	23/30 (77%)	19 (83%)	3 (13%)	1 (4%)	2	0
All	All	315/353 (89%)	283 (90%)	29 (9%)	3 (1%)	15	11

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	53	ASP
1	A	109	LEU
2	B	150	LYS

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	244/274 (89%)	233 (96%)	11 (4%)	27	27
2	B	21/26 (81%)	21 (100%)	0	100	100
All	All	265/300 (88%)	254 (96%)	11 (4%)	30	30

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	56	LYS
1	A	78	ASP
1	A	81	SER
1	A	94	ARG
1	A	111	SER
1	A	186	TYR
1	A	193	HIS
1	A	235	ARG
1	A	238	ILE
1	A	253	ARG
1	A	274	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are

no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	RAM	A	402	-	10,10,11	2.05	3 (30%)	14,14,16	3.91	7 (50%)
3	UDP	A	401	-	24,26,26	0.93	1 (4%)	37,40,40	0.70	1 (2%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	RAM	A	402	-	-	-	0/1/1/1
3	UDP	A	401	-	-	7/16/32/32	0/2/2/2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	402	RAM	O3-C3	4.32	1.53	1.43
4	A	402	RAM	O5-C1	3.38	1.49	1.43
3	A	401	UDP	PA-O1A	2.89	1.61	1.50
4	A	402	RAM	C2-C3	-2.04	1.49	1.52

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	402	RAM	C3-C4-C5	9.62	124.75	109.77
4	A	402	RAM	O3-C3-C2	5.88	121.26	109.99
4	A	402	RAM	C1-O5-C5	-5.16	101.08	112.78
4	A	402	RAM	C6-C5-C4	-4.79	104.22	113.07
4	A	402	RAM	O4-C4-C5	-3.51	101.89	109.67
4	A	402	RAM	O5-C1-C2	2.54	114.70	110.77
4	A	402	RAM	O5-C5-C6	2.52	112.75	107.33
3	A	401	UDP	O3B-PB-O3A	2.13	111.78	104.64

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	401	UDP	PB-O3A-PA-O5'
3	A	401	UDP	PA-O3A-PB-O1B
3	A	401	UDP	O4'-C4'-C5'-O5'
3	A	401	UDP	C2'-C1'-N1-C6
3	A	401	UDP	O4'-C1'-N1-C6
3	A	401	UDP	O4'-C1'-N1-C2
3	A	401	UDP	C4'-C5'-O5'-PA


There are no ring outliers.


2 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	402	RAM	8	0
3	A	401	UDP	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be

highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	294/323 (91%)	1.23	59 (20%) <b>1</b> <b>1</b>	35, 69, 128, 155	0
2	B	25/30 (83%)	2.94	13 (52%) <b>0</b> <b>0</b>	44, 114, 144, 149	0
All	All	319/353 (90%)	1.37	72 (22%) <b>0</b> <b>0</b>	35, 71, 132, 155	0

All (72) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	157	ASN	11.0
1	A	196	THR	7.1
2	B	151	PHE	7.0
2	B	1	ALA	6.8
2	B	160	SER	6.5
1	A	150	GLN	6.1
2	B	156	GLU	6.0
1	A	151	ASP	5.7
1	A	154	PHE	5.6
1	A	54	SER	5.6
1	A	188	ARG	5.5
1	A	191	VAL	5.5
1	A	156	ASP	5.2
1	A	319	ASP	5.2
2	B	159	PRO	5.0
1	A	148	ILE	4.9
1	A	157	LEU	4.8
1	A	203	LEU	4.6
2	B	149	PRO	4.6
1	A	197	GLU	4.5
1	A	239	SER	4.2
1	A	153	LEU	4.1
1	A	41	ASP	3.9
1	A	103	LEU	3.8

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	A	45	ILE	3.7
1	A	320	CYS	3.6
1	A	187	LEU	3.6
1	A	242	LEU	3.5
1	A	185	GLU	3.5
1	A	247	LEU	3.5
1	A	63	ARG	3.5
1	A	194	LEU	3.4
2	B	154	TRP	3.3
1	A	195	ARG	3.3
1	A	40	TYR	3.2
1	A	44	LEU	3.2
1	A	55	SER	3.2
1	A	278	ALA	3.1
1	A	27	ARG	3.1
1	A	285	ILE	3.1
2	B	155	ASP	3.0
1	A	204	LYS	3.0
1	A	158	ASN	3.0
2	B	165	TYR	2.9
1	A	315	LEU	2.9
1	A	236	GLN	2.7
1	A	35	ARG	2.6
1	A	80	ILE	2.6
1	A	316	LEU	2.6
1	A	238	ILE	2.6
1	A	169	HIS	2.6
1	A	296	SER	2.5
2	B	150	LYS	2.5
1	A	198	LEU	2.5
1	A	155	GLY	2.5
1	A	131	TYR	2.5
1	A	227	LYS	2.5
1	A	200	ALA	2.5
1	A	225	ILE	2.4
1	A	192	ALA	2.4
1	A	37	LEU	2.4
1	A	318	SER	2.4
1	A	57	SER	2.4
1	A	184	THR	2.3
2	B	162	ALA	2.2
1	A	234	TYR	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	91	LEU	2.2
2	B	161	SER	2.1
1	A	290	PHE	2.1
1	A	53	ASP	2.0
1	A	289	PRO	2.0
1	A	168	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	RAM	A	402	10/11	0.60	0.19	59,76,94,94	0
3	UDP	A	401	25/25	0.93	0.11	50,67,81,85	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around RAM A 402:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.