



# wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 27, 2020 – 03:14 PM BST

PDB ID : 6TY5  
Title : Crystal structure of human TLR8 in complex with Compound 11  
Authors : Faller, M.; Zink, F.  
Deposited on : 2020-01-15  
Resolution : 2.79 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.13  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.79 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	807	
1	B	807	
2	C	5	
2	E	5	
2	F	5	
2	H	5	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	D	3	 100%
3	G	3	 100%

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 12526 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Toll-like receptor 8.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	738	5838	3735	987	1097	19	0	0	0
1	B	743	5885	3759	1001	1106	19	0	0	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	828	GLU	-	expression tag	UNP Q9NR97
A	829	PHE	-	expression tag	UNP Q9NR97
A	830	LEU	-	expression tag	UNP Q9NR97
A	831	VAL	-	expression tag	UNP Q9NR97
A	832	PRO	-	expression tag	UNP Q9NR97
A	833	ARG	-	expression tag	UNP Q9NR97
B	828	GLU	-	expression tag	UNP Q9NR97
B	829	PHE	-	expression tag	UNP Q9NR97
B	830	LEU	-	expression tag	UNP Q9NR97
B	831	VAL	-	expression tag	UNP Q9NR97
B	832	PRO	-	expression tag	UNP Q9NR97
B	833	ARG	-	expression tag	UNP Q9NR97

- Molecule 2 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	C	5	61	34	2	25	0	0	0
2	E	5	61	34	2	25	0	0	0
2	F	5	61	34	2	25	0	0	0

*Continued on next page...*


Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	H	5	61	34	2	25	0	0	0

- Molecule 3 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	D	3	39	22	2	15	0	0	0
3	G	3	39	22	2	15	0	0	0

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
4	A	1	14	8	1	5	0	0
4	A	1	14	8	1	5	0	0
4	A	1	14	8	1	5	0	0
4	A	1	14	8	1	5	0	0
4	A	1	14	8	1	5	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	B	1	Total	C	N	O	0	0
			14	8	1	5		
4	B	1	Total	C	N	O	0	0
			14	8	1	5		
4	B	1	Total	C	N	O	0	0
			14	8	1	5		
4	B	1	Total	C	N	O	0	0
			14	8	1	5		
4	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 5 is 5-methyl-7-(7-methyl-2-piperidin-4-yl-indazol-5-yl)furo[3,2-c]pyridin-4-one (three-letter code: O0W) (formula: C<sub>21</sub>H<sub>22</sub>N<sub>4</sub>O<sub>2</sub>) (labeled as "Ligand of Interest" by author).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	B	1	Total	C	N	O	0	0
			27	21	4	2		
5	B	1	Total	C	N	O	0	0
			27	21	4	2		


- Molecule 6 is water.

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
6	A	122	Total 122	O 122	0	0
6	B	137	Total 137	O 137	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Toll-like receptor 8



- Molecule 1: Toll-like receptor 8



- Molecule 2: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



Chain C: 

MAG1  
MAG2  
BGLA3  
MAN4  
MAN5

- Molecule 2: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E: 

MAG1  
MAG2  
BGLA3  
MAN4  
MAN5

- Molecule 2: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F: 

MAG1  
MAG2  
BGLA3  
MAN4  
MAN5

- Molecule 2: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain H: 

MAG1  
MAG2  
BGLA3  
MAN4  
MAN5

- Molecule 3: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D: 

MAG1  
MAG2  
BGLA3

- Molecule 3: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G: 

MAG1  
MAG2  
BGLA3

## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	164.81Å 87.72Å 153.46Å 90.00° 120.46° 90.00°	Depositor
Resolution (Å)	74.64 - 2.79 74.64 - 2.79	Depositor EDS
% Data completeness (in resolution range)	96.1 (74.64-2.79) 96.1 (74.64-2.79)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.86 (at 2.77Å)	Xtrriage
Refinement program	BUSTER 2.11.7 (3-OCT-2019)	Depositor
R, $R_{free}$	0.237 , 0.264 0.241 , 0.270	Depositor DCC
$R_{free}$ test set	2315 reflections (5.11%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	72.3	Xtrriage
Anisotropy	0.480	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.28 , 43.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.000 for h,-k,-h-l	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	12526	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	79.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.41% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, NAG, O0W, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.22	0/5955	0.43	0/8089
1	B	0.22	0/6002	0.43	0/8151
All	All	0.22	0/11957	0.43	0/16240

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5838	0	5726	27	0
1	B	5885	0	5802	29	0
2	C	61	0	52	0	0
2	E	61	0	52	0	0
2	F	61	0	52	0	0
2	H	61	0	52	0	0
3	D	39	0	34	0	0
3	G	39	0	34	0	0
4	A	84	0	78	0	0
4	B	84	0	78	0	0
5	B	54	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	A	122	0	0	1	0
6	B	137	0	0	0	0
All	All	12526	0	11960	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 56 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:373:HIS:HA	1:B:400:ASN:HB3	1.83	0.59
1:A:373:HIS:HA	1:A:400:ASN:HB3	1.84	0.59
1:B:45:VAL:HB	1:B:65:VAL:HA	1.86	0.57
1:A:343:ASP:OD2	1:A:469:HIS:HD2	1.88	0.57
1:B:343:ASP:OD2	1:B:469:HIS:HD2	1.88	0.57

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	724/807 (90%)	673 (93%)	49 (7%)	2 (0%)	41 72
1	B	733/807 (91%)	683 (93%)	48 (6%)	2 (0%)	41 72
All	All	1457/1614 (90%)	1356 (93%)	97 (7%)	4 (0%)	41 72

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	330	GLY
1	B	330	GLY

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	378	VAL
1	B	378	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	658/751 (88%)	649 (99%)	9 (1%)	67 90
1	B	665/751 (88%)	658 (99%)	7 (1%)	73 92
All	All	1323/1502 (88%)	1307 (99%)	16 (1%)	71 92

5 of 16 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	465	SER
1	A	701	LEU
1	B	388	GLN
1	A	420	LEU
1	B	416	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	594	ASN
1	A	629	ASN
1	B	593	HIS
1	A	593	HIS
1	B	469	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

26 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	NAG	C	1	1,2	14,14,15	0.30	0	17,19,21	0.47	0
2	NAG	C	2	2	14,14,15	0.29	0	17,19,21	0.59	0
2	BMA	C	3	2	11,11,12	0.24	0	15,15,17	0.71	1 (6%)
2	MAN	C	4	2	11,11,12	0.32	0	15,15,17	0.80	1 (6%)
2	MAN	C	5	2	11,11,12	0.31	0	15,15,17	0.81	1 (6%)
3	NAG	D	1	1,3	14,14,15	0.29	0	17,19,21	0.65	0
3	NAG	D	2	3	14,14,15	0.30	0	17,19,21	0.40	0
3	BMA	D	3	3	11,11,12	0.29	0	15,15,17	0.52	0
2	NAG	E	1	1,2	14,14,15	0.29	0	17,19,21	0.58	1 (5%)
2	NAG	E	2	2	14,14,15	0.28	0	17,19,21	0.50	0
2	BMA	E	3	2	11,11,12	0.26	0	15,15,17	0.45	0
2	MAN	E	4	2	11,11,12	0.30	0	15,15,17	0.68	1 (6%)
2	MAN	E	5	2	11,11,12	0.31	0	15,15,17	0.73	1 (6%)
2	NAG	F	1	1,2	14,14,15	0.28	0	17,19,21	0.48	0
2	NAG	F	2	2	14,14,15	0.27	0	17,19,21	0.60	0
2	BMA	F	3	2	11,11,12	0.24	0	15,15,17	0.72	1 (6%)
2	MAN	F	4	2	11,11,12	0.30	0	15,15,17	0.83	1 (6%)
2	MAN	F	5	2	11,11,12	0.31	0	15,15,17	0.80	1 (6%)
3	NAG	G	1	1,3	14,14,15	0.28	0	17,19,21	0.63	0
3	NAG	G	2	3	14,14,15	0.29	0	17,19,21	0.40	0
3	BMA	G	3	3	11,11,12	0.28	0	15,15,17	0.50	0
2	NAG	H	1	1,2	14,14,15	0.29	0	17,19,21	0.57	1 (5%)
2	NAG	H	2	2	14,14,15	0.28	0	17,19,21	0.51	0
2	BMA	H	3	2	11,11,12	0.26	0	15,15,17	0.54	0
2	MAN	H	4	2	11,11,12	0.31	0	15,15,17	0.68	1 (6%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	MAN	H	5	2	11,11,12	0.32	0	15,15,17	0.75	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	C	2	2	-	0/6/23/26	0/1/1/1
2	BMA	C	3	2	-	0/2/19/22	0/1/1/1
2	MAN	C	4	2	-	0/2/19/22	0/1/1/1
2	MAN	C	5	2	-	0/2/19/22	0/1/1/1
3	NAG	D	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	D	2	3	-	0/6/23/26	0/1/1/1
3	BMA	D	3	3	-	0/2/19/22	0/1/1/1
2	NAG	E	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	E	2	2	-	0/6/23/26	0/1/1/1
2	BMA	E	3	2	-	0/2/19/22	0/1/1/1
2	MAN	E	4	2	-	0/2/19/22	0/1/1/1
2	MAN	E	5	2	-	0/2/19/22	0/1/1/1
2	NAG	F	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	F	2	2	-	0/6/23/26	0/1/1/1
2	BMA	F	3	2	-	0/2/19/22	0/1/1/1
2	MAN	F	4	2	-	0/2/19/22	0/1/1/1
2	MAN	F	5	2	-	0/2/19/22	0/1/1/1
3	NAG	G	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	G	2	3	-	0/6/23/26	0/1/1/1
3	BMA	G	3	3	-	0/2/19/22	0/1/1/1
2	NAG	H	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	H	2	2	-	0/6/23/26	0/1/1/1
2	BMA	H	3	2	-	0/2/19/22	0/1/1/1
2	MAN	H	4	2	-	0/2/19/22	0/1/1/1
2	MAN	H	5	2	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	4	MAN	C1-O5-C5	3.05	116.33	112.19
2	C	5	MAN	C1-O5-C5	3.04	116.31	112.19
2	F	5	MAN	C1-O5-C5	2.97	116.22	112.19

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	4	MAN	C1-O5-C5	2.96	116.21	112.19
2	H	5	MAN	C1-O5-C5	2.71	115.87	112.19


There are no chirality outliers.

There are no torsion outliers.


There are no ring outliers.


No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.














## 5.6 Ligand geometry [i](#)

14 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
4	NAG	A	1015	1	14,14,15	0.28	0	17,19,21	0.65	1 (5%)
5	O0W	B	1121	-	24,31,31	2.42	9 (37%)	26,46,46	1.32	2 (7%)
4	NAG	A	1019	1	14,14,15	0.29	0	17,19,21	0.51	0
4	NAG	B	1116	1	14,14,15	0.27	0	17,19,21	0.64	1 (5%)
4	NAG	A	1009	1	14,14,15	0.28	0	17,19,21	0.41	0
4	NAG	B	1120	1	14,14,15	0.28	0	17,19,21	0.50	0
4	NAG	A	1018	1	14,14,15	0.31	0	17,19,21	0.99	2 (11%)
4	NAG	A	1017	1	14,14,15	0.30	0	17,19,21	0.62	1 (5%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	NAG	A	1016	1	14,14,15	0.29	0	17,19,21	0.58	1 (5%)
5	O0W	B	1101	-	24,31,31	2.33	9 (37%)	26,46,46	1.61	5 (19%)
4	NAG	B	1119	1	14,14,15	0.28	0	17,19,21	0.97	2 (11%)
4	NAG	B	1110	1	14,14,15	0.28	0	17,19,21	0.54	0
4	NAG	B	1117	1	14,14,15	0.28	0	17,19,21	0.58	1 (5%)
4	NAG	B	1118	1	14,14,15	0.29	0	17,19,21	0.59	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	1015	1	-	0/6/23/26	0/1/1/1
5	O0W	B	1121	-	-	0/4/16/16	0/5/5/5
4	NAG	A	1019	1	-	0/6/23/26	0/1/1/1
4	NAG	B	1116	1	-	0/6/23/26	0/1/1/1
4	NAG	A	1009	1	-	0/6/23/26	0/1/1/1
4	NAG	B	1120	1	-	0/6/23/26	0/1/1/1
4	NAG	A	1018	1	-	0/6/23/26	0/1/1/1
4	NAG	A	1017	1	-	0/6/23/26	0/1/1/1
4	NAG	A	1016	1	-	0/6/23/26	0/1/1/1
5	O0W	B	1101	-	-	0/4/16/16	0/5/5/5
4	NAG	B	1119	1	-	0/6/23/26	0/1/1/1
4	NAG	B	1110	1	-	0/6/23/26	0/1/1/1
4	NAG	B	1117	1	-	0/6/23/26	0/1/1/1
4	NAG	B	1118	1	-	0/6/23/26	0/1/1/1

The worst 5 of 18 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	1121	O0W	C8-C7	5.86	1.49	1.39
5	B	1101	O0W	C8-C7	5.38	1.49	1.39
5	B	1101	O0W	C14-C15	-4.67	1.33	1.43
5	B	1121	O0W	C14-C15	-4.65	1.33	1.43
5	B	1121	O0W	C2-N1	-4.42	1.31	1.35

The worst 5 of 17 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	1101	O0W	C21-C22-C17	-3.71	107.53	110.44

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1018	NAG	C1-O5-C5	3.27	116.63	112.19
4	B	1119	NAG	C1-O5-C5	3.26	116.61	112.19
5	B	1101	O0W	C19-C18-C17	-3.13	107.99	110.44
5	B	1121	O0W	C2-N1-N5	-2.76	110.05	111.70


There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	738/807 (91%)	0.13	13 (1%) 68 61	48, 77, 101, 118	0
1	B	743/807 (92%)	0.19	18 (2%) 59 49	46, 78, 124, 141	0
All	All	1481/1614 (91%)	0.16	31 (2%) 63 54	46, 77, 115, 141	0

The worst 5 of 31 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	571	ALA	5.0
1	A	707	LEU	4.2
1	A	710	PHE	3.9
1	B	761	THR	3.7
1	B	32	ARG	3.6

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	MAN	H	5	11/12	0.58	0.23	87,87,88,88	0
2	MAN	F	5	11/12	0.62	0.19	95,96,96,96	0
2	MAN	C	5	11/12	0.67	0.23	95,96,96,96	0
2	MAN	F	4	11/12	0.67	0.38	95,96,96,96	0
2	MAN	E	5	11/12	0.72	0.25	85,86,86,86	0

*Continued on next page...*






*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	BMA	G	3	11/12	0.79	0.24	80,81,81,81	0
2	MAN	H	4	11/12	0.81	0.14	86,87,87,87	0
3	BMA	D	3	11/12	0.81	0.32	71,72,72,72	0
2	BMA	F	3	11/12	0.82	0.14	90,92,94,94	0
2	MAN	C	4	11/12	0.82	0.21	95,95,96,96	0
2	BMA	C	3	11/12	0.84	0.12	91,93,94,94	0
2	BMA	H	3	11/12	0.85	0.15	82,84,85,86	0
2	BMA	E	3	11/12	0.85	0.18	80,82,83,84	0
2	MAN	E	4	11/12	0.86	0.20	84,85,85,85	0
3	NAG	G	2	14/15	0.88	0.15	73,75,77,78	0
3	NAG	D	2	14/15	0.90	0.20	65,66,68,69	0
2	NAG	C	1	14/15	0.93	0.19	78,80,81,82	0
2	NAG	H	1	14/15	0.95	0.14	69,70,71,73	0
2	NAG	F	2	14/15	0.95	0.17	82,83,85,88	0
2	NAG	F	1	14/15	0.95	0.20	77,78,79,81	0
2	NAG	C	2	14/15	0.95	0.15	84,85,87,89	0
2	NAG	E	1	14/15	0.96	0.15	63,64,66,68	0
2	NAG	E	2	14/15	0.96	0.15	70,72,74,77	0
3	NAG	G	1	14/15	0.97	0.17	66,67,69,71	0
3	NAG	D	1	14/15	0.97	0.14	59,60,61,63	0
2	NAG	H	2	14/15	0.97	0.17	74,75,77,80	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around Chain C:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around Chain E:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around Chain F:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







**Electron density around Chain H:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around Chain D:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	NAG	B	1118	14/15	0.70	0.24	85,86,87,87	0
4	NAG	B	1119	14/15	0.74	0.34	93,94,94,94	0
4	NAG	A	1016	14/15	0.74	0.33	86,86,87,87	0
4	NAG	B	1117	14/15	0.78	0.35	79,80,80,80	0
4	NAG	A	1017	14/15	0.79	0.32	94,94,95,95	0
4	NAG	A	1018	14/15	0.80	0.43	85,86,86,86	0
4	NAG	A	1015	14/15	0.82	0.24	91,91,92,92	0
4	NAG	A	1019	14/15	0.82	0.33	108,109,109,109	0
4	NAG	B	1120	14/15	0.87	0.32	115,115,115,115	0
4	NAG	B	1110	14/15	0.87	0.19	78,78,79,79	0
4	NAG	B	1116	14/15	0.94	0.16	84,85,85,85	0
4	NAG	A	1009	14/15	0.94	0.15	67,67,68,68	0
5	O0W	B	1101	27/27	0.96	0.17	56,57,57,57	0
5	O0W	B	1121	27/27	0.97	0.17	58,59,60,60	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.5 Other polymers [i](#)

There are no such residues in this entry.