



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 5, 2024 – 03:35 AM EST

PDB ID : 1U0C
Title : Y33C Mutant of Homing endonuclease I-CreI
Authors : Sussman, D.; Chadsey, M.; Fauce, S.; Engel, A.; Bruett, A.; Monnat, R.;
Stoddard, B.L.; Seligman, L.M.
Deposited on : 2004-07-13
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	1-C	24	
1	2-C	24	
2	1-D	24	
2	2-D	24	
3	1-A	163	
3	1-B	163	
3	2-A	163	
3	2-B	163	

2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 7062 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called 5'-D(*GP*CP*TP*AP*AP*AP*CP*GP*TP*CP*GP*TP*GP*AP*GP*AP*CP*AP*GP*TP*TP*AP*CP*G)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
1	1-C	24	Total 493	C 235	N 95	O 140	P 23	0	0	0
1	2-C	24	Total 493	C 235	N 95	O 140	P 23	0	0	0

- Molecule 2 is a DNA chain called 5'-D(*CP*GP*TP*AP*AP*CP*TP*GP*TP*CP*TP*CP*AP*CP*GP*AP*CP*GP*TP*TP*TP*AP*GP*C)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	1-D	24	Total 485	C 233	N 85	O 144	P 23	0	0	0
2	2-D	24	Total 485	C 233	N 85	O 144	P 23	0	0	0

- Molecule 3 is a protein called DNA endonuclease I-CreI.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	1-A	152	Total 1230	C 790	N 209	O 229	S 2	0	0	0
3	2-A	152	Total 1230	C 790	N 209	O 229	S 2	0	0	0
3	1-B	152	Total 1230	C 790	N 209	O 229	S 2	0	0	0
3	2-B	152	Total 1230	C 790	N 209	O 229	S 2	0	0	0

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	33	CYS	TYR	engineered mutation	UNP P05725
A	42	THR	ALA	engineered mutation	UNP P05725

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	47	GLU	GLN	engineered mutation	UNP P05725
A	110	GLU	TRP	engineered mutation	UNP P05725
A	111	GLN	ARG	engineered mutation	UNP P05725
B	333	CYS	TYR	engineered mutation	UNP P05725
B	342	THR	ALA	engineered mutation	UNP P05725
B	347	GLU	GLN	engineered mutation	UNP P05725
B	410	GLU	TRP	engineered mutation	UNP P05725
B	411	GLN	ARG	engineered mutation	UNP P05725

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	1-C	1	Total Mg 1 1	0	0
4	2-C	1	Total Mg 1 1	0	0
4	1-A	1	Total Mg 1 1	0	0
4	2-A	1	Total Mg 1 1	0	0
4	1-B	1	Total Mg 1 1	0	0
4	2-B	1	Total Mg 1 1	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	1-C	11	Total O 11 11	0	0
5	2-C	12	Total O 12 12	0	0
5	1-D	17	Total O 17 17	0	0
5	2-D	18	Total O 18 18	0	0
5	1-A	34	Total O 34 34	0	0
5	2-A	32	Total O 32 32	0	0
5	1-B	28	Total O 28 28	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	2-B	28	Total	O	0	0
			28	28		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 5'-D(*GP*CP*TP*AP*AP*AP*CP*GP*TP*CP*GP*TP*GP*AP*GP*AP*CP*AP*GP*TP*TP*AP*CP*G)-3'

Chain 1-C: 




- Molecule 1: 5'-D(*GP*CP*TP*AP*AP*AP*CP*GP*TP*CP*GP*TP*GP*AP*GP*AP*CP*AP*GP*TP*TP*AP*CP*G)-3'

Chain 2-C: 



- Molecule 2: 5'-D(*CP*GP*TP*AP*AP*CP*TP*GP*TP*CP*TP*CP*AP*CP*GP*AP*CP*GP*TP*TP*TP*AP*GP*C)-3'

Chain 1-D: 



- Molecule 2: 5'-D(*CP*GP*TP*AP*AP*CP*TP*GP*TP*CP*TP*CP*AP*CP*GP*AP*CP*GP*TP*TP*TP*AP*GP*C)-3'

Chain 2-D: 




- Molecule 3: DNA endonuclease I-CreI

Chain 1-A: 






- Molecule 3: DNA endonuclease I-CreI



- Molecule 3: DNA endonuclease I-CreI



- Molecule 3: DNA endonuclease I-CreI



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	43.09Å 68.02Å 87.18Å 90.00° 92.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.50 43.56 – 2.50	Depositor EDS
% Data completeness (in resolution range)	(Not available) (50.00-2.50) 96.1 (43.56-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.77 (at 2.51Å)	Xtrriage
Refinement program	CNS	Depositor
R, R_{free}	0.214 , 0.277 0.272 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	38.2	Xtrriage
Anisotropy	0.240	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.50 , 44.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.043 for h,-k,-l	Xtrriage
F_o, F_c correlation	0.86	EDS
Total number of atoms	7062	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.02% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	1-C	0.30	0/554	0.71	0/854
1	2-C	0.29	0/554	0.67	0/854
2	1-D	0.29	0/542	0.75	0/834
2	2-D	0.29	0/542	0.71	0/834
3	1-A	0.43	0/1252	0.67	0/1690
3	1-B	0.42	0/1252	0.68	1/1690 (0.1%)
3	2-A	0.43	0/1252	0.67	0/1690
3	2-B	0.42	0/1252	0.68	1/1690 (0.1%)
All	All	0.39	0/7200	0.69	2/10136 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	1-C	0	1
1	2-C	0	2
2	1-D	0	2
2	2-D	0	1
All	All	0	6

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	1-B	376	TYR	N-CA-C	-5.33	96.61	111.00
3	2-B	376	TYR	N-CA-C	-5.33	96.61	111.00

There are no chirality outliers.

5 of 6 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	1-C	514	DA	Sidechain
2	1-D	567	DC	Sidechain
2	1-D	568	DG	Sidechain
1	2-C	517	DC	Sidechain
1	2-C	518	DA	Sidechain

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	1-C	493	0	271	26	0
1	2-C	493	0	271	26	0
2	1-D	485	0	273	24	0
2	2-D	485	0	273	17	0
3	1-A	1230	0	1264	62	0
3	1-B	1230	0	1264	64	0
3	2-A	1230	0	1264	61	0
3	2-B	1230	0	1264	55	0
4	1-A	1	0	0	0	0
4	1-B	1	0	0	0	0
4	1-C	1	0	0	0	0
4	2-A	1	0	0	0	0
4	2-B	1	0	0	0	0
4	2-C	1	0	0	0	0
5	1-A	34	0	0	2	0
5	1-B	28	0	0	2	0
5	1-C	11	0	0	2	0
5	1-D	17	0	0	0	0
5	2-A	32	0	0	2	0
5	2-B	28	0	0	2	0
5	2-C	12	0	0	0	0
5	2-D	18	0	0	1	0
All	All	7062	0	6144	291	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 22.

The worst 5 of 291 close contacts within the same asymmetric unit are listed below, sorted by

their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:505:DA:H2''	1:C:506:DA:H5''	1.50	0.91
3:A:4:LYS:HE2	3:A:4:LYS:H	1.38	0.87
3:A:4:LYS:HE2	3:A:4:LYS:H	1.38	0.87
3:B:322:SER:OG	3:B:324:ILE:HD11	1.75	0.87
3:B:322:SER:OG	3:B:324:ILE:HD11	1.75	0.87

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	1-A	150/163 (92%)	139 (93%)	11 (7%)	0	100	100
3	1-B	150/163 (92%)	138 (92%)	11 (7%)	1 (1%)	22	39
3	2-A	150/163 (92%)	139 (93%)	11 (7%)	0	100	100
3	2-B	150/163 (92%)	138 (92%)	11 (7%)	1 (1%)	22	39
All	All	600/652 (92%)	554 (92%)	44 (7%)	2 (0%)	41	61

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	1-B	450	ALA
3	2-B	450	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	1-A	139/150 (93%)	131 (94%)	8 (6%)	20	38
3	1-B	139/150 (93%)	125 (90%)	14 (10%)	7	14
3	2-A	139/150 (93%)	131 (94%)	8 (6%)	20	38
3	2-B	139/150 (93%)	125 (90%)	14 (10%)	7	14
All	All	556/600 (93%)	512 (92%)	44 (8%)	12	24

5 of 44 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	2-A	128	TRP
3	2-B	381	ILE
3	2-A	152	LEU
3	2-B	368	ARG
3	2-B	401	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 13 such sidechains are listed below:

Mol	Chain	Res	Type
3	2-A	26	GLN
3	2-A	31	GLN
3	2-B	350	GLN
3	2-A	103	ASN
3	2-B	331	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 6 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers

Unable to reproduce the depositors R factor - this section is therefore empty.