



Full wwPDB NMR Structure Validation Report ⓘ

Nov 8, 2023 – 05:57 AM EST

PDB ID : 8U1T
BMRB ID : 31104
Title : SARS-CoV-2 Envelope Protein Transmembrane Domain: Dimeric Structure
Determined by Solid-State NMR
Authors : Zhang, R.; Qin, H.; Prasad, R.; Fu, R.; Zhou, H.X.; Cross, T.
Deposited on : 2023-09-02

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
wwPDB-RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
wwPDB-ShiftChecker : v1.2
BMRB Restraints Analysis : v1.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLID-STATE NMR

The overall completeness of chemical shifts assignment is 3%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	29	 72% 14% 10%
1	B	29	 76% 14% 10%

2 Ensemble composition and analysis

This entry contains 14 models. Model 9 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:13-A:37, B:12-B:37 (51)	0.44	9

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 2 single-model clusters were found.

Cluster number	Models
1	1, 3, 4, 9, 10, 11, 13, 14
2	8, 12
3	2, 5
Single-model clusters	6; 7

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 864 atoms, of which 466 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Envelope small membrane protein.

Mol	Chain	Residues	Atoms					Trace
			Total	C	H	N	O	
1	A	26	432	142	233	27	30	0
1	B	26	432	142	233	27	30	0

There are 6 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	9	SER	-	expression tag	UNP P0DTC4
A	10	ASN	-	expression tag	UNP P0DTC4
A	11	ALA	-	expression tag	UNP P0DTC4
B	9	SER	-	expression tag	UNP P0DTC4
B	10	ASN	-	expression tag	UNP P0DTC4
B	11	ALA	-	expression tag	UNP P0DTC4

4 Residue-property plots


4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Envelope small membrane protein



- Molecule 1: Envelope small membrane protein




4.2 Scores per residue for each member of the ensemble


Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1

- Molecule 1: Envelope small membrane protein




- Molecule 1: Envelope small membrane protein



4.2.2 Score per residue for model 2

- Molecule 1: Envelope small membrane protein




- Molecule 1: Envelope small membrane protein




4.2.3 Score per residue for model 3

- Molecule 1: Envelope small membrane protein




- Molecule 1: Envelope small membrane protein



4.2.4 Score per residue for model 4

- Molecule 1: Envelope small membrane protein



- Molecule 1: Envelope small membrane protein




4.2.5 Score per residue for model 5

- Molecule 1: Envelope small membrane protein




- Molecule 1: Envelope small membrane protein




4.2.6 Score per residue for model 6

- Molecule 1: Envelope small membrane protein




- Molecule 1: Envelope small membrane protein




4.2.7 Score per residue for model 7

- Molecule 1: Envelope small membrane protein




- Molecule 1: Envelope small membrane protein



4.2.8 Score per residue for model 8

- Molecule 1: Envelope small membrane protein



- Molecule 1: Envelope small membrane protein




4.2.9 Score per residue for model 9 (medoid)

- Molecule 1: Envelope small membrane protein




- Molecule 1: Envelope small membrane protein




4.2.10 Score per residue for model 10

- Molecule 1: Envelope small membrane protein



- Molecule 1: Envelope small membrane protein



4.2.11 Score per residue for model 11

- Molecule 1: Envelope small membrane protein

Chain A:  69% 14% 10%



- Molecule 1: Envelope small membrane protein

Chain B:  72% 17% 10%



4.2.12 Score per residue for model 12


- Molecule 1: Envelope small membrane protein

Chain A:  62% 24% 10%



- Molecule 1: Envelope small membrane protein

Chain B:  76% 7% 7% 10%



4.2.13 Score per residue for model 13

- Molecule 1: Envelope small membrane protein

Chain A:  66% 21% 10%



- Molecule 1: Envelope small membrane protein

Chain B:  72% 17% 10%



4.2.14 Score per residue for model 14


- Molecule 1: Envelope small membrane protein

Chain A:  66% 17% . . 10%



- Molecule 1: Envelope small membrane protein

Chain B:  59% 31% 10%



5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the 100 calculated structures, 14 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR NIH	structure calculation	
NAMD	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	22
Number of shifts mapped to atoms	22
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	3%

Note: This is a solid-state NMR structure, where hydrogen atoms are typically not assigned a chemical shift value, which may lead to lower completeness of assignment measure.

6 Model quality i

6.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	1.58±0.09	1±1/193 (0.6± 0.4%)	1.94±0.11	4±2/265 (1.5± 0.8%)
1	B	1.56±0.09	1±1/201 (0.4± 0.6%)	1.97±0.10	5±2/276 (1.8± 0.8%)
All	All	1.57	27/5516 (0.5%)	1.96	127/7574 (1.7%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	0.4±0.6
1	B	0.0±0.0	0.5±0.6
All	All	0	12

All unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
1	A	16	SER	CA-CB	7.99	1.65	1.52	10	4
1	B	27	LEU	CA-CB	6.82	1.69	1.53	2	1
1	B	17	VAL	CB-CG2	6.29	1.66	1.52	5	1
1	B	23	PHE	CB-CG	-6.27	1.40	1.51	5	1
1	A	23	PHE	CB-CG	6.04	1.61	1.51	10	1
1	B	20	PHE	CE2-CZ	5.83	1.48	1.37	14	1
1	A	26	PHE	CG-CD1	5.68	1.47	1.38	4	1
1	A	23	PHE	CG-CD2	5.62	1.47	1.38	9	2
1	B	23	PHE	CE1-CZ	5.48	1.47	1.37	13	1
1	A	20	PHE	CB-CG	5.41	1.60	1.51	5	1
1	A	17	VAL	N-CA	5.38	1.57	1.46	2	1
1	B	26	PHE	CB-CG	-5.37	1.42	1.51	9	1
1	B	20	PHE	CB-CG	-5.27	1.42	1.51	4	1
1	B	18	LEU	N-CA	-5.26	1.35	1.46	4	1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
1	B	19	LEU	CA-CB	5.20	1.65	1.53	10	1
1	B	13	ILE	N-CA	5.20	1.56	1.46	14	1
1	A	26	PHE	CE2-CZ	5.12	1.47	1.37	11	1
1	A	34	LEU	N-CA	5.11	1.56	1.46	3	1
1	A	23	PHE	CE2-CZ	5.10	1.47	1.37	7	1
1	A	20	PHE	CG-CD2	5.08	1.46	1.38	13	1
1	B	37	LEU	CA-CB	5.04	1.65	1.53	5	1
1	B	20	PHE	CG-CD1	5.04	1.46	1.38	5	1
1	A	37	LEU	N-CA	-5.01	1.36	1.46	14	1

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	B	26	PHE	CB-CG-CD2	13.61	130.32	120.80	5	8
1	A	20	PHE	CB-CG-CD2	10.44	128.11	120.80	3	2
1	A	20	PHE	CB-CG-CD1	-10.09	113.73	120.80	9	5
1	B	23	PHE	CB-CG-CD1	-9.51	114.14	120.80	11	6
1	A	26	PHE	CB-CG-CD1	-9.23	114.34	120.80	13	3
1	B	26	PHE	CB-CG-CD1	-9.08	114.45	120.80	14	2
1	B	23	PHE	CB-CG-CD2	8.54	126.78	120.80	2	3
1	A	14	VAL	CA-CB-CG1	7.66	122.39	110.90	1	1
1	B	17	VAL	CA-CB-CG2	-7.37	99.85	110.90	8	1
1	B	20	PHE	CB-CG-CD2	-7.20	115.76	120.80	6	4
1	A	23	PHE	CB-CG-CD1	7.17	125.82	120.80	1	3
1	A	36	ALA	N-CA-CB	-6.98	100.32	110.10	9	1
1	B	20	PHE	CG-CD2-CE2	-6.94	113.16	120.80	12	1
1	A	14	VAL	CA-CB-CG2	-6.89	100.57	110.90	7	2
1	A	29	VAL	CG1-CB-CG2	-6.72	100.15	110.90	1	1
1	B	25	VAL	CA-CB-CG2	-6.51	101.13	110.90	7	2
1	B	29	VAL	CA-CB-CG1	6.47	120.60	110.90	11	2
1	A	30	THR	N-CA-CB	6.28	122.23	110.30	5	4
1	B	20	PHE	CG-CD1-CE1	-6.19	113.99	120.80	2	1
1	B	32	ALA	N-CA-CB	-6.19	101.43	110.10	14	1
1	A	19	LEU	CB-CG-CD1	6.19	121.52	111.00	13	1
1	A	21	LEU	CB-CG-CD2	6.18	121.51	111.00	8	2
1	B	27	LEU	O-C-N	-6.16	112.84	122.70	10	2
1	A	21	LEU	N-CA-CB	5.99	122.39	110.40	9	1
1	A	24	VAL	CA-CB-CG1	5.99	119.88	110.90	6	1
1	A	26	PHE	CB-CG-CD2	-5.98	116.61	120.80	14	1
1	A	29	VAL	CA-CB-CG1	5.92	119.78	110.90	8	1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	B	12	LEU	N-CA-CB	-5.92	98.56	110.40	11	1
1	B	29	VAL	CB-CA-C	5.92	122.64	111.40	4	1
1	B	35	THR	N-CA-CB	5.85	121.41	110.30	2	1
1	A	19	LEU	O-C-N	-5.83	113.37	122.70	13	2
1	B	35	THR	CA-CB-OG1	5.83	121.23	109.00	14	1
1	B	20	PHE	CB-CG-CD1	-5.82	116.72	120.80	7	1
1	B	31	LEU	CB-CG-CD2	5.77	120.81	111.00	13	2
1	B	24	VAL	CA-CB-CG2	-5.75	102.28	110.90	7	2
1	B	24	VAL	CG1-CB-CG2	-5.75	101.71	110.90	1	1
1	A	36	ALA	O-C-N	-5.71	113.57	122.70	10	1
1	B	19	LEU	N-CA-CB	5.69	121.78	110.40	8	1
1	A	20	PHE	CZ-CE2-CD2	-5.59	113.39	120.10	13	1
1	B	28	LEU	CB-CA-C	5.58	120.81	110.20	14	1
1	B	14	VAL	CG1-CB-CG2	-5.58	101.97	110.90	12	2
1	A	24	VAL	O-C-N	-5.57	113.78	122.70	11	1
1	A	23	PHE	O-C-N	-5.55	113.82	122.70	2	1
1	A	23	PHE	CB-CG-CD2	5.55	124.69	120.80	14	2
1	B	26	PHE	CD1-CE1-CZ	-5.54	113.46	120.10	13	1
1	A	30	THR	CA-CB-CG2	-5.53	104.66	112.40	12	1
1	B	12	LEU	O-C-N	-5.51	113.88	122.70	3	2
1	A	28	LEU	CB-CG-CD2	-5.50	101.65	111.00	5	1
1	B	14	VAL	CA-CB-CG2	-5.50	102.66	110.90	2	1
1	B	35	THR	OG1-CB-CG2	-5.48	97.39	110.00	5	2
1	B	19	LEU	CB-CG-CD2	5.47	120.30	111.00	9	1
1	B	22	ALA	O-C-N	-5.46	113.97	122.70	2	1
1	B	35	THR	CA-CB-CG2	-5.46	104.76	112.40	4	1
1	B	24	VAL	O-C-N	-5.45	113.98	122.70	2	1
1	B	20	PHE	O-C-N	-5.44	113.99	122.70	1	1
1	B	21	LEU	CB-CG-CD2	-5.38	101.85	111.00	10	2
1	B	21	LEU	CB-CG-CD1	5.37	120.13	111.00	10	1
1	A	35	THR	CA-CB-CG2	-5.37	104.89	112.40	7	2
1	B	26	PHE	CG-CD2-CE2	5.35	126.69	120.80	10	1
1	B	30	THR	N-CA-CB	5.27	120.31	110.30	3	1
1	A	18	LEU	O-C-N	-5.25	114.31	122.70	2	1
1	A	20	PHE	CG-CD2-CE2	5.24	126.56	120.80	1	1
1	B	21	LEU	CB-CA-C	5.23	120.13	110.20	10	2
1	B	35	THR	O-C-N	-5.23	114.33	122.70	12	1
1	A	30	THR	O-C-N	-5.22	114.34	122.70	9	1
1	A	36	ALA	CB-CA-C	5.22	117.93	110.10	7	1
1	A	33	ILE	CA-CB-CG2	-5.21	100.49	110.90	6	1
1	B	23	PHE	CB-CA-C	5.20	120.79	110.40	8	1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	33	ILE	O-C-N	-5.17	114.43	122.70	2	1
1	A	24	VAL	CA-CB-CG2	-5.14	103.19	110.90	11	2
1	A	23	PHE	CG-CD2-CE2	5.12	126.43	120.80	5	1
1	B	22	ALA	CB-CA-C	5.12	117.78	110.10	1	1
1	B	16	SER	N-CA-CB	5.12	118.17	110.50	4	1
1	B	36	ALA	O-C-N	-5.09	114.56	122.70	4	1
1	A	31	LEU	CB-CG-CD1	-5.07	102.39	111.00	9	1
1	A	29	VAL	O-C-N	-5.05	114.61	122.70	12	1
1	A	20	PHE	O-C-N	-5.05	114.62	122.70	9	1
1	A	37	LEU	CB-CA-C	5.04	119.77	110.20	6	1
1	A	29	VAL	CA-CB-CG2	-5.03	103.36	110.90	10	1
1	B	25	VAL	CA-CB-CG1	5.02	118.43	110.90	8	1
1	A	18	LEU	CB-CA-C	-5.00	100.69	110.20	5	1

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	B	26	PHE	Sidechain	3
1	A	26	PHE	Sidechain	2
1	B	23	PHE	Sidechain	2
1	A	20	PHE	Sidechain	2
1	B	20	PHE	Sidechain	2
1	A	23	PHE	Sidechain	1

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	B	199	233	232	0±1
1	A	191	222	222	0±0
All	All	5460	6370	6356	5

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:B:13:ILE:HD12	1:B:13:ILE:H	0.71	1.45	6	4
1:B:13:ILE:H	1:B:13:ILE:CD1	0.43	2.23	6	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	24/29 (83%)	24±0 (100±1%)	0±0 (0±1%)	0±0 (0±0%)	100	100
1	B	24/29 (83%)	24±0 (100±1%)	0±0 (0±1%)	0±0 (0±0%)	100	100
All	All	672/812 (83%)	670 (100%)	2 (0%)	0 (0%)	100	100

There are no Ramachandran outliers.

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	22/25 (88%)	21±1 (95±2%)	1±1 (5±2%)	27	76
1	B	23/25 (92%)	22±1 (97±4%)	1±1 (3±4%)	40	87
All	All	630/700 (90%)	603 (96%)	27 (4%)	33	81

All 13 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	21	LEU	5
1	B	13	ILE	4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	37	LEU	3
1	A	19	LEU	2
1	A	15	ASN	2
1	A	18	LEU	2
1	B	21	LEU	2
1	B	33	ILE	2
1	B	29	VAL	1
1	A	35	THR	1
1	B	19	LEU	1
1	A	28	LEU	1
1	B	25	VAL	1

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 3% for the well-defined parts and 3% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: *assigned_chemical_shifts_1*

7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	22
Number of shifts mapped to atoms	22
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	22

7.1.2 Chemical shift referencing (i)

No chemical shift referencing corrections were calculated (not enough data).

7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 3%, i.e. 22 atoms were assigned a chemical shift out of a possible 774. 0 out of 27 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	22/255 (9%)	0/102 (0%)	0/102 (0%)	22/51 (43%)
Sidechain	0/459 (0%)	0/317 (0%)	0/140 (0%)	0/2 (0%)
Aromatic	0/60 (0%)	0/30 (0%)	0/30 (0%)	0/0 (—%)
Overall	22/774 (3%)	0/449 (0%)	0/272 (0%)	22/53 (42%)

Note: This is a solid-state NMR structure, where hydrogen atoms are typically not assigned a chemical shift value, which may lead to lower completeness of assignment measure.

The following table shows the completeness of the chemical shift assignments for the full structure.

The overall completeness is 3%, i.e. 22 atoms were assigned a chemical shift out of a possible 792. 0 out of 28 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹H	¹³C	¹⁵N
Backbone	22/260 (8%)	0/104 (0%)	0/104 (0%)	22/52 (42%)
Sidechain	0/472 (0%)	0/326 (0%)	0/144 (0%)	0/2 (0%)
Aromatic	0/60 (0%)	0/30 (0%)	0/30 (0%)	0/0 (—%)
Overall	22/792 (3%)	0/460 (0%)	0/278 (0%)	22/54 (41%)

Note: This is a solid-state NMR structure, where hydrogen atoms are typically not assigned a chemical shift value, which may lead to lower completeness of assignment measure.

7.1.4 Statistically unusual chemical shifts [i](#)


The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

List Id	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	22	ALA	N	224.50	106.13 – 140.55	29.4
1	A	32	ALA	N	224.50	106.13 – 140.55	29.4
1	A	36	ALA	N	215.30	106.13 – 140.55	26.7
1	A	18	LEU	N	223.00	102.77 – 140.89	26.5
1	A	19	LEU	N	223.00	102.77 – 140.89	26.5
1	A	21	LEU	N	223.00	102.77 – 140.89	26.5
1	A	27	LEU	N	223.00	102.77 – 140.89	26.5
1	A	28	LEU	N	223.00	102.77 – 140.89	26.5
1	A	31	LEU	N	223.00	102.77 – 140.89	26.5
1	A	34	LEU	N	223.00	102.77 – 140.89	26.5
1	A	20	PHE	N	220.00	99.93 – 140.82	24.4
1	A	23	PHE	N	220.00	99.93 – 140.82	24.4
1	A	26	PHE	N	220.00	99.93 – 140.82	24.4
1	A	13	ILE	N	220.00	100.55 – 142.30	23.6
1	A	33	ILE	N	220.00	100.55 – 142.30	23.6
1	A	14	VAL	N	223.00	99.23 – 142.92	23.3
1	A	17	VAL	N	223.00	99.23 – 142.92	23.3
1	A	24	VAL	N	223.00	99.23 – 142.92	23.3
1	A	25	VAL	N	223.00	99.23 – 142.92	23.3
1	A	29	VAL	N	223.00	99.23 – 142.92	23.3
1	A	30	THR	N	222.60	91.89 – 138.78	22.9
1	A	35	THR	N	206.30	91.89 – 138.78	19.4

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:



8 NMR restraints analysis

No restraints data found

9 Distance violation analysis

No distance restraints data found

10 Dihedral-angle violation analysis

No dihedral-angle restraints found