

wwPDB X-ray Structure Validation Summary Report (i)

Sep 17, 2023 – 02:49 AM EDT

:	4U3F
:	Cytochrome bc1 complex from chicken with designed inhibitor bound
:	Huang, LS.; Zhu, XL.; Yang, G.F.; Berry, E.A.
:	2014-07-21
:	3.23 Å(reported)
	: : : :

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.35.1
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 3.23 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Motria	Whole archive	Similar resolution			
Metric	$(\# {\rm Entries})$	$(\# \text{Entries, resolution range}(\text{\AA}))$			
R _{free}	130704	1619 (3.28-3.20)			
Clashscore	141614	1755 (3.28-3.20)			
Ramachandran outliers	138981	1728 (3.28-3.20)			
Sidechain outliers	138945	1727 (3.28-3.20)			
RSRZ outliers	127900	1567 (3.28-3.20)			

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	Λ	446	2%	1.00/	
	A	440	<u> </u>	16%	••
1	Ν	446	81%	16%	••
0	D	4.4.1	2%		
	D	441	72% 21%	•	5%
2	Ο	441	73% 20%) •	•
9	C	200	%		
3		380	84%	15%	•



Mol	Chain	Length	Quality of chain			
3	Р	380	% 83%	15% •		
4	D	241	% 86%	13% •		
4	Q	241	86%	13% •		
5	Е	196	77%	20% •		
5	R	196	76%	20% ••		
6	F	110	79%	13% 8%		
6	S	110	79%	12% • 8%		
7	G	81	% 81%	16% ••		
7	Т	81	85%	11% ••		
8	Н	77	.% • 78%	10% 12%		
8	U	77	68%	19% • 12%		
9	Ι	76	9% 39% 12% 7%	42%		
9	V	76	9% 37% 13% 5%	45%		
10	J	61	10%	11%		
10	W	61	5%	20% ••		

Continued from previous page...

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
11	PEE	Р	411	-	-	-	Х
3	FME	С	1	-	-	-	Х



2 Entry composition (i)

There are 21 unique types of molecules in this entry. The entry contains 32790 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Mitochondrial ubiquinol-cytochrome-c reductase complex core protein i.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	А	443	Total 3442	C 2157	N 606	O 658	S 21	0	0	0
1	Ν	442	Total 3437	C 2154	N 605	O 657	S 21	0	0	0

• Molecule 2 is a protein called Mitochondrial ubiquinol-cytochrome-c reductase complex core protein 2.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
9	B 491	491	Total	С	Ν	0	\mathbf{S}	0	0	0
Z D	421	3137	1971	544	613	9	0	0	0	
9	0	499	Total	С	Ν	0	S	0	0	0
2 0	422	3143	1974	545	614	10	0	0	0	

• Molecule 3 is a protein called Cytochrome b.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
3	С	380	Total 3021	C 2025	N 478	O 505	S 13	0	0	0
3	Р	379	Total 3022	C 2025	N 480	O 505	S 12	0	1	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
С	1	FME	-	expression tag	UNP P18946
Р	1	FME	-	expression tag	UNP P18946

• Molecule 4 is a protein called Mitochondrial cytochrome c1, heme protein.



Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
4	4 D	241	Total	С	Ν	0	\mathbf{S}	0	Ο	0
4			1898	1212	327	347	12		0	0
4	0	241	Total	С	Ν	0	S	0	0	0
4	4 Q	241	1898	1212	327	347	12	0	0	0

• Molecule 5 is a protein called Cytochrome b-c1 complex subunit Rieske, mitochondrial.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
Б	r D	196	Total	С	Ν	0	S	0	0	0
D E	Ľ		1513	952	263	292	6			
5	D	106	Total	С	Ν	0	S	0	0	0
D R	n	190	1509	950	263	290	6	0		U

• Molecule 6 is a protein called Cytochrome b-c1 complex subunit 7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	F	101	Total	С	Ν	0	S	0	0	0
0	Ľ	101	891	570	159	159	3	0	0	0
6	q	101	Total	С	Ν	Ο	\mathbf{S}	0	0	0
6 0		101	891	570	159	159	3	0	0	0

• Molecule 7 is a protein called Mitochondrial ubiquinol-cytochrome c reductase ubiquinonebinding protein qp-c.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
7	C	C 80		С	Ν	Ο	0	0	0
1	G	80	672	437	119	116	0	0	0
7	т	70	Total	С	Ν	Ο	0	0	0
1	L	19	662	432	117	113	0	0	0

• Molecule 8 is a protein called Cytochrome b-c1 complex subunit 6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
8	Н	68	Total 562	C 343	N 104	0 110	${f S}{5}$	0	0	0
8	U	68	Total 558	C 341	N 104	O 108	${ m S}{ m 5}$	0	0	0

• Molecule 9 is a protein called Cytochrome b-c1 complex subunit Rieske, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
9	Ι	44	Total 266	C 157	N 54	O 53	${ m S} { m 2}$	0	0	1



Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
0	V	49	Total	С	Ν	Ο	\mathbf{S}	0	0	1
9 V	v	42	265	157	55	51	2	0	0	1

There are 30 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
Ι	1	AME	-	expression tag	UNP Q5ZLR5
Ι	29	UNK	LEU	see remark 999	UNP Q5ZLR5
Ι	30	UNK	ALA	see remark 999	UNP Q5ZLR5
Ι	31	UNK	PRO	see remark 999	UNP Q5ZLR5
Ι	32	UNK	ALA	see remark 999	UNP Q5ZLR5
Ι	33	UNK	ALA	see remark 999	UNP Q5ZLR5
Ι	37	UNK	LEU	see remark 999	UNP Q5ZLR5
Ι	38	UNK	ARG	see remark 999	UNP Q5ZLR5
Ι	39	UNK	ALA	see remark 999	UNP Q5ZLR5
Ι	40	UNK	GLU	see remark 999	UNP Q5ZLR5
Ι	41	UNK	LYS	see remark 999	UNP Q5ZLR5
Ι	42	UNK	VAL	see remark 999	UNP Q5ZLR5
Ι	43	UNK	VAL	see remark 999	UNP Q5ZLR5
Ι	44	UNK	LEU	see remark 999	UNP Q5ZLR5
Ι	45	UNK	ASP	see remark 999	UNP Q5ZLR5
V	-1	AME	-	expression tag	UNP Q5ZLR5
V	27	UNK	LEU	see remark 999	UNP Q5ZLR5
V	28	UNK	ALA	see remark 999	UNP Q5ZLR5
V	29	UNK	PRO	see remark 999	UNP Q5ZLR5
V	30	UNK	ALA	see remark 999	UNP Q5ZLR5
V	31	UNK	ALA	see remark 999	UNP Q5ZLR5
V	32	UNK	LEU	see remark 999	UNP Q5ZLR5
V	33	UNK	ARG	see remark 999	UNP Q5ZLR5
V	37	UNK	ALA	see remark 999	UNP Q5ZLR5
V	38	UNK	GLU	see remark 999	UNP Q5ZLR5
V	39	UNK	LYS	see remark 999	UNP Q5ZLR5
V	40	UNK	VAL	see remark 999	UNP Q5ZLR5
V	41	UNK	VAL	see remark 999	UNP Q5ZLR5
V	42	UNK	LEU	see remark 999	UNP Q5ZLR5
V	43	UNK	ASP	see remark 999	UNP Q5ZLR5

• Molecule 10 is a protein called Mitochondrial ubiquinol-cytochrome c reductase 7.2 kda protein.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace	
10	т	61	Total	С	Ν	0	0	0	0
10	J	01	497	321	87	89	0	0	
10	117	60	Total	С	Ν	0	0	0	1
10	vv	00	479	311	86	82	0	0	

• Molecule 11 is 1,2-dioleoyl-sn-glycero-3-phosphoethanolamine (three-letter code: PEE) (formula: $C_{41}H_{78}NO_8P$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
11	А	1	Total C N O P 26 16 1 8 1	0	0
11	С	1	Total C N O P 49 39 1 8 1	0	0
11	С	1	Total C N O P 11 5 1 4 1	0	0
11	С	1	Total C N O P 47 37 1 8 1	0	0
11	С	1	Total C O 15 13 2	0	0
11	Е	1	Total C N O P 48 38 1 8 1	0	0
11	Е	1	Total C 15 15	0	0
11	Ν	1	Total C N O P 8 2 1 4 1	0	0
11	Р	1	Total C O 15 13 2	0	0



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf						
11	D	1	Total C N O P	0	0						
	Г	L	49 39 1 8 1	0	0						
11	D	1	Total C N O P	0	0						
	Г	L	41 31 1 8 1	0	0						
11	D	1	Total C O	0	0						
	1	L	12 10 2	0	0						
11	D	1	Total C N O P	0	0						
	1	L	25 15 1 8 1	0	0						
11	D	1	Total C N O P	0	0						
	n		49 39 1 8 1	U	U						

Continued from previous page...

• Molecule 12 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
12	2 C	1	Total	С	Fe	Ν	Ο	0	0
12	U	1	43	34	1	4	4	0	0
19	С	1	Total	С	Fe	Ν	0	0	0
12	U	1	43	34	1	4	4	0	0
10	D	1	Total	С	Fe	Ν	0	0	0
12	1	1	43	34	1	4	4	0	0
19	D	1	Total	С	Fe	Ν	0	0	0
	L_		43	34	1	4	4	0	U

• Molecule 13 is methyl (2E)-3-methoxy-2-(2-{[(5-methoxy-1,3-benzothiazol-2-yl)sulfanyl]met hyl}phenyl)prop-2-enoate (three-letter code: Y52) (formula: $C_{20}H_{19}NO_4S_2$).





Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
13	С	1	Total	С	Ν	0	S	0	0
10	U	T	27	20	1	4	2	0	0
12	D	1	Total	С	Ν	Ο	S	0	0
10	1	L	27	20	1	4	2	0	U

• Molecule 14 is UBIQUINONE-10 (three-letter code: U10) (formula: $C_{59}H_{90}O_4$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
14	С	1	Total C O 19 15 4	0	0
14	Р	1	Total C O 19 15 4	0	0



• Molecule 15 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula: $C_6H_{13}NO_4S$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf		
15	С	1	Total 12	C 6	N 1	0 4	S 1	0	0

• Molecule 16 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Ato	\mathbf{pms}		ZeroOcc	AltConf
16	С	1	Total 6	С 3	O 3	0	0



Continued from previous page...

Mol	Chain	Residues	Ato	\mathbf{ms}		ZeroOcc	AltConf
16	Р	1	Total 6	$\begin{array}{c} \mathrm{C} \\ \mathrm{3} \end{array}$	O 3	0	0

• Molecule 17 is HEME C (three-letter code: HEC) (formula: $C_{34}H_{34}FeN_4O_4$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
17	D	1	Total 43	С 34	Fe 1	N 4	0 4	0	0
17	Q	1	Total 43	C 34	Fe 1	N 4	0 4	0	0





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
18	Л	1	Total	С	Ο	Р	0	0
10	D	1	42	23	17	2	0	0
18	С	1	Total	С	Ο	Р	0	0
10	o G	1	40	21	17	2	0	0
19	0	1	Total	С	Ο	Р	0	0
10	Q	L	42	23	17	2	0	0
18	Т	1	Total	С	Ο	Р	0	0
10	1	L	40	21	17	2	0	0

• Molecule 19 is octyl beta-D-glucopyranoside (three-letter code: BOG) (formula: $C_{14}H_{28}O_6$).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
19	D	1	Total C O 20 14 6	0	0
19	Р	1	Total C O 20 14 6	0	0
19	Q	1	Total C O 20 14 6	0	0

• Molecule 20 is FE2/S2 (INORGANIC) CLUSTER (three-letter code: FES) (formula: Fe_2S_2).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
20	Е	1	TotalFeS422	0	0
20	R	1	TotalFeS422	0	0

• Molecule 21 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
21	А	2	Total O 2 2	0	0
21	В	1	Total O 1 1	0	0
21	С	2	Total O 2 2	0	0
21	F	1	Total O 1 1	0	0
21	Ν	1	Total O 1 1	0	0



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
21	О	2	Total O 2 2	0	0
21	Р	2	Total O 2 2	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Mitochondrial ubiquinol-cytochrome-c reductase complex core protein i





• Molecule 2: Mitochondrial ubiquinol-cytochrome-c reductase complex core protein 2



• Molecule 2: Mitochondrial ubiquinol-cytochrome-c reductase complex core protein 2



• Molecule 4: Mitochondrial cytochrome c1, heme protein







 \bullet Molecule 10: Mitochondrial ubiquinol-cytochrome c
 reductase 7.2 kda protein



• Molecule 10: Mitochondrial ubiquinol-cytochrome c reductase 7.2 kda protein





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	167.25Å 181.48Å 239.63Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
$\mathbf{Posolution} \left(\overset{\circ}{\mathbf{A}} \right)$	15.00 - 3.23	Depositor
Resolution (A)	54.30 - 3.23	EDS
% Data completeness	98.3 (15.00-3.23)	Depositor
(in resolution range)	82.2 (54.30-3.23)	EDS
R_{merge}	(Not available)	Depositor
R _{sym}	0.30	Depositor
$< I/\sigma(I) > 1$	$1.05 (at 3.26 \text{\AA})$	Xtriage
Refinement program	PHENIX (phenix.refine: dev_1745)	Depositor
B B.	0.236 , 0.284	Depositor
$\mathbf{n}, \mathbf{n}_{free}$	0.243 , 0.286	DCC
R_{free} test set	2286 reflections $(1.98%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	49.5	Xtriage
Anisotropy	0.509	Xtriage
Bulk solvent $k_{sol}(e/A^3)$, $B_{sol}(A^2)$	0.34 , 55.2	EDS
L-test for $twinning^2$	$ < L > = 0.48, < L^2 > = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.87	EDS
Total number of atoms	32790	wwPDB-VP
Average B, all atoms $(Å^2)$	59.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 2.07% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: MES, PEE, FME, HEC, FES, GOL, CDL, U10, BOG, Y52, HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond	lengths	Bond	angles
	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	0.21	0/3513	0.40	0/4760
1	Ν	0.21	0/3508	0.39	0/4753
2	В	0.21	0/3192	0.40	0/4330
2	0	0.22	0/3198	0.40	0/4339
3	С	0.22	0/3114	0.39	0/4263
3	Р	0.22	0/3102	0.39	0/4245
4	D	0.20	0/1956	0.37	0/2658
4	Q	0.20	0/1956	0.37	0/2658
5	Е	0.20	0/1547	0.38	0/2103
5	R	0.20	0/1543	0.38	0/2098
6	F	0.21	0/911	0.35	0/1219
6	S	0.21	0/911	0.35	0/1219
7	G	0.22	0/694	0.39	0/941
7	Т	0.22	0/684	0.39	0/929
8	Н	0.21	0/570	0.36	0/763
8	U	0.21	0/566	0.40	0/758
9	Ι	0.20	0/208	0.48	0/279
9	V	0.21	0/215	0.56	0/288
10	J	0.21	0/508	0.34	0/682
10	W	0.21	0/490	0.35	0/660
All	All	0.21	0/32386	0.39	0/43945

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	А	0	1

There are no bond length outliers.



There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	А	216	PHE	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	3442	0	3354	41	0
1	N	3437	0	3349	40	0
2	В	3137	0	3131	55	0
2	0	3143	0	3135	56	0
3	С	3021	0	3068	35	0
3	Р	3022	0	3064	39	0
4	D	1898	0	1846	17	0
4	Q	1898	0	1846	16	0
5	Е	1513	0	1480	21	0
5	R	1509	0	1476	22	0
6	F	891	0	893	6	0
6	S	891	0	893	8	0
7	G	672	0	653	11	0
7	Т	662	0	645	10	0
8	Н	562	0	545	3	0
8	U	558	0	541	8	0
9	Ι	266	0	223	14	0
9	V	265	0	239	11	0
10	J	497	0	490	3	0
10	W	479	0	478	6	0
11	А	26	0	26	1	0
11	С	122	0	174	2	0
11	Е	63	0	99	5	0
11	N	8	0	7	0	0
11	Р	142	0	193	7	0
11	R	49	0	75	5	0
12	С	86	0	60	7	0
12	Р	86	0	60	8	0



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
13	С	27	0	19	2	0
13	Р	27	0	19	1	0
14	С	19	0	17	1	0
14	Р	19	0	17	3	0
15	С	12	0	13	0	0
16	С	6	0	8	1	0
16	Р	6	0	8	1	0
17	D	43	0	30	0	0
17	Q	43	0	30	0	0
18	D	42	0	28	1	0
18	G	40	0	24	1	0
18	Q	42	0	28	0	0
18	Т	40	0	24	3	0
19	D	20	0	28	0	0
19	Р	20	0	28	3	0
19	Q	20	0	28	0	0
20	Ε	4	0	0	0	0
20	R	4	0	0	0	0
21	А	2	0	0	0	0
21	В	1	0	0	1	0
21	С	2	0	0	0	0
21	F	1	0	0	0	0
21	N	1	0	0	0	0
21	0	2	0	0	1	0
21	Р	2	0	0	0	0
All	All	32790	0	32392	385	0

Continued from previous page...

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 385 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
9:V:37:UNK:HG3	9:V:38:UNK:H	1.37	0.90
9:V:49:LEU:HB2	9:V:55:MET:HB3	1.57	0.85
9:I:37:UNK:HG3	9:I:38:UNK:N	1.94	0.81
9:I:37:UNK:HG3	9:I:38:UNK:H	1.47	0.78
2:B:361:LYS:HD2	2:B:403:ASP:HA	1.67	0.76

There are no symmetry-related clashes.



5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	441/446~(99%)	421 (96%)	16 (4%)	4 (1%)	17 52
1	Ν	440/446 (99%)	419 (95%)	17 (4%)	4 (1%)	17 52
2	В	419/441 (95%)	397~(95%)	18 (4%)	4 (1%)	15 50
2	Ο	420/441 (95%)	397 (94%)	17 (4%)	6 (1%)	11 43
3	С	378/380~(100%)	365~(97%)	11 (3%)	2~(0%)	29 64
3	Р	378/380~(100%)	364 (96%)	12 (3%)	2(0%)	29 64
4	D	239/241~(99%)	226 (95%)	11 (5%)	2(1%)	19 55
4	Q	239/241~(99%)	225 (94%)	12 (5%)	2(1%)	19 55
5	Е	194/196~(99%)	168 (87%)	20 (10%)	6(3%)	4 24
5	R	194/196~(99%)	175 (90%)	14 (7%)	5(3%)	5 28
6	F	99/110 (90%)	97~(98%)	2 (2%)	0	100 100
6	S	99/110 (90%)	98~(99%)	1 (1%)	0	100 100
7	G	78/81~(96%)	74 (95%)	4 (5%)	0	100 100
7	Т	77/81~(95%)	72 (94%)	5 (6%)	0	100 100
8	Н	66/77~(86%)	64 (97%)	2(3%)	0	100 100
8	U	66/77~(86%)	56~(85%)	9 (14%)	1 (2%)	10 41
9	Ι	28/76~(37%)	17 (61%)	9(32%)	2~(7%)	1 7
9	V	28/76~(37%)	19 (68%)	7 (25%)	2(7%)	1 7
10	J	59/61~(97%)	55 (93%)	3 (5%)	1 (2%)	9 39
10	W	58/61~(95%)	54 (93%)	3 (5%)	1 (2%)	9 39
All	All	4000/4218 (95%)	3763 (94%)	193 (5%)	44 (1%)	14 48

5 of 44 Ramachandran outliers are listed below:

Mol	Chain	Res	Type	
2	В	390	GLY	
Continued on next nage				



Continued from previous page...

	v	1	1 0
Mol	Chain	\mathbf{Res}	Type
2	0	19	PRO
2	0	226	ILE
2	0	390	GLY
9	V	59	SER

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the side chain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	entiles
1	А	365/368~(99%)	342 (94%)	23~(6%)	18	50
1	Ν	365/368~(99%)	339~(93%)	26 (7%)	14	45
2	В	331/347~(95%)	312 (94%)	19 (6%)	20	54
2	Ο	332/347~(96%)	313 (94%)	19 (6%)	20	54
3	С	328/328~(100%)	309 (94%)	19 (6%)	20	53
3	Р	327/328~(100%)	310 (95%)	17 (5%)	23	56
4	D	200/200~(100%)	191 (96%)	9 (4%)	27	61
4	Q	200/200~(100%)	190 (95%)	10 (5%)	24	58
5	Е	166/166~(100%)	156 (94%)	10 (6%)	19	52
5	R	165/166~(99%)	150 (91%)	15 (9%)	9	32
6	F	93/96~(97%)	89~(96%)	4 (4%)	29	62
6	S	93/96~(97%)	89~(96%)	4 (4%)	29	62
7	G	71/71~(100%)	67~(94%)	4 (6%)	21	55
7	Т	70/71~(99%)	67~(96%)	3 (4%)	29	62
8	Н	64/71~(90%)	60 (94%)	4 (6%)	18	50
8	U	63/71~(89%)	60~(95%)	3~(5%)	25	59
9	Ι	22/45~(49%)	18 (82%)	4 (18%)	1	7
9	V	23/45~(51%)	17 (74%)	6 (26%)	0	1
10	J	49/49~(100%)	46 (94%)	3~(6%)	18	51
10	W	47/49~(96%)	42 (89%)	5 (11%)	6	26



Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	3374/3482~(97%)	3167~(94%)	207~(6%)	18 51

5 of 207 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	Ν	248	LEU
2	0	402	ILE
9	V	55	MET
1	N	350	THR
2	0	160	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 23 such sidechains are listed below:

Mol	Chain	Res	Type
1	Ν	271	HIS
3	Р	313	GLN
2	0	363	GLN
3	Р	332	ASN
3	С	332	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Iol Type Chain Res		Link	B	ond leng	gths	Bond angles			
	Type	Ullalli	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	FME	С	1	3	7,8,10	1.95	1 (14%)	5,8,11	1.39	1 (20%)



In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	FME	С	1	3	-	0/5/8/11	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	$\operatorname{Ideal}(\operatorname{\AA})$
3	С	1	FME	CN-N	-5.04	1.33	1.46

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
3	С	1	FME	CN-N-CA	2.65	121.87	113.64

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

36 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Mol Type Chain Rea		Ros Link	Bond lengths			Bond angles			
Moi Typ	туре	Ullalli	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z >2
12	HEM	Р	403	3	41,50,50	1.46	4 (9%)	45,82,82	1.41	7 (15%)



Mal	Turne	Chain	Dec	Tink	Bo	ond leng	ths	Bond angles		
	туре	Chain	nes		Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2
11	PEE	Е	503	-	14,14,50	1.00	1 (7%)	$13,\!13,\!55$	0.90	0
15	MES	С	506	-	12,12,12	2.26	1 (8%)	14,16,16	1.23	3 (21%)
17	HEC	Q	501	4	32,50,50	1.90	3 (9%)	24,82,82	1.43	3 (12%)
11	PEE	Е	502	-	47,47,50	1.24	4 (8%)	50,52,55	1.15	4 (8%)
19	BOG	Q	503	-	20,20,20	0.55	1 (5%)	25,25,25	0.61	0
20	FES	R	501	5	0,4,4	-	-	-		
11	PEE	Ν	501	-	7,7,50	1.40	1 (14%)	$9,\!9,\!55$	1.25	1 (11%)
11	PEE	R	502	-	48,48,50	1.24	4 (8%)	$51,\!53,\!55$	1.17	6 (11%)
13	Y52	Р	404	-	27,29,29	0.67	0	30,39,39	1.07	2 (6%)
17	HEC	D	501	4	32,50,50	1.89	3 (9%)	24,82,82	1.42	3 (12%)
20	FES	Е	501	5	0,4,4	_	-	_		
18	CDL	Q	502	-	41,41,99	1.30	4 (9%)	47,53,111	1.29	6 (12%)
11	PEE	С	510	-	14,14,50	2.27	3 (21%)	14,14,55	1.20	1 (7%)
13	Y52	С	503	-	27,29,29	0.68	0	30,39,39	1.10	3 (10%)
11	PEE	А	501	-	25,25,50	1.35	2 (8%)	28,30,55	1.56	5 (17%)
11	PEE	Р	409	-	40,40,50	1.18	3 (7%)	43,45,55	1.19	3 (6%)
14	U10	С	504	_	19,19,63	2.12	2 (10%)	23,26,79	1.48	2 (8%)
18	CDL	D	502	-	41,41,99	1.31	4 (9%)	47,53,111	1.31	5 (10%)
11	PEE	Р	407	-	48,48,50	1.22	4 (8%)	51,53,55	1.09	2 (3%)
11	PEE	Р	411	-	24,24,50	1.30	2 (8%)	27,29,55	1.40	4 (14%)
18	CDL	Т	101	-	39,39,99	1.35	4 (10%)	45,51,111	1.38	5 (11%)
11	PEE	С	505	-	48,48,50	1.20	4 (8%)	51,53,55	1.22	4 (7%)
16	GOL	Р	408	-	5,5,5	0.37	0	5,5,5	0.29	0
12	HEM	С	501	3	41,50,50	1.48	5 (12%)	45,82,82	1.44	8 (17%)
11	PEE	С	507	-	10,10,50	0.51	0	11,12,55	0.55	0
14	U10	Р	405	-	$19,\!19,\!63$	2.18	2 (10%)	$23,\!26,\!79$	1.60	4 (17%)
18	CDL	G	101	-	39,39,99	1.34	4 (10%)	45,51,111	1.36	4 (8%)
19	BOG	Р	406	-	20,20,20	0.52	0	25,25,25	0.62	0
19	BOG	D	503	-	20,20,20	0.55	1 (5%)	25,25,25	0.64	0
11	PEE	С	509	-	46,46,50	1.28	4 (8%)	49,51,55	1.19	4 (8%)
11	PEE	Р	410	-	11,11,50	1.77	2(18%)	11,11,55	1.34	1 (9%)
12	HEM	Р	402	3	41,50,50	1.48	5 (12%)	45,82,82	1.44	8 (17%)
16	GOL	С	508	-	$5,\!5,\!5$	0.36	0	$5,\!5,\!5$	0.34	0
11	PEE	Р	401	-	14,14,50	2.26	3 (21%)	14,14,55	1.29	2 (14%)
12	HEM	С	502	3	$41,\!50,\!50$	1.46	5 (12%)	45,82,82	1.43	7 (15%)



In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
12	HEM	Р	403	3	-	4/12/54/54	-
11	PEE	Е	503	-	-	1/12/12/54	-
15	MES	С	506	-	-	3/6/14/14	0/1/1/1
17	HEC	Q	501	4	-	2/10/54/54	-
11	PEE	Е	502	-	-	21/51/51/54	-
19	BOG	Q	503	-	-	2/11/31/31	0/1/1/1
20	FES	R	501	5	_	-	0/1/1/1
11	PEE	N	501	-	-	0/5/5/54	-
11	PEE	R	502	-	-	21/52/52/54	-
13	Y52	Р	404	-	-	0/18/20/20	0/3/3/3
17	HEC	D	501	4	-	2/10/54/54	-
20	FES	Е	501	5	-	-	0/1/1/1
18	CDL	Q	502	-	-	14/51/51/110	-
11	PEE	С	510	-	-	7/12/12/54	-
13	Y52	С	503	-	-	0/18/20/20	0/3/3/3
11	PEE	А	501	-	-	11/29/29/54	-
11	PEE	Р	409	-	_	19/44/44/54	_
14	U10	С	504	-	-	4/11/35/87	0/1/1/1
18	CDL	D	502	-	-	16/51/51/110	-
11	PEE	Р	407	-	-	15/52/52/54	-
11	PEE	Р	411	-	-	9/28/28/54	-
18	CDL	Т	101	-	-	17/49/49/110	-
11	PEE	С	505	-	-	23/52/52/54	-
16	GOL	Р	408	-	-	4/4/4/4	-
12	HEM	С	501	3	-	6/12/54/54	-
11	PEE	С	507	-	-	5/10/10/54	-
14	U10	Р	405	-	-	4/11/35/87	0/1/1/1
18	CDL	G	101	-	-	14/49/49/110	-
19	BOG	Р	406	-	-	3/11/31/31	0/1/1/1
19	BOG	D	503	-	-	0/11/31/31	0/1/1/1
11	PEE	С	509	-	-	17/50/50/54	-
11	PEE	Р	410	-	-	5/9/9/54	-
12	HEM	Р	402	3	-	6/12/54/54	_



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
16	GOL	С	508	-	-	4/4/4/4	-
11	PEE	Р	401	-	-	7/12/12/54	-
12	HEM	С	502	3	-	4/12/54/54	-

Continued from previous page...

The worst 5 of 85 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
14	Р	405	U10	C6-C1	8.65	1.51	1.35
14	С	504	U10	C6-C1	8.36	1.50	1.35
15	С	506	MES	C8-S	-7.58	1.66	1.77
11	С	510	PEE	O4-C10	7.19	1.46	1.22
11	Р	401	PEE	O4-C10	7.15	1.45	1.22

The worst 5 of 107 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
18	Т	101	CDL	OB6-CB5-C51	4.49	121.18	111.50
11	С	505	PEE	O2-C10-C11	4.48	121.17	111.50
14	С	504	U10	C7-C8-C9	-4.45	119.38	126.79
18	Q	502	CDL	OB6-CB5-C51	4.40	120.98	111.50
11	Р	411	PEE	O2-C10-C11	4.40	120.97	111.50

There are no chirality outliers.

5 of 270 torsion outliers are listed below:

\mathbf{Mol}	Chain	\mathbf{Res}	Type	Atoms
11	А	501	PEE	C1-O3P-P-O1P
11	А	501	PEE	C1-O3P-P-O4P
11	А	501	PEE	C4-O4P-P-O1P
11	С	505	PEE	C11-C10-O2-C2
11	С	505	PEE	C1-C2-O2-C10

There are no ring outliers.

23 monomers are involved in 48 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
12	Р	403	HEM	3	0
11	Е	503	PEE	2	0
11	Е	502	PEE	4	0
11	R	502	PEE	5	0



4U3F	
------	--

Mol	Chain	Res	Type	Clashes	Symm-Clashes
13	Р	404	Y52	1	0
13	С	503	Y52	2	0
11	А	501	PEE	1	0
11	Р	409	PEE	3	0
14	С	504	U10	1	0
18	D	502	CDL	1	0
11	Р	407	PEE	2	0
11	Р	411	PEE	1	0
18	Т	101	CDL	3	0
16	Р	408	GOL	1	0
12	С	501	HEM	4	0
14	Р	405	U10	3	0
18	G	101	CDL	1	0
19	Р	406	BOG	3	0
11	С	509	PEE	2	0
12	Р	402	HEM	5	0
16	С	508	GOL	1	0
11	Р	401	PEE	1	0
12	С	502	HEM	3	0

Continued from previous page...

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


























































5.7 Other polymers (i)

There are no such residues in this entry.



5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle RSRZ \rangle$	#RSRZ>2	$OWAB(Å^2)$	Q < 0.9
1	А	443/446~(99%)	0.05	8 (1%) 68 58	28, 53, 78, 101	0
1	Ν	442/446~(99%)	0.15	9 (2%) 65 54	36, 57, 81, 99	0
2	В	421/441~(95%)	0.24	7 (1%) 70 60	47, 67, 102, 123	0
2	Ο	422/441~(95%)	0.17	8 (1%) 66 57	35, 63, 86, 104	0
3	С	379/380~(99%)	-0.17	4 (1%) 80 73	22, 33, 70, 102	0
3	Р	379/380~(99%)	0.03	2 (0%) 91 87	27, 51, 78, 104	0
4	D	241/241 (100%)	-0.14	3 (1%) 79 70	25, 37, 73, 95	0
4	Q	$241/241 \ (100\%)$	0.14	5 (2%) 63 53	39, 59, 85, 117	0
5	Е	196/196~(100%)	1.61	75 (38%) 0 0	28, 114, 166, 182	9 (4%)
5	R	196/196~(100%)	1.00	41 (20%) 1 1	36, 81, 131, 141	8 (4%)
6	F	101/110 (91%)	-0.17	0 100 100	25, 37, 53, 78	0
6	S	101/110 (91%)	0.22	2 (1%) 65 54	46, 59, 86, 102	0
7	G	80/81~(98%)	0.18	1 (1%) 77 68	28, 45, 73, 89	0
7	Т	79/81~(97%)	0.65	9 (11%) 5 4	40, 70, 122, 127	0
8	Н	68/77~(88%)	0.14	1 (1%) 73 64	35, 50, 68, 101	0
8	U	68/77~(88%)	1.45	20 (29%) 0 0	66, 91, 113, 117	0
9	Ι	30/76~(39%)	1.01	7 (23%) 0 1	55, 77, 116, 120	0
9	V	30/76~(39%)	1.22	7 (23%) 0 1	47, 78, 119, 130	0
10	J	61/61~(100%)	0.81	6 (9%) 7 6	34, 46, 79, 129	0
10	W	60/61~(98%)	0.74	3 (5%) 28 18	41, 55, 92, 117	0
All	All	$403\overline{8/4218} \ (95\%)$	0.26	218 (5%) 25 16	22, 56, 107, 182	17 (0%)

The worst 5 of 218 RSRZ outliers are listed below:



Mol	Chain	Res	Type	RSRZ
5	Ε	163	SER	14.2
5	R	163	SER	7.5
5	Ε	162	GLY	7.4
10	J	62	SER	7.2
5	Е	172	ARG	6.6

6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathbf{\AA}^2)$	Q < 0.9
3	FME	С	1	9/11	0.70	0.41	$72,\!88,\!96,\!99$	0

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(Å^2)$	Q<0.9
11	PEE	А	501	26/51	0.72	0.33	27,87,106,123	0
11	PEE	Ν	501	8/51	0.72	0.28	57,79,96,112	0
11	PEE	Р	409	41/51	0.72	0.34	54,69,112,128	0
18	CDL	Q	502	42/100	0.73	0.36	40,99,121,136	0
11	PEE	Р	411	25/51	0.74	0.41	$36,\!60,\!75,\!80$	25
11	PEE	С	507	11/51	0.76	0.38	$25,\!51,\!62,\!63$	11
11	PEE	R	502	49/51	0.76	0.39	33,64,79,88	0
11	PEE	С	509	47/51	0.76	0.32	22,48,88,107	0
11	PEE	Р	401	15/51	0.78	0.37	41,58,88,89	0
19	BOG	Р	406	20/20	0.79	0.27	39,74,96,98	0
14	U10	Р	405	19/63	0.80	0.39	60,75,88,91	0
11	PEE	E	502	48/51	0.81	0.33	36,55,80,86	0
18	CDL	Т	101	40/100	0.81	0.32	44,85,97,98	0

Continued on next page...



4U	3F
----	----

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B -factors($Å^2$)	Q<0.9
11	PEE	Е	503	15/51	0.81	0.33	22,49,63,68	0
18	CDL	D	502	42/100	0.83	0.30	46,80,99,127	0
15	MES	С	506	12/12	0.85	0.33	45,69,109,122	0
18	CDL	G	101	40/100	0.86	0.27	24,56,69,73	0
11	PEE	Р	407	49/51	0.86	0.28	45,68,84,89	0
11	PEE	Р	410	12/51	0.86	0.26	30,37,46,47	0
14	U10	С	504	19/63	0.86	0.32	45,59,67,70	0
11	PEE	С	510	15/51	0.87	0.25	25,54,81,84	0
19	BOG	D	503	20/20	0.88	0.31	29,46,54,57	0
11	PEE	С	505	49/51	0.91	0.22	22,43,54,57	0
16	GOL	С	508	6/6	0.91	0.24	34,35,38,43	0
19	BOG	Q	503	20/20	0.92	0.24	34,54,62,63	0
16	GOL	Р	408	6/6	0.93	0.26	42,43,49,51	0
13	Y52	Р	404	27/27	0.95	0.22	38,46,58,63	0
13	Y52	С	503	27/27	0.96	0.18	25,33,45,51	0
12	HEM	С	501	43/43	0.96	0.23	18,24,32,35	0
12	HEM	Р	402	43/43	0.96	0.26	26,32,43,53	0
17	HEC	D	501	43/43	0.96	0.21	26,34,40,42	0
17	HEC	Q	501	43/43	0.96	0.23	44,51,63,71	0
12	HEM	Р	403	43/43	0.96	0.25	$27,\!33,\!50,\!53$	0
20	FES	R	501	4/4	0.96	0.13	71,84,92,107	0
12	HEM	С	502	43/43	0.97	0.24	19,24,32,37	0
20	FES	Е	501	4/4	0.98	0.11	92,103,103,106	4

Continued from previous page...

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




























































































6.5 Other polymers (i)

There are no such residues in this entry.

