

Full wwPDB X-ray Structure Validation Report (i)

Jun 25, 2024 – 10:27 AM EDT

NA ternary complex

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 2.92 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R_{free}	130704	2307 (2.94-2.90)
Clashscore	141614	2531 (2.94-2.90)
Ramachandran outliers	138981	2462 (2.94-2.90)
Sidechain outliers	138945	2464 (2.94-2.90)
RSRZ outliers	127900	2248 (2.94-2.90)
RNA backbone	3102	1001 (3.18-2.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of c	hain		
1	А	1130	5% 80%		15%) • •
2	В	112	59%	18%	12%	12%
3	С	38	^{3%} 71%		24%	5%
4	D	8	88%			12%



2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 11907 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called CRISPR-associated endonuclease C2c1.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace		
1	А	1085	Total 8864	$ m C \ 5569$	N 1653	O 1611	S 10	Se 21	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	0	SER	-	expression tag	UNP T0D7A2
А	570	ALA	ASP	engineered mutation	UNP T0D7A2
А	848	ALA	GLU	engineered mutation	UNP T0D7A2
А	977	ALA	ASP	engineered mutation	UNP T0D7A2

• Molecule 2 is a RNA chain called sgRNA.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
2	В	99	Total 2127	C 949	N 396	O 683	Р 99	0	0	0

• Molecule 3 is a DNA chain called Target DNA strand.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
3	С	36	Total	C	N 105	0	Р	0	0	0
			732	349	125	222	36			

• Molecule 4 is a DNA chain called Non-target DNA strand.

Mol	Chain	Residues		At	\mathbf{oms}			ZeroOcc	AltConf	Trace
4	D	8	Total 162	C 79	N 26	O 50	Р 7	0	0	0

• Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O_4S).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	А	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
5	А	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
5	А	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0

• Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	А	6	Total O 6 6	0	0
6	В	1	Total O 1 1	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: CRISPR-associated endonuclease C2c1

• Molecule 2: sgRNA

VAL LEU GLN GLN ASP SER ASP CYS GLU ASN THR THR THR





A106 A107 C111

• Molecule 3: Target DNA strand



4 Data and refinement statistics (i)

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants	116.75Å 184.81Å 214.29Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
$Resolution(\AA)$	49.35 - 2.92	Depositor
Resolution (A)	102.52 - 2.92	EDS
% Data completeness	99.3 (49.35-2.92)	Depositor
(in resolution range)	99.3 (102.52 - 2.92)	EDS
R _{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$1.47 (at 2.91 \text{\AA})$	Xtriage
Refinement program	PHENIX (1.11.1_2575: ???)	Depositor
D D .	0.205 , 0.244	Depositor
n, n_{free}	0.207 , 0.243	DCC
R_{free} test set	2425 reflections $(4.82%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	80.1	Xtriage
Anisotropy	0.123	Xtriage
Bulk solvent $k_{sol}(e/A^3), B_{sol}(A^2)$	0.31 , 54.7	EDS
L-test for $twinning^2$	$ < L >=0.49, < L^2>=0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	11907	wwPDB-VP
Average B, all atoms $(Å^2)$	94.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 2.87% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bo	nd lengths	Bond angles	
	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	0.24	1/9036~(0.0%)	0.39	0/12139
2	В	0.16	0/2380	0.74	0/3706
3	С	0.50	0/817	0.95	0/1258
4	D	0.49	0/180	1.02	0/277
All	All	0.26	1/12413~(0.0%)	0.55	0/17380

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	$\operatorname{Ideal}(\operatorname{\AA})$
1	A	58	GLU	CD-OE2	6.91	1.33	1.25

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	8864	0	8758	98	0
2	В	2127	0	1075	21	0
3	С	732	0	408	10	0
4	D	162	0	94	2	0
5	А	15	0	0	0	0
6	А	6	0	0	1	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	В	1	0	0	0	0
All	All	11907	0	10335	113	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (113) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom 1	Atom 2	Interatomic	Clash	
Atom-1	Atom-2	distance (Å)	overlap (Å)	
1:A:130:ASP:HB3	1:A:133:ALA:HB2	1.62	0.82	
1:A:1085:ILE:HD13	1:A:1100:VAL:HG22	1.67	0.77	
1:A:425:LYS:HE2	1:A:434:VAL:HG11	1.73	0.71	
1:A:678:VAL:HG21	1:A:693:GLU:HG2	1.72	0.71	
1:A:755:GLY:HA2	2:B:33:C:H4'	1.73	0.70	
1:A:331:ARG:NH2	3:C:-1:DT:O4	2.24	0.69	
1:A:742:ARG:NH2	2:B:35:A:OP1	2.27	0.67	
1:A:746:ARG:NH2	2:B:60:C:OP1	2.27	0.64	
1:A:866:GLN:HE22	3:C:-2:DT:H4'	1.63	0.63	
1:A:994:ILE:HD12	1:A:998:ARG:HB3	1.80	0.63	
1:A:761:ILE:HD13	1:A:805:LYS:HG2	1.79	0.63	
1:A:991:ASP:O	1:A:1020:ARG:NH1	2.29	0.62	
1:A:432:ARG:NH1	6:A:1301:HOH:O	2.32	0.62	
2:B:68:C:H42	2:B:85:G:H1	1.45	0.61	
1:A:397:LEU:H	1:A:459:ILE:HD11	1.67	0.59	
1:A:569:VAL:HB	1:A:847:LEU:HD23	1.84	0.59	
1:A:569:VAL:HG22	1:A:579:ILE:HG22	1.84	0.59	
1:A:616:ARG:NH1	1:A:819:GLU:OE1	2.34	0.58	
1:A:397:LEU:HD12	1:A:458:PRO:HG2	1.84	0.58	
1:A:260:GLN:HB3	1:A:263:LEU:HD13	1.84	0.58	
1:A:418:ILE:HD13	1:A:502:LEU:HD23	1.84	0.58	
1:A:893:MSE:HE3	1:A:988:LEU:HA	1.86	0.58	
1:A:900:ARG:NH2	3:C:-6:DC:O2	2.37	0.57	
1:A:507:ARG:NH2	3:C:21:DG:OP1	2.38	0.57	
3:C:10:DC:H2"	3:C:11:DT:H5"	1.87	0.56	
1:A:33:VAL:HG12	1:A:383:PRO:HG3	1.87	0.55	
1:A:754:GLY:HA3	1:A:760:GLN:HG3	1.89	0.55	
1:A:123:LYS:HE2	1:A:175:THR:HG21	1.88	0.54	
1:A:614:HIS:ND1	2:B:6:G:OP1	2.41	0.54	
3:C:-1:DT:H2'	3:C:-1:DT:O2	2.08	0.53	
2:B:86:U:H5'	2:B:87:G:H5'	1.88	0.53	
1:A:918:ARG:HG3	2:B:13:G:OP1	2.09	0.53	



		Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
2:B:69:U:O2'	2:B:70:U:O5'	2.26	0.53
1:A:218:ARG:NH1	4:D:5:DG:OP2	2.28	0.52
1:A:22:LEU:HD22	1:A:481:ILE:HD13	1.92	0.52
1:A:15:MSE:HE1	1:A:439:VAL:HA	1.92	0.52
1:A:54:ASN:HA	1:A:789:LYS:HE2	1.92	0.51
1:A:745:ILE:HD11	2:B:34:A:OP1	2.09	0.51
1:A:824:VAL:HB	1:A:838:LYS:HD3	1.92	0.51
1:A:131:LYS:HG3	1:A:189:PRO:HG3	1.92	0.51
1:A:607:ASN:OD1	1:A:607:ASN:N	2.43	0.51
1:A:415:ARG:NH2	2:B:27:G:OP1	2.44	0.50
1:A:745:ILE:HD12	1:A:745:ILE:H	1.77	0.50
1:A:688:TRP:HB2	1:A:729:GLN:HE22	1.77	0.49
1:A:188:LYS:HB3	1:A:189:PRO:HD3	1.95	0.49
1:A:464:ARG:HG2	1:A:471:HIS:CE1	2.47	0.49
1:A:1000:ARG:HG2	1:A:1076:VAL:HG22	1.95	0.49
1:A:902:ASP:HB3	1:A:906:GLY:H	1.77	0.49
1:A:591:LYS:HB3	1:A:596:GLY:HA2	1.95	0.49
1:A:418:ILE:HG13	1:A:441:ILE:HD11	1.95	0.48
1:A:306:TRP:HZ2	1:A:322:ILE:HD11	1.77	0.48
1:A:397:LEU:HG	1:A:405:THR:HG22	1.96	0.48
2:B:106:A:H2'	2:B:107:A:C8	2.48	0.48
3:C:-7:DG:N3	3:C:-6:DC:H2'	2.29	0.48
1:A:245:LEU:HB3	1:A:371:LEU:HD13	1.95	0.48
1:A:865:ASN:HA	1:A:868:MSE:HE3	1.95	0.48
2:B:32:C:O2'	2:B:33:C:OP1	2.30	0.48
1:A:444:SER:HA	2:B:29:G:C8	2.49	0.47
1:A:29:VAL:HA	1:A:220:MSE:HE2	1.95	0.47
1:A:125:LEU:HD23	1:A:218:ARG:HG2	1.96	0.47
1:A:746:ARG:HH22	2:B:60:C:P	2.37	0.47
1:A:405:THR:HG23	1:A:421:HIS:NE2	2.28	0.47
1:A:635:ARG:HA	1:A:638:ARG:HD2	1.96	0.47
3:C:-7:DG:C2	3:C:-6:DC:H2'	2.49	0.47
1:A:588:ASP:OD1	1:A:588:ASP:N	2.47	0.47
2:B:87:G:O2'	2:B:88:G:H5"	2.16	0.46
2:B:20:U:H5	2:B:86:U:H2'	1.81	0.46
1:A:174:ARG:NH1	1:A:177:ASP:OD1	2.48	0.46
1:A:8:VAL:HG13	1:A:441:ILE:HG23	1.98	0.46
1:A:79:ARG:NH2	1:A:192:ARG:HG2	2.30	0.46
1:A:12:LEU:HD23	1:A:15:MSE:HE3	1.98	0.45
1:A:15:MSE:HE1	1:A:440:PRO:HD2	1.98	0.45
1:A:679:ASP:HB3	1:A:689:ARG:HH11	1.81	0.45

Continued from previous page...



A + a 1	A 4 a ma 2	Interatomic	Clash
Atom-1	Atom-2	distance (\AA)	overlap (Å)
1:A:453:ARG:NE	1:A:460:ALA:HB1	2.32	0.45
1:A:904:ARG:NH2	2:B:5:A:OP2	2.49	0.45
1:A:32:GLY:HA3	1:A:220:MSE:SE	2.67	0.45
1:A:1009:GLU:HG3	1:A:1011:VAL:HG13	1.99	0.45
1:A:828:ASP:C	1:A:830:ARG:H	2.20	0.45
1:A:853:TYR:CZ	1:A:867:LEU:HD21	2.52	0.45
1:A:914:ARG:HE	1:A:959:ILE:HD13	1.81	0.44
1:A:397:LEU:HD23	1:A:397:LEU:HA	1.74	0.44
1:A:602:PHE:CZ	1:A:986:GLN:HB2	2.53	0.44
1:A:22:LEU:HD13	1:A:500:VAL:HG11	2.00	0.44
1:A:624:GLY:HA3	1:A:754:GLY:O	2.18	0.44
1:A:1028:LYS:NZ	1:A:1040:GLU:OE1	2.48	0.44
1:A:69:LYS:HD3	1:A:96:LEU:HD13	2.00	0.43
1:A:419:ARG:HD3	1:A:436:ASP:OD1	2.19	0.43
3:C:-7:DG:H4'	3:C:-6:DC:OP1	2.17	0.43
1:A:1029:VAL:HG12	1:A:1041:ARG:HB2	2.00	0.43
1:A:108:PRO:HD2	1:A:229:MSE:HE2	2.00	0.43
1:A:142:ALA:HB1	4:D:5:DG:H21	1.83	0.43
3:C:27:DC:H3'	3:C:28:DA:C8	2.53	0.43
1:A:490:HIS:HB3	1:A:492:HIS:CD2	2.54	0.42
1:A:734:ARG:HD2	1:A:771:LEU:HD21	2.00	0.42
1:A:1030:PHE:CE1	1:A:1040:GLU:HG2	2.55	0.42
1:A:765:GLU:OE1	1:A:798:ARG:NH2	2.52	0.42
1:A:832:LYS:NZ	2:B:85:G:H5"	2.35	0.42
2:B:33:C:H5"	2:B:34:A:H5"	2.02	0.41
1:A:449:ASN:HA	1:A:464:ARG:HB2	2.03	0.41
1:A:15:MSE:CE	1:A:440:PRO:HD2	2.50	0.41
1:A:29:VAL:HG21	1:A:392:THR:HB	2.02	0.41
1:A:83:ASN:ND2	1:A:189:PRO:HD2	2.35	0.41
1:A:919:CYS:HB3	1:A:928:PHE:CE1	2.56	0.41
1:A:1:MSE:HE3	1:A:509:GLN:HB2	2.03	0.41
1:A:312:ASP:OD1	1:A:312:ASP:N	2.50	0.41
1:A:503:ASN:ND2	2:B:93:A:O2'	2.53	0.41
1:A:64:THR:HG23	1:A:67:GLU:H	1.86	0.41
1:A:305:LYS:HA	1:A:308:LYS:HE2	2.03	0.41
1:A:395:ASP:HB2	1:A:401:LEU:HB2	2.03	0.41
1:A:780:VAL:O	1:A:783:GLN:HG2	2.21	0.41
2:B:94:G:H2'	2:B:95:A:C8	2.56	0.40
1:A:1012:LEU:HD23	1:A:1012:LEU:HA	1.95	0.40
1:A:751:ASP:OD1	1:A:751:ASP:N	2.55	0.40

Continued from previous page...

There are no symmetry-related clashes.



5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	1077/1130~(95%)	1021 (95%)	56~(5%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percen	tiles
1	А	930/945~(98%)	884 (95%)	46~(5%)	25	56

All (46) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	72	LEU
1	А	98	LEU
1	А	134	VAL
1	А	190	LEU
1	А	251	ARG
1	А	257	PHE
1	А	270	LEU
1	А	315	PHE
1	А	338	LEU
1	А	392	THR
1	А	397	LEU
1	А	436	ASP
1	А	480	LYS



Mol	Chain	Res	Type
1	А	490	HIS
1	А	491	MSE
1	А	517	GLU
1	А	588	ASP
1	А	590	LEU
1	А	595	LYS
1	А	607	ASN
1	А	629	LYS
1	А	649	LEU
1	А	652	LEU
1	А	706	CYS
1	А	711	TRP
1	А	738	ARG
1	А	741	GLU
1	А	786	ARG
1	А	798	ARG
1	А	812	LEU
1	А	853	TYR
1	А	866	GLN
1	А	890	VAL
1	А	893	MSE
1	А	932	LEU
1	А	950	ASP
1	A	956	GLU
1	А	983	ASN
1	А	988	LEU
1	А	994	ILE
1	A	997	ILE
1	А	1000	ARG
1	A	1007	ASP
1	А	1029	VAL
1	А	1032	THR
1	А	1092	ARG

Continued from previous page...

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	А	866	GLN

5.3.3 RNA (i)



Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	В	97/112~(86%)	20 (20%)	4 (4%)

All (20) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	В	5	А
2	В	6	G
2	В	12	А
2	В	13	G
2	В	16	U
2	В	21	С
2	В	29	G
2	В	33	С
2	В	34	А
2	В	36	U
2	В	46	С
2	В	58	G
2	В	60	С
2	В	67	G
2	В	69	U
2	В	70	U
2	В	83	А
2	В	87	G
2	В	93	А
2	В	101	А

All (4) RNA pucker outliers are listed below:

Mol	Chain	\mathbf{Res}	Type
2	В	5	А
2	В	32	С
2	В	82	G
2	В	86	U

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.



5.6 Ligand geometry (i)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	ol Type Chain	Chain	Dec	Pog Tipk	Bond lengths			Bond angles		
INIOI		nes		Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2	
5	SO4	А	1201	1	4,4,4	0.14	0	$6,\!6,\!6$	0.05	0
5	SO4	А	1203	1	4,4,4	0.14	0	$6,\!6,\!6$	0.06	0
5	SO4	А	1202	-	4,4,4	0.14	0	$6,\!6,\!6$	0.05	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	#RSRZ>2	$\mathbf{OWAB}(\mathrm{\AA}^2)$	Q<0.9
1	А	1064/1130~(94%)	0.50	54 (5%) 28 24	47, 85, 148, 190	0
2	В	99/112~(88%)	0.04	0 100 100	55, 93, 202, 220	0
3	С	36/38~(94%)	-0.17	1 (2%) 53 50	58, 73, 149, 159	0
4	D	8/8~(100%)	-0.15	0 100 100	56, 59, 69, 91	0
All	All	1207/1288~(93%)	0.44	55 (4%) 32 29	47, 85, 155, 220	0

All (55) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	А	495	ARG	13.2
1	А	831	GLY	8.8
1	А	1071	ARG	7.8
1	А	492	HIS	5.6
1	А	160	GLU	4.9
1	А	830	ARG	4.1
1	А	317	LEU	3.6
1	А	150	ARG	3.6
1	А	739	SER	3.6
1	А	488	LEU	3.5
1	А	490	HIS	3.5
1	А	705	ILE	3.5
1	А	489	ALA	3.4
1	А	1044	GLY	3.4
1	А	413	GLU	3.3
1	А	494	ARG	3.3
1	А	832	LYS	3.3
1	А	54	ASN	3.3
1	А	589	GLU	3.2
1	А	471	HIS	3.1
1	А	748	TYR	3.1



Mol	Chain	\mathbf{Res}	Type	RSRZ
1	А	162	GLU	3.0
1	А	930	TRP	2.9
1	А	164	GLU	2.8
1	А	463	PHE	2.8
1	А	472	PHE	2.8
3	С	-7	DG	2.8
1	А	328	ARG	2.8
1	А	161	GLU	2.7
1	А	1073	LYS	2.7
1	А	493	ARG	2.6
1	А	0	SER	2.6
1	А	467	GLY	2.5
1	А	591	LYS	2.5
1	А	542	LEU	2.5
1	А	470	GLN	2.4
1	А	525	VAL	2.4
1	А	58	GLU	2.4
1	А	1072	GLU	2.3
1	А	608	ASP	2.3
1	А	922	GLU	2.3
1	А	631	LEU	2.3
1	А	508	VAL	2.3
1	А	946	LEU	2.2
1	А	476	PHE	2.2
1	А	159	TRP	2.2
1	А	152	ARG	2.2
1	А	265	HIS	2.2
1	А	1029	VAL	2.2
1	А	675	GLU	2.2
1	А	148	TRP	2.1
1	А	688	TRP	2.1
1	А	461	LEU	2.0
1	А	651	TYR	2.0
1	А	364	TYR	2.0

Continued from previous page...

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

DB ATA BANK

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(A^2)$	Q<0.9
5	SO4	А	1201	5/5	0.85	0.14	121,138,142,150	0
5	SO4	А	1203	5/5	0.92	0.24	89,93,103,130	0
5	SO4	А	1202	5/5	0.98	0.16	64,72,90,95	0

6.5 Other polymers (i)

There are no such residues in this entry.

