



wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 24, 2023 – 12:57 PM EDT

PDB ID : 5U8Q
Title : Structure of the ectodomain of the human Type 1 insulin-like growth factor receptor in complex with IGF-I
Authors : Lawrence, M.; Xu, Y.
Deposited on : 2016-12-14
Resolution : 3.27 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.27 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1177 (3.32-3.24)
Clashscore	141614	1044 (3.30-3.26)
Ramachandran outliers	138981	1026 (3.30-3.26)
Sidechain outliers	138945	1025 (3.30-3.26)
RSRZ outliers	127900	1141 (3.32-3.24)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	885	<div style="display: flex; align-items: center;"> <div style="width: 7%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 74%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 16%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 9%; height: 10px; background-color: grey;"></div> </div>
2	B	70	<div style="display: flex; align-items: center;"> <div style="width: 4%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 61%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 7%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 31%; height: 10px; background-color: grey;"></div> </div>
3	H	126	<div style="display: flex; align-items: center;"> <div style="width: 3%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 82%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 12%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 6%; height: 10px; background-color: grey;"></div> </div>
4	L	108	<div style="display: flex; align-items: center;"> <div style="width: 4%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 80%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 18%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 2%; height: 10px; background-color: grey;"></div> </div>
5	C	2	<div style="display: flex; align-items: center;"> <div style="width: 50%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 50%; height: 10px; background-color: yellow;"></div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	E	2	 100%
6	D	4	 50% 50%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	NAG	C	2	-	-	-	X

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 8704 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Insulin-like growth factor 1 receptor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	801	6427	4066	1112	1202	47	0	0	0

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	?	-	THR	deletion	UNP P08069
A	?	-	THR	deletion	UNP P08069
A	?	-	MET	deletion	UNP P08069
A	?	-	SER	deletion	UNP P08069
A	?	-	SER	deletion	UNP P08069
A	?	-	ARG	deletion	UNP P08069
A	?	-	SER	deletion	UNP P08069
A	?	-	ARG	deletion	UNP P08069
A	?	-	ASN	deletion	UNP P08069
A	?	-	THR	deletion	UNP P08069
A	?	-	THR	deletion	UNP P08069
A	?	-	ALA	deletion	UNP P08069
A	?	-	ALA	deletion	UNP P08069
A	?	-	ASP	deletion	UNP P08069
A	?	-	THR	deletion	UNP P08069
A	?	-	TYR	deletion	UNP P08069
A	?	-	ASN	deletion	UNP P08069
A	?	-	ILE	deletion	UNP P08069
A	?	-	THR	deletion	UNP P08069
A	?	-	ASP	deletion	UNP P08069
A	738	ALA	PRO	conflict	UNP P08069
A	739	GLY	GLU	conflict	UNP P08069
A	740	ASN	GLU	conflict	UNP P08069
A	741	ASN	LEU	conflict	UNP P08069

- Molecule 2 is a protein called Insulin-like growth factor I.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	48	375	235	63	70	7	0	0	0

- Molecule 3 is a protein called Fv 24-60 heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	H	118	915	581	150	181	3	0	0	0

- Molecule 4 is a protein called Fv 24-60 light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	L	107	820	511	140	166	3	0	0	0

- Molecule 5 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
5	C	2	28	16	2	10	0	0	0
5	E	2	28	16	2	10	0	0	0

- Molecule 6 is an oligosaccharide called alpha-D-mannopyranose-(1-4)-alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
6	D	4	50	28	2	20	0	0	0

- Molecule 7 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total O S 5 4 1	0	0
7	A	1	Total O S 5 4 1	0	0


- Molecule 8 is D-MALATE (three-letter code: MLT) (formula: $C_4H_6O_5$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	1	Total C O 9 4 5	0	0

- Molecule 9 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula:

C₈H₁₅NO₆).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
9	A	1	Total	C	N	O	0	0
			14	8	1	5		
9	A	1	Total	C	N	O	0	0
			14	8	1	5		
9	A	1	Total	C	N	O	0	0
			14	8	1	5		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Insulin-like growth factor 1 receptor




- Molecule 2: Insulin-like growth factor I



- Molecule 3: Fv 24-60 heavy chain




- Molecule 4: Fv 24-60 light chain



- Molecule 5: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 5: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 6: alpha-D-mannopyranose-(1-4)-alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Deposit
Cell constants a, b, c, α , β , γ	88.69Å 197.66Å 117.65Å 90.00° 90.00° 90.00°	Deposit
Resolution (Å)	22.17 – 3.27 22.17 – 3.27	Deposit EDS
% Data completeness (in resolution range)	98.6 (22.17-3.27) 99.1 (22.17-3.27)	Deposit EDS
R_{merge}	0.19	Deposit
R_{sym}	(Not available)	Deposit
$\langle I/\sigma(I) \rangle$ ¹	1.20 (at 3.30Å)	Xtrige
Refinement program	phenix.refine 1.11.1-2575_1692, PHENIX 1.11.1-2575_1692	Deposit
R, R_{free}	0.260 , 0.303 0.260 , 0.303	Deposit DCC
R_{free} test set	1524 reflections (4.72%)	wwPDB-V
Wilson B-factor (Å ²)	111.0	Xtrige
Anisotropy	0.718	Xtrige
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 97.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.27$	Xtrige
Estimated twinning fraction	No twinning to report.	Xtrige
F_o, F_c correlation	0.92	EDS
Total number of atoms	8704	wwPDB-V
Average B, all atoms (Å ²)	177.0	wwPDB-V

Xtrige's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.55% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MAN, SO4, MLT, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.24	0/6582	0.44	0/8931
2	B	0.24	0/381	0.39	0/511
3	H	0.24	0/936	0.44	0/1270
4	L	0.25	0/838	0.47	0/1136
All	All	0.24	0/8737	0.44	0/11848

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6427	0	6220	81	0
2	B	375	0	347	5	0
3	H	915	0	888	10	0
4	L	820	0	792	13	0
5	C	28	0	25	1	0
5	E	28	0	25	0	0
6	D	50	0	43	0	0
7	A	10	0	0	0	0
8	A	9	0	4	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
9	A	42	0	39	0	0
All	All	8704	0	8383	106	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 106 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:511:GLN:H	1:A:515:GLY:HA3	1.58	0.69
1:A:114:GLU:HG2	1:A:115:LYS:HG3	1.75	0.68
1:A:433:GLU:HA	1:A:436:THR:HB	1.75	0.67
1:A:337:GLY:HA3	1:A:340:ILE:HD11	1.78	0.66
1:A:144:PRO:HG2	1:A:147:GLU:HB2	1.78	0.64

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	789/885 (89%)	723 (92%)	61 (8%)	5 (1%)	25	58
2	B	44/70 (63%)	40 (91%)	4 (9%)	0	100	100
3	H	116/126 (92%)	111 (96%)	5 (4%)	0	100	100
4	L	105/108 (97%)	99 (94%)	6 (6%)	0	100	100
All	All	1054/1189 (89%)	973 (92%)	76 (7%)	5 (0%)	29	62

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	286	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	11	ASN
1	A	379	LEU
1	A	380	GLY
1	A	504	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	715/787 (91%)	706 (99%)	9 (1%)	69	82
2	B	41/57 (72%)	41 (100%)	0	100	100
3	H	100/107 (94%)	100 (100%)	0	100	100
4	L	95/95 (100%)	93 (98%)	2 (2%)	53	75
All	All	951/1046 (91%)	940 (99%)	11 (1%)	71	83

5 of 11 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	624	ASP
1	A	626	TYR
4	L	90	GLN
4	L	29	ILE
1	A	310	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
4	L	90	GLN
4	L	92	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

8 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	NAG	C	1	5,1	14,14,15	0.28	0	17,19,21	0.45	0
5	NAG	C	2	5	14,14,15	0.27	0	17,19,21	0.52	0
6	NAG	D	1	6,1	14,14,15	0.46	0	17,19,21	0.38	0
6	NAG	D	2	6	14,14,15	0.29	0	17,19,21	0.45	0
6	MAN	D	3	6	11,11,12	0.95	0	15,15,17	1.17	2 (13%)
6	MAN	D	4	6	11,11,12	0.72	1 (9%)	15,15,17	1.21	2 (13%)
5	NAG	E	1	5,1	14,14,15	0.28	0	17,19,21	0.44	0
5	NAG	E	2	5	14,14,15	0.23	0	17,19,21	0.43	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	C	1	5,1	-	1/6/23/26	0/1/1/1
5	NAG	C	2	5	-	0/6/23/26	0/1/1/1
6	NAG	D	1	6,1	-	0/6/23/26	0/1/1/1
6	NAG	D	2	6	-	2/6/23/26	0/1/1/1
6	MAN	D	3	6	-	0/2/19/22	0/1/1/1
6	MAN	D	4	6	-	2/2/19/22	0/1/1/1
5	NAG	E	1	5,1	-	2/6/23/26	0/1/1/1
5	NAG	E	2	5	-	1/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	D	4	MAN	C1-C2	2.07	1.56	1.52

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	D	4	MAN	C1-O5-C5	3.57	117.03	112.19
6	D	3	MAN	C1-O5-C5	2.69	115.84	112.19
6	D	3	MAN	O2-C2-C3	-2.25	105.62	110.14
6	D	4	MAN	O2-C2-C3	-2.17	105.78	110.14

There are no chirality outliers.

5 of 8 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	E	1	NAG	O5-C5-C6-O6
5	E	1	NAG	C4-C5-C6-O6
6	D	4	MAN	O5-C5-C6-O6
6	D	4	MAN	C4-C5-C6-O6
6	D	2	NAG	O5-C5-C6-O6


There are no ring outliers.


1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	C	2	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.







5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
9	NAG	A	1004	1	14,14,15	0.25	0	17,19,21	0.54	0
9	NAG	A	1012	1	14,14,15	0.31	0	17,19,21	0.56	0
9	NAG	A	1007	1	14,14,15	0.40	0	17,19,21	0.57	0
7	SO4	A	1001	-	4,4,4	0.14	0	6,6,6	0.05	0
8	MLT	A	1003	-	8,8,8	1.06	0	10,10,10	1.59	1 (10%)
7	SO4	A	1002	-	4,4,4	0.14	0	6,6,6	0.05	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
9	NAG	A	1012	1	-	1/6/23/26	0/1/1/1
9	NAG	A	1004	1	-	1/6/23/26	0/1/1/1
9	NAG	A	1007	1	-	0/6/23/26	0/1/1/1
8	MLT	A	1003	-	-	6/8/8/8	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	A	1003	MLT	O2-C1-C2	3.39	120.17	112.72

There are no chirality outliers.

5 of 8 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
8	A	1003	MLT	C1-C2-C3-C4
8	A	1003	MLT	O3-C2-C3-C4
9	A	1004	NAG	C3-C2-N2-C7
9	A	1012	NAG	C3-C2-N2-C7
8	A	1003	MLT	O2-C1-C2-C3

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	801/885 (90%)	0.42	65 (8%) 12 11	119, 174, 259, 316	0
2	B	48/70 (68%)	0.27	3 (6%) 20 20	140, 171, 205, 214	0
3	H	118/126 (93%)	0.16	4 (3%) 45 43	109, 147, 196, 222	0
4	L	107/108 (99%)	0.23	4 (3%) 41 39	102, 157, 202, 242	0
All	All	1074/1189 (90%)	0.37	76 (7%) 16 16	102, 170, 251, 316	0

The worst 5 of 76 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	171	TYR	7.1
1	A	233	ALA	6.4
1	A	146	LYS	5.8
1	A	799	GLU	5.6
1	A	512	ASP	5.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	NAG	C	2	14/15	0.77	0.47	161,231,239,242	0
6	MAN	D	4	11/12	0.81	0.59	189,225,236,236	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
6	MAN	D	3	11/12	0.83	0.45	219,242,253,261	0
6	NAG	D	1	14/15	0.84	0.18	136,164,184,214	0
5	NAG	C	1	14/15	0.87	0.20	177,211,224,225	0
6	NAG	D	2	14/15	0.87	0.34	168,202,218,239	0
5	NAG	E	1	14/15	0.90	0.27	119,153,169,179	0
5	NAG	E	2	14/15	0.91	0.37	150,177,196,200	0


The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around Chain E:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
9	NAG	A	1007	14/15	0.64	0.33	188,239,290,327	0
7	SO4	A	1002	5/5	0.74	0.36	172,185,208,221	0
9	NAG	A	1012	14/15	0.74	0.39	159,188,214,215	0
9	NAG	A	1004	14/15	0.81	0.61	172,207,225,227	0
7	SO4	A	1001	5/5	0.82	0.27	163,205,216,216	0
8	MLT	A	1003	9/9	0.83	0.31	137,153,160,161	0

6.5 Other polymers [i](#)

There are no such residues in this entry.