

Full wwPDB X-ray Structure Validation Report (i)

Oct 2, 2023 – 11:31 PM EDT

PDB ID	:	6U8D
Title	:	Crystal structure of hepatitis C virus IRES junction IIIabc in complex with
		Fab HCV2
Authors	:	Koirala, D.; Lewicka, A.; Koldobskaya, Y.; Huang, H.; Piccirilli, J.A.
Deposited on	:	2019-09-04
Resolution	:	1.81 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : FAILED	
Xtriage (Phenix) : 1.13	
EDS : FAILED	
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th	2019)
Ideal geometry (proteins) : Engh & Huber (2001)	
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)	
Validation Pipeline (wwPDB-VP) : 2.35.1	

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\hbox{-}RAY\,DIFFRACTION$

The reported resolution of this entry is 1.81 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.



6U8D

2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 5188 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a RNA chain called JIIIabc RNA (68-MER).

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	А	68	Total 1460	C 650	N 270	O 472	Р 68	0	0	0

• Molecule 2 is a protein called Heavy chain of Fab HCV2.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
2	Н	225	Total 1698	C 1076	N 283	O 333	S 6	0	0	0

• Molecule 3 is a protein called Light chain of Fab HCV2.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
3	L	215	Total 1644	C 1025	N 275	O 338	S 6	0	0	0

• Molecule 4 is water.

Mol	Chain	ResiduesAtoms		ZeroOcc	AltConf
4	А	159	Total O 159 159	0	0
4	Н	124	Total O 124 124	0	0
4	L	103	Total O 103 103	0	0

MolProbity and EDS failed to run properly - this section is therefore empty.



3 Data and refinement statistics (i)

Property	Value	Source	
Space group	C 1 2 1	Depositor	
Cell constants	49.70Å 161.22Å 96.98Å	Depositor	
a, b, c, α , β , γ	90.00° 102.66° 90.00°	Depositor	
Resolution (Å)	61.36 - 1.81	Depositor	
% Data completeness	98.7 (61.36-1.81)	Depositor	
(in resolution range)		Depositor	
R_{merge}	0.04	Depositor	
R_{sym}	(Not available)	Depositor	
$< I/\sigma(I) > 1$	$0.71 (at 1.81 \text{\AA})$	Xtriage	
Refinement program	PHENIX (1.14_3260)	Depositor	
R, R_{free}	0.173 , 0.209	Depositor	
Wilson B-factor $(Å^2)$	37.0	Xtriage	
Anisotropy	0.264	Xtriage	
L-test for twinning ²	$< L > = 0.50, < L^2 > = 0.33$	Xtriage	
Estimated twinning fraction	0.011 for h,-k,-h-l	Xtriage	
Total number of atoms	5188	wwPDB-VP	
Average B, all atoms $(Å^2)$	55.0	wwPDB-VP	

EDS failed to run properly - this section is therefore incomplete.

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 5.04% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

4 Model quality (i)

4.1 Standard geometry (i)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts (i)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles (i)

4.3.1 Protein backbone (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA (i)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates (i)

There are no monosaccharides in this entry.

4.6 Ligand geometry (i)

There are no ligands in this entry.

4.7 Other polymers (i)

There are no such residues in this entry.



4.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



5 Fit of model and data (i)

5.1 Protein, DNA and RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.3 Carbohydrates (i)

EDS failed to run properly - this section is therefore empty.

5.4 Ligands (i)

EDS failed to run properly - this section is therefore empty.

5.5 Other polymers (i)

EDS failed to run properly - this section is therefore empty.

